



NATL INST. OF STAND & TECH



A11107 265847













A11103 066665



# NBS SPECIAL PUBLICATION 513

U.S. DEPARTMENT OF COMMERCE / National Bureau of Standards

National Bureau of Standards  
Library, E-01 Admin. Bldg.

OCT 1 1981

191067

QC

100

.457

## Reaction Rate and Photochemical Data for Atmospheric Chemistry - 1977





MAY 24 1978

Not acc

100

17

13

18

2

# Reaction Rate and Photochemical Data for Atmospheric Chemistry - 1977

*Special publication no. 513*

Edited by

Robert F. Hampson, Jr. and David Garvin

National Measurement Laboratory  
National Bureau of Standards  
Washington, D.C. 20234

Sponsored by the  
Department of Transportation  
National Aeronautics & Space Administration  
Office of Standard Reference Data, NBS  
Office of Environmental Measurements, NBS



---

U.S. DEPARTMENT OF COMMERCE, Juanita M. Kreps, Secretary

Dr. Sidney Harman, Under Secretary

Jordan J. Baruch, Assistant Secretary for Science and Technology

NATIONAL BUREAU OF STANDARDS, Ernest Ambler, Director

Issued May 1978

**Library of Congress Cataloging in Publication Data**

Main entry under title:

Reaction rate and photochemical data for atmospheric chemistry, 1977.

(National Bureau of Standards special publication ; 513)

Supt. of Docs. no.: C13.10:513

1. Atmospheric chemistry--Tables. 2. Chemical reaction, Rate of--Tables, etc. 3. Photochemistry--Tables. I. Hampson, Robert F. II. Garvin, David, 1923- III. United States. Dept. of Transportation IV. United States. National Bureau of Standards. V. Series: United States. National Bureau of Standards. Special publication ; 513.

QC100.U57 no. 513 [QC879.6] 602'.1s [551.5'11] 78-606033

**National Bureau of Standards Special Publication 513**

Nat. Bur. Stand. (U.S.) Spec. Publ. 513, 111 pages (May 1978)

CODEN: XNBSAV

U.S. GOVERNMENT PRINTING OFFICE

WASHINGTON: 1978

---

For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402

Stock No. 003-003-01924-1 Price \$2.75

(Add 25 percent additional for other than U.S. mailing).

## FOREWORD

The National Standard Reference Data System was established in 1963 for the purpose of promoting the critical evaluation and dissemination of numerical data of the physical sciences. The program is coordinated by the Office of Standard Reference Data of the National Bureau of Standards, but involves the efforts of many groups in universities, government laboratories, and private industry. The primary aim of the program is to provide compilations of critically evaluated numerical data. These tables are published in the Journal of Physical and Chemical Reference Data, The NSRDS-NBS Publication Series of the National Bureau of Standards, and through other appropriate channels.

The present report consists of tables of data assembled for use in modelling the chemistry of the stratosphere. It represents contributions from the Chemical Kinetics Information Center, other NSRDS data centers, and a number of individual experts. Support for the preparation of those tables has been provided by the Department of Transportation under the High Altitude Pollution Program, by the National Aeronautics and Space Administration under the Upper Atmosphere Research Program, and by the Office of Standard Reference Data, N.B.S. and the Office of Environmental Measurements, N.B.S.

David R. Lide, Jr.  
Chief, Office of Standard  
Reference Data

## TABLE OF CONTENTS

Abstract	1
1. Introduction	1
2. Guide to the Table	3
3. Reaction Index	9
4. Table. Data for chemical reactions and photochemistry of neutral species	16
5. Reference List	67
<u>Appendices:</u>	
1. Chemical Thermodynamic Properties of Selected Molecular Species	90
2. Conversion Tables	104

R. F. Hampson, D. Garvin, Editors

A table of data for gas phase chemical reactions and photochemistry of neutral species is presented. Specifically, it gives preferred values for reaction rate constants, photoabsorption cross sections, and quantum yields of primary photochemical processes and also cites recent experimental work (to December 1977). It is intended to provide the basic physical chemical data needed as input data for calculations modelling atmospheric chemistry. An auxiliary table of thermochemical data for the pertinent chemical species is given in the appendix.

Key words: Air pollution; atmospheric chemistry; chemical kinetics; data evaluation; gas phase; photoabsorption cross section; photochemistry; quantum yield; rate constant.

## 1. Introduction

This technical note consists of a table of data on the kinetics of chemical reactions and the photochemistry of neutral species. It is designed for use in modelling the chemistry of the stratosphere and, to a more limited extent, the polluted troposphere and also the interpretation of laboratory experiments. The table of rate and photochemical data is supplemented by a table of thermochemical data for the species involved.

---

\* This work was supported by the High Altitude Pollution Program of the Federal Aviation Administration, U. S. Department of Transportation, by the Upper Atmospheric Research Program of the National Aeronautics and Space Administration, and by the Office of Standard Reference Data, N.B.S. and the Office of Environmental Measurements, N.B.S.



Earlier versions of this table have been issued as D. Garvin (editor) "Chemical Kinetics Data Survey IV" NBSIR 73-203 (1973); D. Garvin and R. F. Hampson (editors) "Chemical Kinetics Data Survey VII" NBSIR 74-430 (1974); and R. F. Hampson and D. Garvin (editors) "Chemical Kinetic and Photochemical Data for Modelling Atmospheric Chemistry" NBS Technical Note 866 (1975). The table appearing in NBS Technical Note 866 was also published in the U. S. Department of Transportation's Climatic Impact Assessment Program Monograph 1 "The Natural Stratosphere of 1974" E. Reiter, editor (1975). The present table supersedes all earlier versions.

Since the publication of NBS Technical Note 866 in 1975, a major effort in the evaluation of rate and photochemical data was undertaken for the NASA chlorofluoromethane assessment by its Laboratory Measurements Committee of which the present editors were members. The recommendations of this committee covering 104 chemical reactions and 48 photochemical processes are given in NASA Reference Publication 1010 "Chlorofluoromethanes and the Stratosphere" R. D. Hudson, editor, August 1977. These recommendations have been accepted by the editors of this table and are given here as preferred values. These recommendations are indicated either by the entry [ $\dagger$ \*NASA(1977)eval] or by the symbols [ $\dagger$ \*] immediately preceding the author's name in the reference column. The first notation is used to indicate that the committee has synthesized a preferred value from a consideration of all data sets. The latter designation is used when the committee has accepted the results of a particular study or of an existing evaluation.

Recommended values also are given for reactions not considered by the NASA Laboratory Measurements Committee. Some of these are drawn from other modern rate data evaluations, and the sources are indicated in the table. Other recommendations have been prepared for this work.

Rate data evaluation is a small but growing branch of physical chemistry. There are more recommended rate constants than contained in this table, but it does contain most of those which are applicable to atmospheric chemistry. Evaluations and compilations of rate data and on-going programs in these areas have been described in the recent review paper "Evaluation and Compilation of Reaction Rate Data" by R. F. Hampson and D. Garvin, Journal of Physical Chemistry 81, 2317-2319 (1977). A current extensive listing of sources of evaluated rate data, NBS List of Publications 73 "Chemical Kinetics Tables, Data Evaluations and Bibliographies. A Guide to the Literature" is available on request from the NBS Chemical Kinetics Information Center.

Three sets of detailed data sheets have been published by us in the course of preparing these tables. They are R. F. Hampson (editor) "Survey of Photochemical and Rate Data for Twenty-eight Reactions of Interest in Atmospheric Chemistry" J. Phys. Chem. Ref. Data 2, 267 (1973);

R. F. Hampson (editor) "Chemical Kinetics Data Survey VI, NBSIR 73-207 (1973); and D. Garvin (editor) "Chemical Kinetics Data Survey V" NBSIR 73-206 (1973), all recommendations contained in them (and still pertinent) are summarized in the present table.

We have also accepted many recommendations made by R. T. Watson in "Rate Constants for Reactions of  $\text{C}\ell\text{O}_x$  of Atmospheric Interest"

J. Phys. Chem. Ref. Data 6, 871 (1977) and by D. L. Baulch, et al "Evaluated Kinetic Data for High Temperature Reactions" Volume 1 (1972), Volume 2 (1973) and Volume 3 (1976), Butterworth and Co., London.

In addition to recommendations on rate and photochemical data the table includes listings of current research results. These data listings serve several purposes. Some simply record measurements, usually limited in number, on reactions for which it is not yet practical to give recommended values. Others show the data upon which a new recommendation is based. Still others show what has been done on a reaction since its rate constant was last evaluated. These new data may support the recommendation or suggest the need for modifications in the future.

## 2. Guide to the Table

### 2.1 General

This table provides current (December 1977) information on reaction rate constants, quantum yields and absorption cross sections. For many reactions, preferred values are given. The reactions included in the table are summarized in the index of reactions given in section 3.

Most of the important stratospheric reactions are in the tables. A number of the less important ones and some related systems are included, often simply for comparison.

The following remarks summarize the content of the table. It lists 416 reactions and gives a preferred value of the rate constant for 194 reactions; 136 based on recent reviews and 58 based on recent (1975-1977) experimental work. New data entries (1975-1977) are given for 252 reactions, showing substantial activity in the study of gas kinetics.

The following principles have been used in the selection and presentation of data:

- a. Where a preferred value is given for a rate constant or photochemical quantity, it is so indicated by an asterisk [\*] placed ahead of the entry in the reference column.
- b. As indicated earlier an asterisk preceded by a dagger symbol [†\*] is used to indicate those preferred values that have been recommended by the NASA Laboratory Measurements Committee and published in NASA Reference Publication 1010 "Chlorofluoromethanes and the Stratosphere" R. D. Hudson, editor, August 1977.



- c. Where there is a recent published evaluation of the data and there are no newer data, the evaluation is adopted and usually marked with an asterisk. The original data covered by the evaluation are not listed separately. However this latter rule has not been applied when the evaluation has appeared within the past year.
- d. Where there is an evaluation but there are new data, both are listed and a preferred value is indicated (marked by an asterisk).
- e. Where desirable, the available recent data sets are listed, and a selection is made or a preferred value is synthesized from them.

## 2.2 Uncertainty in Recommended Value of a Rate Constant

The uncertainty assigned here to the recommended value of a rate constant is given in the column "Notes and reliability of log k". This is an estimate by the evaluator of the absolute accuracy of the preferred value. It is a subjective judgment derived from intercomparison of data sets, consideration of related reactions studied with the same technique, estimates of how well the parameters could have been controlled, and comparison with theory. It means that in the evaluators judgment, the true value will lie within the indicated limits to a high level of confidence (90 to 95 percent).

Usually the uncertainty is indicated by the term  $D$  in the expression:  $\log_{10} k = C \pm D$ . This is equivalent to the statement that  $k$  is uncertain to a factor of  $F$  where  $D = \log_{10} F$ . The statement that  $k$  has the value  $k_0$  and is uncertain to a factor of  $F$  means that  $k_0/F < k < k_0 F$ .

Other forms used to indicate reliability are the following:

$A < k < B$  means  $k$  lies in the range between  $A$  and  $B$   
 $k < (>) B$  means  $B$  is an upper (lower) limit  
 $k \sim B$  means  $B$  is only a rough guide to value of  $k$   
 $k = A \pm B$  alternative form for stating reliability limits


At times a rate constant expression is quoted from a paper with individual uncertainties for the rate parameters as  $k = (A \pm a) \exp(-B \pm b/T)$ . These measures are those provided by the author and often are indications of precision, not overall reliability.

Some notes are in order, regarding the statement of the uncertainty in the recommended value of the rate constant for those recommended values recommended by the NASA Laboratory Measurements Committee. Those recommendations were made for the limited temperature range 200-300 K. The uncertainty is indicated for  $T = 230$  K, a typical midstratospheric temperature and in some cases allowance for extrapolation from higher temperatures was necessary. This is why occasionally the entry consists of the value of a rate constant measured at 298 K, the symbols [ $\dagger^*$ ] immediately ahead of the entry in the reference column to show that this value has been adopted by the Committee for the temperature range 200-300 K, and a statement of the uncertainty in the preferred value at  $T = 230$  K. This stated uncertainty will generally be larger than the uncertainty in the measured value (measured at 298 K) to allow for the additional uncertainty introduced by extrapolation to  $T = 230$  K.

For the recommendations made by the NASA Laboratory Committee the indicated uncertainties in NASA Reference Publication 1010 have been doubled in the present table in order to make the reliability statement analogous to a "95 percent confidence level".

### 2.3 Conventions Concerning Rate Constants

a. General Convention. Almost all of the reactions in the table are elementary processes. For them the rate expression is derivable from a statement of the reaction, e.g.





$$-d[A]/dt = -(1/2) d[B]/dt = d[AB_2]/dt = k[A][B]^2$$

Note that the stoichiometric coefficient for B, i.e. 2, appears in the denominator before B's rate of change (which is equal to  $2k[A][B]^2$ ) and as a power on the right hand side.

Wherever there may be any doubt an explicit rate expression is given.

b. Dissociation and Combination Reactions. Some reactions of these types are not of integral kinetic order over the stratospheric pressure and temperature range. That is, although they require an energy transfer agent, "M," they are in the "pressure fall-off region". For some such reactions we tabulate rate constants that include the effect of the energy transfer agent and give rates for various altitudes. The reactions are written to demphasize "M," e.g.,




The rate expressions to be used do not have the concentration of M in them, e.g., Rate =  $k[\text{NO}_2][\text{NO}_3]$  and  $k[\text{HNO}_3]$  respectively. The units given for the k's, s<sup>-1</sup> for first order, and cm<sup>+3</sup> molecule<sup>-1</sup>s<sup>-1</sup> for second order, are consistent with this convention.

The altitude, temperature and number density regime used in these cases is

altitude/km	temp./K	log[M]/molecule cm <sup>-3</sup>
15	217	18.60
20	217	18.27
25	222	17.93
30	227	17.58
35	237	17.26
40	251	16.92
45	265	16.60

This table follows the U. S. standard atmosphere, 1976.

c. Forward and Reverse Reactions. In some cases there are no data on a reaction of interest but there are data on the "reverse" reaction. Occasionally, for



an evaluation will use the data on the "reverse" reaction together with an equilibrium constant to obtain the other rate constant. Obviously this is an approximation but it often is a useful method of estimating non-measured physical properties.

The table contains notations to warn the reader when this procedure has been used, such as "based on reverse reaction," or when both reactions  $f$  and  $r$  are listed together " $k_f = k_r K_{eq}$ " or  $k_r = k_f / K_{eq}$ . These expressions, which are those used in the analyses, are based on equating the two rates at equilibrium.

$$R_f = R_r$$

$$k_f[A] = k_r[B]$$

$$K_{eq} = [B]/[A] = k_f/k_r.$$

## 2.4 Convention Concerning Optical Absorption Coefficients

These are reported in the table as "absorption cross sections per molecule, base e". They are defined by the equations.

$$I/I_0 = \exp(-\sigma[N]\ell)$$

$$\sigma = (1/([N]\ell)) \ln(I_0/I)$$

where  $I_0$  and  $I$  are the intensities of incident and transmitted light,  $\sigma$  is the absorption cross section,  $\text{cm}^2\text{molecule}^{-1}$ ,  $[N]$  is the concentration of absorbers,  $\text{molecules cm}^{-3}$ , and  $\ell$  is the path length, cm. Other definitions and units are frequently used. The terms "absorption coefficient" and "extinction coefficient" are common. It is always necessary to know what concentration units, path length units and type of logarithm (base e or base 10) are used in the definition. To convert "cross-sections" to absorption coefficients in  $(\text{atm at } 273 \text{ K})^{-1} \text{ cm}^{-1}$ , base e, multiply by  $2.69 \times 10^{19}$ .

A table of conversion factors is given in the appendix.



### 3. Arrangement of the Table

Data on a reaction appear only once in the table. The normal location for a reaction is determined by its reactants. Each species has been assigned a sequence number (1 to 65) as shown in the index that follows. These sequence numbers are the same as those used in NBS Technical Note 866. New species have been inserted in appropriate positions and assigned sequence numbers such as 34a, 34b, etc. A reaction is filed under the lower numbered species. That is, the reaction of ozone (7) with an oxygen atom (1) is filed under reactions of oxygen atoms. The numbers for the reactants appear at the left margin of the table, preceding the statement of the reaction, e.g. 1,7  $O + O_3 \rightarrow O_2 + O_2$ . These number pairs run in ascending order through the table.

If a reaction is not filed in the location described above, a cross reference is given there. The most common exception is the listing together of data on the forward and reverse reactions of a pair, i.e.  $A \rightarrow B$ ,  $B \rightarrow A$ .

In the index that follows, bimolecular reactions are listed under both reactants. Frequently the listing of reaction partners for a particular species is divided into two parts by three dashes, separating species earlier in the list than the species indexed from those later in the list. Reactions of the particular species with those species listed before the three dashes should be sought under those reaction partners. There is no indexing of products of reactions. A few species are listed in the index for which there are no reactions in the table, in anticipation of expansion of the data set.

# REACTION INDEX

1. O                    Rxn with: O, O(<sup>1</sup>S), O<sub>2</sub>, O<sub>3</sub>, N, NO, NO<sub>2</sub>, NO<sub>3</sub>, N<sub>2</sub>, N<sub>2</sub>O, N<sub>2</sub>O<sub>5</sub>, NH<sub>2</sub>, NH<sub>3</sub>, H, HO, HO<sub>2</sub>, H<sub>2</sub>, H<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub>, HNO<sub>2</sub>, HNO<sub>3</sub>, SO, SO<sub>2</sub>, SO<sub>3</sub>, HS, H<sub>2</sub>S, CS, CS<sub>2</sub>, OCS, ClO, BrO, OClO, HCl, HBr, NO<sub>3</sub>Cl, Cl<sub>2</sub>, Cl<sub>2</sub>O, CO, CO<sub>2</sub>, CN, CHO, CH<sub>2</sub>O, CH<sub>3</sub>, CH<sub>3</sub>NO<sub>2</sub>, CH<sub>3</sub>ONO, CH<sub>4</sub>, C<sub>2</sub>H<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, C<sub>3</sub>H<sub>6</sub>, alkane, C<sub>6</sub>H<sub>6</sub>, C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>, CH<sub>3</sub>Cl
  
2. O(<sup>1</sup>D)               Rxn with: O<sub>2</sub>, O<sub>3</sub>, NO, NO<sub>2</sub>, N<sub>2</sub>, N<sub>2</sub>O, NH<sub>3</sub>, H<sub>2</sub>, H<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub>, HCl, CO<sub>2</sub>, CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, CF<sub>2</sub>Cl<sub>2</sub>, CFC1<sub>3</sub>
  
3. O(<sup>1</sup>S)               Rxn with: O, O<sub>2</sub>, O<sub>3</sub>, NO, NO<sub>2</sub>, N<sub>2</sub>, N<sub>2</sub>O, NH<sub>3</sub>, H<sub>2</sub>O, CO<sub>2</sub>, CH<sub>4</sub>
  
4. O<sub>2</sub>                   Rxn with: O, O(<sup>1</sup>D), O(<sup>1</sup>S) - - -  
                           hv, N, NO, NO<sub>2</sub>, NO<sub>3</sub>, N<sub>2</sub>, H, H<sub>2</sub>, H<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub>, HNO, SO, HS, Cl, CN, CHO, CH<sub>3</sub>, CH<sub>3</sub>O
  
5. O<sub>2</sub>(<sup>1</sup>Δ)             Rxn with: O<sub>2</sub>, O<sub>3</sub>, N, NO, N<sub>2</sub>, H, SO, SO<sub>2</sub>, H<sub>2</sub>S, CO, CF<sub>2</sub>Cl<sub>2</sub>, CCl<sub>4</sub>, CH<sub>3</sub>Cl, CH<sub>2</sub>Cl<sub>2</sub>, CHCl<sub>3</sub>
  
6. O<sub>2</sub>(<sup>1</sup>Σ)             Rxn with: O<sub>2</sub>, N<sub>2</sub>, H<sub>2</sub>O
  
7. O<sub>3</sub>                   Rxn with: O, O(<sup>1</sup>D), O(<sup>1</sup>S), O<sub>2</sub>(<sup>1</sup>Δ) - - -  
                           hv, M, N, NO, NO<sub>2</sub>, H, HO, HO<sub>2</sub>, SO, SO<sub>2</sub>, H<sub>2</sub>S, Cl, Br, ClO, BrO, CO, CH<sub>2</sub>O, CH<sub>3</sub>, CH<sub>3</sub>O<sub>2</sub>, CH<sub>3</sub>ONO, CH<sub>4</sub>, C<sub>2</sub>H<sub>4</sub>, C<sub>3</sub>H<sub>6</sub>, allene, butenes, butadiene, C<sub>2</sub>Cl<sub>2</sub>H<sub>2</sub>

8. N                    Rxn with: O, O<sub>2</sub>, O<sub>2</sub>(<sup>1</sup>Δ), O<sub>3</sub> - - -  
N, NO, NO<sub>2</sub>, HO, SO, SO<sub>3</sub>, OC1O
  
9. NO                   Rxn with: O, O(<sup>1</sup>D), O(<sup>1</sup>S), O<sub>2</sub>, O<sub>2</sub>(<sup>1</sup>Δ), O<sub>3</sub>, N - - -  
hv, M, NO, NO<sub>2</sub> + H<sub>2</sub>O, NO<sub>3</sub>, NH, NH<sub>2</sub>, H, HO,  
HO<sub>2</sub>, H<sub>2</sub>, H<sub>2</sub>O, H<sub>2</sub>O<sub>2</sub>, Cl, ClO, BrO, OC1O,  
CH<sub>3</sub>, CH<sub>3</sub>O, CH<sub>3</sub>O<sub>2</sub>
  
10. NO<sub>2</sub>                Rxn with: O, O(<sup>1</sup>D), O(<sup>1</sup>S), O<sub>2</sub>, O<sub>3</sub>, N, NO + H<sub>2</sub>O - - -  
hv, M, NO<sub>2</sub>, NO<sub>3</sub>, NH<sub>2</sub>, NH<sub>3</sub>, H, HO, HO<sub>2</sub>,  
SO<sub>2</sub>, Cl, ClO, CH<sub>3</sub>, CH<sub>3</sub>O, CH<sub>3</sub>O<sub>2</sub>
  
11. NO<sub>3</sub>                Rxn with: O, O<sub>2</sub>, NO, NO<sub>2</sub> - - -  
hv, M, NO<sub>3</sub>, H<sub>2</sub>O, SO<sub>2</sub>
  
12. N<sub>2</sub>                   Rxn with: O, O(<sup>1</sup>D), O(<sup>1</sup>S), O<sub>2</sub>, O<sub>2</sub>(<sup>1</sup>Δ), O<sub>2</sub>(<sup>1</sup>Σ) - - -  
M, HO
  
13. N<sub>2</sub>O                Rxn with: O, O(<sup>1</sup>D), O(<sup>1</sup>S) - - -  
hv, M, H, HO, ClO, CO
  
14. N<sub>2</sub>O<sub>5</sub>              Rxn with: O - - - hv, M, H<sub>2</sub>O, SO<sub>2</sub>
  
15. NH                   Rxn with: NO - - -
  
16. NH<sub>2</sub>                Rxn with: O, NO - - - H, HO, H<sub>2</sub>, H<sub>2</sub>O
  
17. NH<sub>3</sub>                Rxn with: O, O(<sup>1</sup>D), O(<sup>1</sup>S), NO<sub>2</sub> - - -  
M, H, HO, ClO



- 17a.  $N_2H_4$       Rxn with: H
18. H      Rxn with:  $O$ ,  $O_2$ ,  $O_2(^1\Delta)$ ,  $O_3$ , NO,  $NO_2$ ,  $N_2O$ ,  $NH_2$ ,  $NH_3$ ,  
 $N_2H_4$  - - -  
H, HO,  $HO_2$ ,  $H_2O$ ,  $H_2O_2$ , HNO,  $HNO_2$ ,  $HNO_3$ ,  
 $SO_2$ , HS,  $H_2S$ , COS, OClO, HCl, NOCl,  $Cl_2$ , CO,  
 $CO_2$ ,  $CH_2O$ ,  $CH_3OOH$ ,  $CH_3ONO$ ,  $CH_3Cl$
19. HO      Rxn with:  $O$ ,  $O_3$ , N, NO,  $NO_2$ ,  $N_2$ ,  $N_2O$ ,  $NH_2$ ,  $NH_3$ , H - - -  
M, HO,  $HO_2$ ,  $H_2$ ,  $D_2$ ,  $H_2O$ ,  $H_2O_2$ , HNO,  $HNO_2$ ,  
 $HNO_3$ ,  $SO_2$ ,  $H_2S$ , Cl, HCl, DCl, HBr,  $NO_3Cl$ ,  
CO,  $CH_2O$ ,  $CH_3OH$ ,  $CH_3NO_2$ ,  $CH_3ONO$ ,  $CH_4$ ,  $C_2H_2$ ,  
 $C_2H_4$ ,  $C_2H_6$ ,  $C_3H_6$ ,  $C_4H_{10}$ , alkane,  $C_6H_6$ ,  
 $C_6H_5CH_3$ ,  $CF_2Cl_2$ ,  $CFCl_3$ ,  $CCl_4$ ,  $CH_3Cl$ ,  
 $CH_2Cl_2$ ,  $CHCl_3$ ,  $CH_iCl_jF_k$
20.  $HO_2$       Rxn with:  $O$ ,  $O_3$ , NO,  $NO_2$ , H, HO - - -  
M,  $HO_2$ ,  $H_2$ ,  $H_2O$ ,  $SO_2$ , Cl, Br, ClO, CO,  $CH_2O$ ,  
 $CH_3O_2$ ,  $C_2H_4$ ,  $C_2H_6$ ,  $C_3H_8$ ,  $C_4H_{10}$
21.  $H_2$       Rxn with:  $O$ ,  $O(^1D)$ ,  $O_2$ , NO,  $NH_2$ , HO,  $HO_2$  - - -  
M, Cl, ClO
22.  $H_2O$       Rxn with:  $O$ ,  $O(^1D)$ ,  $O(^1S)$ ,  $O_2$ ,  $O_2(^1\Sigma)$ , NO,  $NO_3$ ,  $N_2O_5$ ,  
 $NH_2$ , H, HO,  $HO_2$  - - -  $SO_3$
23.  $H_2O_2$       Rxn with:  $O$ ,  $O(^1D)$ ,  $O_2$ , NO, H, HO - - -  
 $h\nu$ , M, Cl, Br

24.	HNO	Rxn with: O <sub>2</sub> , H, HO - - - M, HNO
25.	HNO <sub>2</sub>	Rxn with: O, H, HO - - - hv
26.	HNO <sub>3</sub>	Rxn with: O, H, HO - - - hv, M, Cl
26a.	S	Rxn with: CS <sub>2</sub> , COS
27.	SO	Rxn with: O, O <sub>2</sub> , O <sub>2</sub> ( <sup>1</sup> Δ), O <sub>3</sub> , N - - - SO, SO <sub>3</sub>
28.	SO <sub>2</sub>	Rxn with: O, O <sub>2</sub> ( <sup>1</sup> Δ), O <sub>3</sub> , NO <sub>2</sub> , NO <sub>3</sub> , N <sub>2</sub> O <sub>5</sub> , H, HO, HO <sub>2</sub> - - - CH <sub>3</sub>
29.	SO <sub>3</sub>	Rxn with: O, N, H <sub>2</sub> O, SO - - -
30.	S <sub>2</sub> O	Rxn with: .
31.	HS	Rxn with: O, O <sub>2</sub> , H - - - HS
32.	H <sub>2</sub> S	Rxn with: O, O <sub>2</sub> ( <sup>1</sup> Δ), O <sub>3</sub> , H, HO - - -
33.	HSO <sub>3</sub>	Rxn with:
34.	H <sub>2</sub> SO <sub>4</sub>	Rxn with:
34a.	CS	Rxn with: O - - -
34b.	CS <sub>2</sub>	Rxn with: O, S - - -
34c.	COS	Rxn with: O, H, S - - -

35. Cl Rxn with:  $O_2$ ,  $O_3$ , NO,  $NO_2$ , HO,  $HO_2$ ,  $H_2$ ,  $H_2O_2$ ,  
 $HNO_3$  - - - Cl,  $ClO$ ,  $ClOO$ ,  $NOCl$ ,  $NO_2Cl$ ,  
 $NO_3Cl$ ,  $Cl_2O$ ,  $CH_4$ ,  $C_2H_6$
- Br Rxn with:  $O_3$ ,  $HO_2$ ,  $H_2O_2$  - - -
36.  $ClO$  Rxn with:  $O$ ,  $O_3$ , NO,  $NO_2$ ,  $N_2O$ ,  $NH_3$ ,  $HO_2$ ,  $H_2$  - - -  
 $h\nu$ ,  $ClO$ ,  $BrO$ , CO,  $CH_4$ ,  $C_2H_2$ ,  $C_2H_4$
- BrO Rxn with:  $O$ ,  $O_3$ , NO,  $ClO$  - - - BrO
37.  $ClOO$  Rxn with: Cl - - -  $h\nu$ , M
- $OCIO$  Rxn with:  $O$ , N, NO, H, Cl - - -  $h\nu$
38.  $ClO_3$  Rxn with:
39. HCl Rxn with:  $O$ ,  $O(^1D)$ , H, HO - - -  $h\nu$
- HBr Rxn with:  $O$ , HO - - -
40.  $HOCl$  Rxn with:  $h\nu$
41.  $HOClO$  Rxn with:
42.  $NOCl$  Rxn with: H, Cl - - -  $h\nu$
43.  $NO_2Cl$  Rxn with: Cl - - -  $h\nu$
- 43a.  $NO_3Cl$  Rxn with:  $O$ , HO, Cl - - -  $h\nu$
44.  $Cl_2$  Rxn with:  $O$ , H - - -  $h\nu$
- 44a.  $Cl_2O$  Rxn with:  $O$ , Cl - - -

45.	CO	Rxn with: O, O <sub>2</sub> ( <sup>1</sup> Δ), O <sub>3</sub> , N <sub>2</sub> O, H, HO, HO <sub>2</sub> , ClO - - - CH <sub>3</sub> O
46.	CO <sub>2</sub>	Rxn with: O, O( <sup>1</sup> D), O( <sup>1</sup> S), H - - - M
46a.	CN	Rxn with: O, O <sub>2</sub> - - -
47.	CHO	Rxn with: O, O <sub>2</sub> - - -
48.	CH <sub>2</sub> O	Rxn with: O, O <sub>3</sub> , H, HO, HO <sub>2</sub> - - - hν
49.	CH <sub>3</sub>	Rxn with: O, O <sub>2</sub> , O <sub>3</sub> , NO, NO <sub>2</sub> , SO <sub>2</sub> - - -
50.	CH <sub>3</sub> O	Rxn with: O <sub>2</sub> , NO, NO <sub>2</sub> , CO
51.	CH <sub>3</sub> O <sub>2</sub>	Rxn with: O <sub>3</sub> , NO, NO <sub>2</sub> , HO <sub>2</sub> - - - CH <sub>3</sub> O <sub>2</sub>
52.	CH <sub>3</sub> OH	Rxn with: HO - - -
53.	CH <sub>3</sub> OOH	Rxn with: H - - -
54.	CH <sub>3</sub> NO <sub>2</sub>	Rxn with: O, HO - - -
	CH <sub>3</sub> ONO	Rxn with: O, O <sub>3</sub> , H, HO - - - hν
55.	CH <sub>3</sub> NO <sub>3</sub>	Rxn with:
56.	CH <sub>4</sub>	Rxn with: O, O( <sup>1</sup> D), O( <sup>1</sup> S), O <sub>3</sub> , HO, Cl, ClO - - -
56a.	C <sub>2</sub> H <sub>2</sub>	Rxn with: HO, ClO - - -
57.	C <sub>2</sub> H <sub>4</sub>	Rxn with: O, O <sub>3</sub> , HO, HO <sub>2</sub> , ClO - - -

58.  $C_2H_6$  Rxn with:  $O$ ,  $O(^1D)$ ,  $HO$ ,  $HO_2$ ,  $Cl$  - - -
- \*59.  $C_3H_6$  Rxn with:  $O$ ,  $O_3$ ,  $HO$  - - -
- +60.  $C_3H_8$  Rxn with:  $O$ ,  $HO$ ,  $HO_2$  - - -
- ∇60a.  $C_6H_6$  Rxn with:  $O$ ,  $HO$  - - -
61.  $CF_2Cl_2$  Rxn with:  $O(^1D)$ ,  $O_2(^1\Delta)$ ,  $HO$  - - -  $h\nu$
62.  $CFC1_3$  Rxn with:  $O(^1D)$ ,  $HO$  - - -  $h\nu$
63.  $CCl_4$  Rxn with:  $HO$  - - -  $h\nu$
64.  $CH_iCl_j$  Rxn with:  $O$ ,  $O_2(^1\Delta)$ ,  $O_3$ ,  $H$ ,  $HO$  - - -
- √65.  $CH_iCl_jF_k$  Rxn with:  $HO$  - - -

\* and higher alkenes  
 † and higher alkanes  
 ∇ and other aromatics  
 ✓ and other halocarbons

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
1.1M	$\phi + \phi + M \rightarrow \phi_2 + M$ *Campbell, Gray (1973)	298	$4.8 \times 10^{-33}$	M = N <sub>2</sub> ±0.03
		196	$11.1 \times 10^{-33}$	±0.02
	*Johnston (1968) review	1000-8000	$3.80 \times 10^{-30} T^{-1} \exp(-170/T)$	M = O <sub>2</sub> ±0.3
	Taylor (1975) review	2000-10000	$1.7 \times 10^{-32} T^{-1/2}$	M = N <sub>2</sub>
			$2.2 \times 10^{-28} T^{-3/2}$	M = O <sub>2</sub>
			$6.2 \times 10^{-28} T^{-3/2}$	M = O
			$8.3 \times 10^{-33} T^{-1/2}$	M = N, N <sup>+</sup>
			$5.2 \times 10^{-35} \exp(900/T)$	M = Ar ±0.1 at 190K
1.3	Baulch, et al (1976) review	190-4000	see O(1S) + O →	
1.4M	$O(^1P) + \phi_2 + M \rightarrow \phi_3 + M$ (f)			
7,M	$\phi_3 + M \rightarrow \phi + \phi_2 + M$ (r)			
	*This Survey			
		200-346	$k_f = 6.6 \times 10^{-35} \exp(510/T)$	M = Ar ±0.7
			Rel. M eff.: Ar(1.0), N <sub>2</sub> (1.6), O <sub>2</sub> (1.7), H <sub>2</sub> O(15)	
		300	$k_f = 5.5 \times 10^{-34}$	M = N <sub>2</sub>
	Baulch, et al (1976) review	200-1000	$k_r = 4.1 \times 10^{-10} \exp(-11430/T)$	M = Ar
		200-1000	$k_f = 4.6 \times 10^{-35} \exp(1050/T)$	M = O <sub>2</sub> ±0.1
	Johnston (1968) eval.		$k_r = 1.65 \times 10^{-9} \exp(-11400/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	±0.1
			Rel. M eff.: O <sub>3</sub> (1.0), Ar(0.25), O <sub>2</sub> (0.44), N <sub>2</sub> (0.30)	
	Hule, Herron, Davis (1972)	200-346	$k_f = 6.6 \times 10^{-35} \exp(510/T)$	M = Ar
			Rel. M efficiencies: Ar(1.0), He(0.9), N <sub>2</sub> (1.7)	
	Mulcahy, Williams (1968)	213-386	$k_f = 4.7 \times 10^{-35} \exp(840/T)$	M = Ar
			Rel. M eff.: Ar(1.0), He(0.8), CO <sub>2</sub> (3.4), O <sub>2</sub> (1.1)	
	Meaburn, et al (1968)	300	$k_f = 1.0 \times 10^{-33}$	M = CO <sub>2</sub>
			Rel. M eff.: CO <sub>2</sub> (1.0), C <sub>6</sub> H <sub>6</sub> (0.44), N <sub>2</sub> O(0.88)	
	Stuhl, Niki (1971)	300	$k_f = 5.4 \times 10^{-34} \text{ cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$	M = N <sub>2</sub>
			Rel. M efficiencies: N <sub>2</sub> (1.0), O <sub>2</sub> (1.18), C <sub>6</sub> H <sub>6</sub> (1.24)	
	Donovan, Husain, Kirsch (1970)	300	$k_f = 5.0 \times 10^{-34} \text{ cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$	M = Ar
			Rel. M efficiencies: Ar(1.0), Kr(0.98), He(0.92)	
			$k_f = 8 \times 10^{-34}$	M = N <sub>2</sub>
	Hippler, Troe (1971)	300	$k_f = 4.4 \times 10^{-34}$	M = Ar, Rel. eff. Ar(1.0), N <sub>2</sub> (1.6)
	Slanger, Black (1970)	300	$k_f = 1.24 \times 10^{-34}$	M = O <sub>2</sub>
	Francis (1969)	300	$k_f = 2.28 \times 10^{-34}$	M = Ar
	Sauer (1967)	300	$k_f = 5.4 \pm 1.2 \times 10^{-34}$	M = N <sub>2</sub>
	Ball and Larkin (1973)	295	Rel. efficiencies: N <sub>2</sub> (1.0), O <sub>2</sub> (1.09), Ar(0.78)	
			$k_f \sim 8 \times 10^{-34} \text{ cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$	M = N <sub>2</sub> (a)
	Hippler, Schippert, Troe (1975)	300	$k_f(\infty) = 2.8 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	(a,b)

[illegible]





No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of $\log k$
<b>Hippler, Schippert, Troe (1975)</b>				
1,11	$\text{O} + \text{NO}_3 \rightarrow \text{O}_2 + \text{NO}_2$	300	$k_r \sim 8 \times 10^{-42} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ , $M = \text{N}_2$	(a)
1,12	Graham, Johnston (1978)		$k_f = 8.0 \times 10^{-32} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ , $M = \text{N}_2$	(b)
1,12M	$\text{O} + \text{N}_2 \rightarrow \text{N} + \text{NO}$		$k_f(\text{a}) = 2.0 \times 10^{-11} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$	
13,M	$\text{N}_2\text{O} + \text{M} \rightarrow \text{N}_2 + \text{O} + \text{M}$ (r)		(a) No reliable data. Value based on $k_f$	
	Baulch, et al (1973) review		(b) 2d order high pressure limit	
		1300-2500	$k_f = 3.9 \times 10^{-35} \exp(-10400/T) \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$	
			$\text{M} = \text{Ar}$	
			$k_r = 8.3 \times 10^{-10} \exp(-29000/T) \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$	$\pm 0.2$ (a)
		900-2100	$\text{M} = \text{Ar}$	
			$k_f = 5.5 \times 10^{-15} \exp(-11330/T) \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$	$\pm 0.2$ (a,b)
			$k_r = 1.3 \times 10^{-11} \exp(-30000/T)\text{s}^{-1}$	$\pm 0.2$ (c)
			(a) $k_f = k_r K_{\text{eq}}$	
			(b) 2d order high pressure limit	
			(c) 1st order high pressure limit	
	Schofield (1973) review	300-568	$k_f \leq 5 \times 10^{-38} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$	upper limit only
		800-2100	$k_r = 1.4 \times 10^{11} \exp(-30,000/T)\text{s}^{-1}$	1st order limit
1,13	$\text{O} + \text{N}_2\text{O} \rightarrow \text{N}_2 + \text{O}_2$ (1)			
1,13	$\text{O} + \text{N}_2\text{O} \rightarrow \text{NO} + \text{NO}$ (2)			
9,9	$\text{NO} + \text{NO} \rightarrow \text{N}_2\text{O} + \text{O}$ (2r)			
	Baulch, et al (1973) review			
		1200-2000	$k_1 = 1.7 \times 10^{-10} \exp(-14.1 \times 10^3/T)$	$\pm 0.4$ (a)
			$k_2 = 1.7 \times 10^{-10} \exp(-14.1 \times 10^3/T)$	$\pm 0.3$
		1200-2000	$k_{2r} = 2.2 \times 10^{-12} \exp(-32100/T)$	$\pm 0.7$ (b)
			(a) Based on $k_1/k_2 = 1$	
			(b) $d[\text{NO}]/dt = -2k_{2r}[\text{NO}]^2$	
	Taylor (1975) review	2000-10000	$k_{2r} = 0.4T^{5/2} \exp(-43000/T)$	
1,14	$\text{O} + \text{N}_2\text{O}_5 \rightarrow \text{products}$			
	Graham, Johnston (1978)	300	$< 2 \times 10^{-14}$	preliminary
	Davis (1974)	300	$< 2 \times 10^{-13}$	
1,16	$\text{O} + \text{NH}_2 \rightarrow \text{HN} + \text{H}$ (a)			
	$\rightarrow \text{HO} + \text{NH}$ (b)			
	Gehring, et al (1973)	300	$3.5 \times 10^{-12} (k_a + k_b)$	
1,17	$\text{O} + \text{NH}_3 \rightarrow \text{HO} + \text{NH}_2$ (f)			
16,19	$\text{HO} + \text{NH}_2 \rightarrow \text{O} + \text{NH}_3$ (r)			
	*Kurylo, et al (1969)			
	Baulch, et al (1973) review	361-677	$k_f = 6.6 \times 10^{-12} \exp(-3300/T)$	$\pm 0.3$
		300-1000	$k_f = 2.5 \times 10^{-12} \exp(-3020/T)$	$\pm 0.2$
		300-1000	$k_r = k_f/K_{\text{eq}} = 1 \times 10^{-13}$	$\pm 0.2$ (a)

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of log k
1.18M	Albers, et al (1965) Kondratiev (1970) review	300-1000 350-1000	$k_f = 2 \times 10^{-12} \exp(-3000/T)$ $k_r = 1.8 \times 10^{-12} \exp(-2500/T)$ Data of Albers, et al provide an upper limit when extrapolated to 220K (a) $\text{H}\theta + \text{NH}_2 \rightarrow \text{NH} + \text{H}_2\theta$ may be preferred channel.	
1.19 4.18	$\theta + \text{H} + \text{M} \rightarrow \text{H}\theta + \text{M}$ Schofield (1973) review Baulch, et al (1972) review $\theta + \text{H}\theta \rightarrow \text{H} + \theta_2$ (f) $\theta_2 + \text{H} \rightarrow \theta + \text{H}\theta$ (r)	1000-3000	$\sim 2 \times 10^{-32} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ M = Ar No recommendation	$\pm 1$ estimate
1.19M 20.M	†*Wilson (1972) review †Baulch, et al (1972) review Baulch, et al (1972) review $\theta + \text{H}\theta + \text{M} \rightarrow \text{H}\theta_2 + \text{M}$ (f) $\text{H}\theta_2 + \text{M} \rightarrow \theta + \text{H}\theta + \text{M}$ (r)	300-2000 700-2500 300	$k_f = 4.2 \pm 1.7 \times 10^{-11}$ $k_r = 3.7 \times 10^{-10} \exp(-8450/T)$ $k_f = 3.8 \pm 1.7 \times 10^{-11}$	$\pm 0.3$ $\pm 0.1$ $\pm 0.17$
1.20	Baulch, et al (1972) review $\theta + \text{H}\theta_2 \rightarrow \text{H}\theta + \theta_2$ †*Burrows, Harris, Thrush (1977)	293	no recommendation for forward or reverse rxn $3.5 \times 10^{-11}$ (a) Based on value for $k(\theta + \text{H}\theta)$ and $k(\text{H}\theta + \text{H}_2\theta_2)$	$\pm 0.3$ at 230K (a)
1.21 18.19	Lloyd (1974) review $\theta + \text{H}_2 \rightarrow \text{H}\theta + \text{H}$ (f) $\text{H} + \text{H}\theta \rightarrow \theta + \text{H}_2$ (r) Baulch, et al (1972) review	$\sim 300$ 400-2000	$8 \times 10^{-11} \exp(-500/T)$	estimate
1.22 1.22	Dubinsky, McKenney (1975) Schott, et al (1974) Campbell, Handy (1975) $\theta + \text{H}_2\theta \rightarrow \text{H}\theta + \text{H}\theta$ $\theta + \text{H}_2\theta \rightarrow \text{H} + \text{H}\theta_2$	347-742 1400-1900 363-490	$k_f = 3.0 \times 10^{-14} (T) \exp(-4480/T)$ $k_r = k_f/K_{\text{eq}} = 1.4 \times 10^{-14} (T) \exp(-3500/T)$ $k_f = 8.8 \times 10^{-12} \exp(-4200/T)$ $k_f/k_{\text{ref}} = 3.6 \pm 0.7$ Ref rxn is $\theta_2 + \text{H} \rightarrow \theta + \text{H}\theta$ $k_f = 5.1 \times 10^{-11} \exp(-4950/T)$ See reverse reaction	$\pm 0.1$ $\pm 0.15$
1.23	Baulch, et al (1972) review $\theta(3P) + \text{H}_2\theta_2 = \text{H}\theta_2 + \theta\text{H}$ (a) $= \text{H}_2\theta + \theta_2$ (b) †*Davis, Wong, Schiff (1974) Foner, Hudson (1962) Albers, et al (1971)	283-373 300 370-800	no recommendation $k(a+b) = 2.75 \times 10^{-12} \exp(-2135/T)$ $\leq 4 \times 10^{-15}$ $k(a+b) = 4.6 \times 10^{-11} \exp(-3220/T)$ (a) Although there is no evidence, products are most likely $\theta\text{H} + \text{H}\theta_2$ as the channel to produce $\text{H}_2\theta + \theta_2$ requires a complex rearrangement.	$\pm 0.3$ (a)
1.25	$\theta + \text{HN}\theta_2 \rightarrow \text{H}\theta + \text{N}\theta_2$ This survey		No data. Probably faster than $\theta + \text{HN}\theta_3$ , since it is 94 kJ/mol more exothermic.	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of $\log k$
1,26	$\text{O} + \text{HN}\text{O}_3 \rightarrow \text{HO} + \text{NO}_3$ *Chapman, Wayne (1974) Hampson, et al (1973) review Morley, Smith (1972) Baulch, et al (1973) review	300 300 300 300	$< 3 \times 10^{-17}$ $< 1.5 \times 10^{-14}$ $< 1.3 \times 10^{-14}$ $< 10^{-14}$	
1,27	$\text{O} + \text{SO} \rightarrow \text{SO}_2 + \text{hv}$ Baulch, et al (1976) review		no recommendation	approximate upper limit
1,27M 28,M	$\text{O} + \text{SO} + \text{M} \rightarrow \text{SO}_2 + \text{M}$ (f) $\text{SO}_2 + \text{M} \rightarrow \text{O} + \text{SO} + \text{M}$ (r) *Baulch, et al (1976) review	298	$k_f = 1.9 \times 10^{-31} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ $k_r$ no recommendation $k_f = 8.8 \times 10^{-31}$ , $\text{M} = \text{Ar}$ $k_r = 4.2 \times 10^{-10} \exp(-55000/\text{T})$	$\text{M} = \text{Ar}$ $\pm 0.15$ $\pm 0.3$ $\pm 0.3$
1,28 4,27	$\text{O} + \text{SO}_2 \rightarrow \text{O}_2 + \text{SO}$ (f) $\text{O}_2 + \text{SO} \rightarrow \text{O} + \text{SO}_2$ (r) *Baulch, et al (1976) review	440-2100 440-2100 300	$k_f = 2.1 \times 10^{-10}\text{T}^{-0.5} \exp(-9980/\text{T})$ $k_r = 7.5 \times 10^{-13} \exp(-3250/\text{T})$ $k_r = < 8 \times 10^{-17}$	$\pm 0.25$ (a) $\pm 0.25$
1,28M	Breckenridge, Miller (1972) Schofield (1973) review	400-2500	$k_r = 3.0 \times 10^{-13} \exp(-2800/\text{T})$ (a) $k_f = k_r \times K_{\text{eq}}$	$\pm 0.3$
1,29	$\text{O} + \text{SO}_2 + \text{M} \rightarrow \text{SO}_3 + \text{M}$ *Davis (1976) Schofield (1973) review	220-353 250-1000	$3.4 \times 10^{-32} \exp(-1130/\text{T}) \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ $\text{N}_2(1.0)$ , $\text{He}(1.45)$ , $\text{Ar}(1.87)$ , $\text{SO}_2(56)$ $1 \times 10^{-33} \exp(\pm 500/\text{T}) \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ $\text{M} = \text{O}_2$ , $\text{N}_2$ , $\text{Ar}$ , $\text{He}$	$\text{M} = \text{N}_2$ $\pm 0.3$ at 300K $\pm 0.6$ at other temp.
1,29	Atkinson, Pitts (1974) Westenberg, deHaas(1975d) Baulch, et al (1976) review $\text{O} + \text{SO}_3 \rightarrow \text{products}$ Westenberg, deHaas (1975c)	299-392 248-415	$9.2 \times 10^{-32} \exp(-1000/\text{T})$ , $\text{M} = \text{N}_2\text{O}$ $1.07 \times 10^{-31} \exp(-1400/\text{T})$ , $\text{M} = \text{He}$ Pel. eff: $\text{He}(1.0)$ , $\text{N}_2(2.4)$ , $\text{SO}_2(9.5)$ no recommendation	(a)
1,31	Schofield (1973) review Jacob, Winkler (1972) Baulch, et al (1976) review $\text{O} + \text{HS} \rightarrow \text{H} + \text{SO}$ Cupitt, Glass (1975) $\text{O} + \text{H}_2\text{S} \rightarrow \text{HO} + \text{HS}$ *Whytock, et al (1976)	1480-1550 300-500 295 263-495	$1.4 \times 10^{-31} \exp(785/\text{T}) \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ Rel. eff: $\text{He}(1.0)$ , $\text{N}_2(1.4)$ (a) Reaction $\text{O} + \text{SO}_3 + \text{M} \rightarrow \text{products}$ found to be 3rd order up to $P = 7$ Torr; $\text{SO}_4$ not detected $5 \times 10^{-10} \exp(-6000/\text{T})$ $3 \times 10^{-16} \exp(-500/\text{T})$ no recommendation $1.6 \pm 0.5 \times 10^{-10}$ $7.24 \times 10^{-12} \exp(-1660/\text{T})$	uncertain        $\pm 0.1$



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
1,34a	Hollinden, Kurylo, Timmons (1970)	205-300	2.9 x 10 <sup>-13</sup> exp(-750/T)	(a)
	Baulch, et al (1976) review		no recommendation	
	Schofield (1973) review	200-350	6.3 x 10 <sup>-13</sup> exp(-920/T)	±0.3
	δ + CS → Cδ + S		(a) Assumes stoichiometry of 3.5	
1,34b	*Baulch, et al (1976) review	300	2.2 x 10 <sup>-11</sup>	±0.2
	Bida, et al (1976)	300	2.24 x 10 <sup>-11</sup>	
	Slagle, et al (1975)	305	2.06 x 10 <sup>-11</sup>	
	δ + CS <sub>2</sub> → CS + Sδ (a) → δCS + S (b)			
1,34c	*Baulch, et al (1976) review	200-1000	k <sub>a</sub> = 3.7 x 10 <sup>-11</sup> exp(-700/T)	±0.15, T < 360K
	Wei, Timmons (1975)	218-293	k = 2.8 x 10 <sup>-11</sup> exp(-640/T)	
	Slagle, et al (1974)	302	k = 4.0 x 10 <sup>-12</sup>	
	Graham, Gutman (1976)	249-500	k <sub>b</sub> /k = 0.093 k <sub>b</sub> /k decreases from 0.098 at 249K to 0.081 at 500K	
1,36	δ + δCS → Sδ + Cδ			
	Manning, et al (1976)	296	1.10 ± 0.1 x 10 <sup>-14</sup> , P = 50 Torr 1.49 ± 0.1 x 10 <sup>-14</sup> , P = 2 torr	
	Wei, Timmons (1975)	239-404	2.0 x 10 <sup>-11</sup> exp(-2150/T)	
	Klemm, Stief (1974)	263-502	1.65 x 10 <sup>-11</sup> exp(-2165/T)	
1,36Br	Baulch, et al (1976) review	190-1200	2.6 x 10 <sup>-11</sup> exp(-2250/T)	±0.2 T < 600K ±0.5 T < 600K
	δ + Clδ → Cl + δ <sub>2</sub>			
	†NASA (1977) eval	200-300	7.7 x 10 <sup>-11</sup> exp(-130/T)	±0.12
	Clyne, Nip (1976)	220-425	1.07 x 10 <sup>-10</sup> exp(-224/T)	(a)
1,37	Zahniser, Kaufman (1977)	218-295	(a) Selected by Watson (1977) review k/k <sub>ref</sub> = (1.55 ± 0.17)exp(246 ± 30/T) (b) k <sub>ref</sub> = k(Cl + δ <sub>2</sub> )	(b)
	δ + Brδ → Br + δ <sub>2</sub>			
	†NASA (1977) eval	200-300	3 x 10 <sup>-11</sup>	±0.8 at 230K
	δ + δClδ → Clδ + δ <sub>2</sub>			
1,39	†NASA (1977) eval	200-300	2 x 10 <sup>-11</sup> exp(-1100/T)	±0.6 at 230K (a)
	δ + δClδ → Clδ + δ <sub>2</sub>	298	5 x 10 <sup>-13</sup>	±0.2
	†NASA (1977) eval		(a) Based on room temp value of Bemand, et al	
	Bemand, et al (1973)			
19,35	δ + HCl → Hδ + Cl (f)			
	Hδ + Cl → δ + HCl (r)			
	†Watson (1977) eval			
	†NASA (1977) eval			
1,39	Balakhrnin, et al (1971)	200-300	k <sub>f</sub> = 1.14 x 10 <sup>-11</sup> exp(-3370/T)	±0.3 at 230K
	Brown, Smith (1974)	200-300	k <sub>r</sub> = k <sub>f</sub> x K <sub>eq</sub> = 1.0 x 10 <sup>-11</sup> exp(-2970/T)	±0.3 at 230K
	Wong, Belles (1972)	295-371	k <sub>f</sub> = 1.74 ± 0.6 x 10 <sup>-12</sup> exp(-2260/T)	
	δ + HCl(v = 1) → Hδ + Cl (a) → δ + HCl(v = 0) (b)	293-440 356-628	k <sub>f</sub> = 2.5 x 10 <sup>-12</sup> exp(-2970 ± 150/T) k <sub>f</sub> = 1.9 ± 0.3 x 10 <sup>-11</sup> exp (-3580/T)	

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of log k
1,39Br	Arnoldi, Wolfrum (1974) Brown, Glass, Smith (1975)	300 196-400	$3.6 \pm 1.2 \times 10^{-12}$ $6.2 \times 10^{-12} \exp(-500/T)$ (a) Total rate constant for sum of two reaction paths.	(a) (a)
1,43a	$\phi + \text{HBr} \rightarrow \text{H}\phi + \text{Br}$ +NASA (1977) eval Brown, Smith (1975) Takacs, Glass (1973) $\phi + \text{NO}_2\text{Cl} \rightarrow \text{products}$ +NASA (1977) eval Molina, et al (1977) Kurylo (1977) Ravishankara, et al (1977)	200-300 267-430 298 200-300 213-295 225-273 245	$7.6 \times 10^{-12} \exp(-1570/T)$ $4.0 \times 10^{-12} \exp(-1360 \pm 50/T)$ $(4.4 \pm 1.0) \times 10^{-14}$ $3.0 \times 10^{-12} \exp(-808/T)$ $3.4 \times 10^{-12} \exp(-840/T)$ $1.9 \times 10^{-12} \exp(-692/T)$ $2 \times 10^{-13}$ (a) Recommended expression given by Kurylo (1977) based on his work and that of Molina, et al (1977)	$\pm 0.8$ at 230K $\pm 0.15$ (a)
1,44	$\phi + \text{Cl}_2 \rightarrow \text{Cl}\phi + \text{Cl}$ *Clyne, et al (1976)	174-602	$4.2 \times 10^{-12} \exp(-1370/T)$ (a) Recommendation based on this study (299-502K) and other recent work. Also recommended by Watson (1977) review.	(a)
1,44a	$\phi + \text{Cl}_2\phi \rightarrow 2\text{Cl}\phi$ Watson (1977) review	300	$1.4 \times 10^{-11}$	$\pm 0.1$
1,45M 45M	$\phi + \text{C}\phi + \text{M} \rightarrow \text{C}\phi_2 + \text{M}$ (f) $\text{C}\phi_2 + \text{M} \rightarrow \phi + \text{C}\phi + \text{M}$ (r) *Baulch, et al (1976) review	250-500 296	$k_f = 6.5 \times 10^{-33} \exp(-2180/T)$ M = C $\phi$ $k_r = 2.3 \times 10^{-36}$ M = N $_2$ no recommendation for $k_r$	$\pm 0.2$ at 250K inc. to $\pm 0.5$ at 500K
1,46 4,45	$\phi + \text{C}\phi_2 \rightarrow \text{C}\phi + \phi_2$ (f) $\phi_2 + \text{C}\phi \rightarrow \phi + \text{C}\phi_2$ (r) *Baulch, et al (1976) review	1500-3000 1500-3000	$k_f = 2.8 \times 10^{-11} \exp(-26500/T)$ $k_r = 4.2 \times 10^{-12} \exp(-24000/T)$ (a) $k_f = k_r K_{\text{eq}}$	$\pm 0.3$ (a) $\pm 0.3$
1,46a	$\phi + \text{CN} \rightarrow \text{C}\phi + \text{N}$ Schmatjko, Wolfrum (1976)	295	$1.8 \times 10^{-11}$	
1,47	Albers, et al (1975) $\phi + \text{CH}\phi \rightarrow \text{C}\phi_2 + \text{H}$ (a) $\rightarrow \text{C}\phi + \text{H}\phi$ (b) *Washida, et al (1974) $\phi + \text{CH}_2\phi \rightarrow \text{CH}\phi + \phi\text{H}$	275-387 297	$2.0 \times 10^{-11}$ $2.1 \pm 0.4 \times 10^{-10} (k_a + k_b)$	
1,48	+NASA (1977) eval Herron, Hule (1973) review	200-300 300	$2 \times 10^{-11} \exp(-1450/T)$ $1.5 \times 10^{-13}$	$\pm 0.4$ at 230K (a) $\pm 0.15$

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
	Mack, Thrush (1973)	300	1.5 x 10 <sup>-13</sup> (a) "A factor" chosen; E <sub>act</sub> calc to fit k(700K)	±0.05
1.49	φ + CH <sub>3</sub> → CH <sub>2</sub> φ + H (a) → CHφ + Hφ (b) Washida, Bayes (1976) Washida, et al (1973)	259-341 300	k <sub>a</sub> = 1.0 ± 0.2 x 10 <sup>-10</sup> k <sub>b</sub> negligible k <sub>b</sub> /k <sub>a</sub> < 0.05	
	Slagle, Pruss, Gutman (1974)	300	k <sub>a</sub> = 1.85 ± 0.28 x 10 <sup>-10</sup>	
	Morris, Niki (1972)	300	k <sub>a</sub> /k(φ + tetramethylethylene) = 1.5	
	Peeters, Mahnen (1973)	1100-1900	k <sub>a</sub> = 2.2 x 10 <sup>-10</sup> exp(-1000/T)	
	Bowman (1975)	1875-2240	k <sub>a</sub> = 1.7 x 10 <sup>-10</sup>	
1.54	Biordi, et al (1975) φ + CH <sub>3</sub> NO <sub>2</sub> → products	1550-1725	k <sub>a</sub> = 1.7 x 10 <sup>-10</sup>	
1.54	Campbell, Goodman (1975a) φ + CH <sub>3</sub> ONO → products	295	3.3 x 10 <sup>-15</sup>	
1.56	Davidson, Thrush (1975) φ + CH <sub>4</sub> → products	300-410	2.3 x 10 <sup>-11</sup> exp(-2620/T)	
1.57	*Herron, Huie (1973) review φ + C <sub>2</sub> H <sub>4</sub> → CH <sub>3</sub> + HCφ (a) → CH <sub>2</sub> CO + H <sub>2</sub> (b)	350-1000	3.5 x 10 <sup>-11</sup> exp (-4550/T)	±0.11
	*Herron, Huie (1973) evaluation	200-500	5.5 x 10 <sup>-12</sup> exp (-565/T) (k <sub>a</sub> + k <sub>b</sub> )	±0.08
	Pruss, Slagle, Gutman (1974)	300	k <sub>b</sub> = 3.81 ± 0.95 x 10 <sup>-14</sup> k <sub>b</sub> /(k <sub>a</sub> + k <sub>b</sub> ) = 0.05	(a)
	Atkinson, Pitts (1974)	300-392	5.6 x 10 <sup>-12</sup> exp(-640 ± 100/T) (k <sub>a</sub> + k <sub>b</sub> )	
	Singleton, Cvetanovic (1976)	298-486	1.16 x 10 <sup>-11</sup> exp(-845/T)	
			a) Calculated using above recommended value for (k <sub>a</sub> + k <sub>b</sub> )	
1.58	φ + C <sub>2</sub> H <sub>6</sub> → products	300-650	4.1 x 10 <sup>-11</sup> exp (-3200/T)	±0.11
1.59	*Herron, Huie (1973) evaluation φ + C <sub>3</sub> H <sub>6</sub> → products	200-500	4.1 x 10 <sup>-12</sup> exp (-38/T)	
	Atkinson, Pitts (1974)	300-392	3.45 x 10 <sup>-12</sup> exp(θ ± 150/T)	
	Singleton, Cvetanovic (1976)	298-483	1.26 x 10 <sup>-11</sup> exp(-363/T)	
1.60	φ + alkane → Hφ + alkyl radical Herron, Huie (1969)	250-600	k = [0.8 exp(-2900/T) N <sub>p</sub> + 2.2 exp(-2250/T) N <sub>s</sub> + 2.6 exp(-1650/T) N <sub>t</sub> ] x 10 <sup>-11</sup> Where N <sub>p</sub> , N <sub>s</sub> , and N <sub>t</sub> are the number of primary, secondary, and tertiary hydrogen atoms, respectively. Do not use formula for CH <sub>4</sub> .	±0.08
1.60a	φ + C <sub>6</sub> H <sub>6</sub> → products			
	Colussi, et al (1975)	298-462	1.81 x 10 <sup>-11</sup> exp(-2115/T)	
	Atkinson, Pitts (1975a)	300-392	1.84 x 10 <sup>-11</sup> exp(-2000/T)	



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of log k
1,64	$\theta + \text{C}_6\text{H}_5\text{CH}_3 \rightarrow \text{products}$ Colussi, et al (1975) Atkinson, Pitts (1975a) $\theta + \text{CH}_3\text{Cl} \rightarrow \text{H}\theta + \text{CH}_2\text{Cl}$ Westenberg, deHaas (1975a) Barassin, Combourieu (1974)	298-462 300-392  500-1000 298-443	$3.82 \times 10^{-11} \exp(-1940/T)$ $1.36 \times 10^{-11} \exp(-1550/T)$  $5.8 \times 10^{-11} \exp(-4550/T)$ $2.2 \times 10^{-11} \exp(-3450/T)$ (a) Also measured $k(\theta + \text{CH}_3\text{Br})/k(\theta + \text{CH}_3\text{Cl})$ to be unity over same temp. range	(a)
2,4	$\theta(^1\text{D}_2) + \theta_2 \rightarrow \theta_2(^1\Sigma_g^+) + \theta(^3\text{P})$ †Streit, et al (1976) Fletcher, Husain (1976b)	104-354 300	$2.9 \times 10^{-11} \exp(67/T)$ $5.3 \times 10^{-11}$	0.1
2,7	$\theta(^1\text{D}_2) + \theta_3 \rightarrow \theta_2 + \theta_2$ (a) $\rightarrow \theta_2 + 2\theta$ (b)	300	$k_a/k_b \sim 1$	$\pm 0.1$
2,9	†Streit, et al (1976) Heidner, et al (1973) †Cvetanovic (1975) review $\theta(^1\text{D}_2) + \text{N}\theta \rightarrow \text{N}\theta + \theta(^3\text{P})$ This survey Heidner, Husain (1973)	103-393 300 300  300 300	$2.4 \times 10^{-10} (k_a + k_b)$ $2.7 \times 10^{-10} (k_a + k_b)$ $k_a/k_b \sim 1$  $3.5 \times 10^{-11}$ $8.5 \times 10^{-11}$ (a) $0.41 \times k$ of Heidner and Husain; scaled to match other selected $\theta(^1\text{D})$ rates	$\pm 0.1$     $\pm 0.3$ (a)
2,10	$\theta(^1\text{D}_2) + \text{N}\theta_2 \rightarrow \text{N}\theta + \theta_2$ This survey Heidner, Husain (1973)	300 300	$1.0 \times 10^{-10}$ $2.3 \times 10^{-10}$ (a) $0.41 \times k$ of Heidner and Husain; scaled to match other selected $\theta(^1\text{D})$ rates	$\pm 0.2$ (a)
2,12	$\theta(^1\text{D}_2) + \text{N}_2 \rightarrow \text{N}_2 + \theta(^3\text{P})$ †Streit, et al (1976) Heidner, et al (1973) $\theta(^1\text{D}_2) + \text{N}_2 + \text{M} \rightarrow \text{N}_2\theta + \text{M}$ †NASA (1977) eval	104-354 300  300	$2.0 \times 10^{-11} \exp(107/T)$ $6.9 \times 10^{-11}$  $3.5 \times 10^{-37} \text{ cm}^6 \text{molecule}^{-2} \text{s}^{-1}$ (a) Derived from data in Kajimoto and Cvetanovic (1976a) and value of $k(\theta(^1\text{D}) + \text{N}_2)$ this survey	$\pm 0.1$   (a)
2,13	$\theta(^1\text{D}_2) + \text{N}_2\theta \rightarrow \text{N}_2 + \theta_2$ (a) $\rightarrow 2\text{N}\theta$ (b) †Davidson, et al (1977) Heidner, Husain (1973) †Cvetanovic (1975) review	204-359 300 300	$1.1 \times 10^{-10} (k_a + k_b)$ $2.2 \times 10^{-10} (k_a + k_b)$ $k_a/k_b = 1$	0.1
2,17	$\theta(^1\text{D}_2) + \text{NH}_3 \rightarrow \text{NH}_2 + \text{H}\theta$ †Davidson, et al (1977) Fletcher, Husain (1976a)	204-354 300	$2.5 \times 10^{-10}$ $6.3 \times 10^{-10}$	$\pm 0.1$

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Notes and Reliability of $\log k$
2,21	$\phi(^1\text{D}_2) + \text{H}_2 \rightarrow \text{H}\phi + \text{H}$ *Davidson, et al (1977) Heidner, Husain (1973)	204-352 300	$9.9 \times 10^{-11}$ $2.7 \times 10^{-10}$	
2,22	$\phi(^1\text{D}_2) + \text{H}_2\phi \rightarrow 2\text{H}\phi$ *Streit, et al (1976) Heidner, et al (1973)	253-353 300	$2.3 \times 10^{-10}$ $3.0 \times 10^{-10}$	$\pm 0.1$
2,23	$\phi(^1\text{D}_2) + \text{H}_2\phi_2 \rightarrow \text{H}\phi + \text{H}\phi_2$ Fletcher, Husain (1976a)	300	$5.2 \times 10^{-10}$	
2,39	$\phi(^1\text{D}_2) + \text{HCl} \rightarrow \text{products}$ *Davidson, et al (1977)	199-379	$1.4 \times 10^{-10}$	$\pm 0.1$
2,45	$\phi(^1\text{D}_2) + \text{C}\phi \rightarrow \text{C}\phi + \phi(^3\text{P})$ *This survey Heidner, et al (1973)	300 300	$3 \times 10^{-11}$ $7.3 \times 10^{-11}$	$\pm 0.3$ (a)
2,46	$\phi(^1\text{D}_2) + \text{C}\phi_2 \rightarrow \text{C}\phi_2 + \phi(^3\text{P})$ *Streit, et al (1976)		(a) $0.41 \times k$ of Heidner, et al; scaled to match other selected $\phi(^1\text{D})$ rates	
2,56	Fletcher, Husain (1976b) $\phi(^1\text{D}_2) + \text{CH}_4 \rightarrow \text{CH}_3 + \text{H}\phi$ (a) $\rightarrow \text{CH}_2\phi + \text{H}_2$ (b)	139-200 200-354 300	$1.2 \times 10^{-10}$ $6.8 \times 10^{-11} \exp(117/T)$ $1.7 \times 10^{-10}$	$\pm 0.1$ $\pm 0.1$
2,58	*Davidson, et al (1977) Heidner, Husain (1973) †Cvetanovic (1975) review $\phi(^1\text{D}_2) + \text{C}_2\text{H}_6 \rightarrow \text{products}$	198-357 300 300	$1.4 \times 10^{-10} (k_a + k_b)$ $3.1 \times 10^{-10} (k_a + k_b)$ $k_a/k_b = 10$	
	*This survey Fletcher, Husain (1976a)	300 300	$3 \times 10^{-10}$ $7.3 \times 10^{-10}$	$\pm 0.2$ (a)
2,61	$\phi(^1\text{D}_2) + \text{CF}_2\text{Cl}_2 \rightarrow \text{products}$ *NASA (1977) eval Fletcher, Husain (1976b)	300 300	(a) $0.41 \times k$ of Fletcher and Husain; scaled to match other selected $\phi(^1\text{D})$ rates	
2,62	$\phi(^1\text{D}_2) + \text{CFCl}_3 \rightarrow \text{products}$ *NASA (1977) eval Fletcher, Husain (1976b)	300 300	$2 \times 10^{-10}$ $4.8 \times 10^{-10}$	$\pm 0.2$ (a)
			(a) $0.41 \times k$ of Fletcher and Husain; scaled to match other selected $\phi(^1\text{D})$ rates	
3,1	$\phi(^1\text{S}) + \phi(^3\text{P}) \rightarrow ?$ Slanger, Black (1976a)	300 300	$2.3 \times 10^{-10}$ $5.5 \times 10^{-10}$	$\pm 0.2$ (a)
3,4	$\phi(^1\text{S}) + \phi_2 \rightarrow ?$	200-365 200-377	(a) $0.41 \times k$ of Fletcher and Husain; scaled to match other selected $\phi(^1\text{D})$ rates $5 \times 10^{-11} \exp(-305/T)$ $4.3 \times 10^{-12} \exp(-850/T)$	$\pm 0.15$ a

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of Log k
3,7	$\text{O}(^1\text{S}) + \text{O}_3 \rightarrow ?$	300	$5.8 \times 10^{-10}$	$\pm 0.07$ a
3,9	$\text{O}(^1\text{S}) + \text{NO} \rightarrow ?$	200-291	$3.2 \times 10^{-11} (\text{T})^{0.5}$	$\pm 0.05$ a
3,10	$\text{O}(^1\text{S}) + \text{NO}_2 \rightarrow ?$	300	$5 \times 10^{-10}$	$\pm 0.2$ a
3,12	$\text{O}(^1\text{S}) + \text{N}_2 \rightarrow ?$	200-380	$< 5 \times 10^{-17}$	a
3,13	$\text{O}(^1\text{S}) + \text{N}_2\text{O} \rightarrow ?$			
3,17	Slanger, Black (1976b) $\text{O}(^1\text{S}) + \text{NH}_3 \rightarrow ?$	200-368 300	$3.8 \times 10^{-11} \exp(-420/\text{T})$ $5 \times 10^{-10}$	$\pm 0.2$ a
3,22	$\text{O}(^1\text{S}) + \text{H}_2\text{O} \rightarrow ?$			
3,46	Bingham, et al (1976) $\text{O}(^1\text{S}) + \text{CO}_2 \rightarrow ?$	300 200-450	$1.3 \times 10^{-10}$ $3.1 \times 10^{-11} \exp(-1320/\text{T})$	$\pm 0.15$ a
3,56	$\text{O}(^1\text{S}) + \text{CH}_4 \rightarrow ?$	300	$2 \times 10^{-14}$	$\pm 0.4$ a
4,hv	$\text{O}_2 + \text{h}\nu \rightarrow \text{O}(^3\text{P}) + \text{O}(^3\text{P})$			
4,hv	Hampson, et al (1973a) review $\text{O}_2 + \text{h}\nu \rightarrow \text{O}(^3\text{P}) + \text{O}(^1\text{D})$		$\phi = 1$ $175 < \lambda < 200 \text{ nm}$	
4,hv	Hampson, et al (1973a) review $\text{O}_2 + \text{h}\nu \rightarrow 2 \text{ oxygen atoms}$		$\phi = 1$ $133 < \lambda < 175 \text{ nm}$	
4,hv	Hampson, et al (1973a) review $\text{O}_2 + \text{h}\nu$		$\phi = 1$ $106 < \lambda < 200 \text{ nm}$	
4,8	$\text{O}_2 + \text{N} \rightarrow \text{NO} + \text{O}$		Recommended values of absorption cross section $135 < \lambda < 243 \text{ nm}$	
4,9,9	$\text{O}_2 + \text{NO} + \text{NO} \rightarrow \text{NO}_2 + \text{NO}_2$ *Baulch, et al (1973) review Stedman, Niki (1973)	273-660 300	See reverse reaction $-\text{d}[\text{NO}]/\text{dt} = \text{d}[\text{NO}_2]/\text{dt} = 2k_1[\text{NO}]^2[\text{O}_2]$ $3.3 \times 10^{-39} \exp(530/\text{T}) \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ $2.0 \times 10^{-38}$	$\pm 0.2$ $\pm 0.1$
4,9,10	$\text{O}_2 + \text{NO} + \text{NO}_2 \rightarrow \text{NO}_2 + \text{NO}_3$		See reverse reaction	
4,9	$\text{O}_2 + \text{NO} \rightarrow \text{NO}_2 + \text{O}$		See reverse reaction	
4,9M	$\text{O}_2 + \text{NO} + \text{M} \rightarrow \text{NO}_3 + \text{M}$		No recommendation	
4,10	Baulch, et al (1973) review $\text{O}_2 + \text{NO}_2 \rightarrow \text{NO} + \text{O}_3$		See reverse reaction	
4,11	$\text{O}_2 + \text{NO}_3 \rightarrow \text{NO}_2 + \text{O}_3$		See reverse reaction	
4,12	$\text{O}_2 + \text{N}_2 \rightarrow \text{N}_2\text{O} + \text{O}$			
	*Baulch, et al (1973) review	1200-2000	$1.0 \times 10^{-10} \exp(-55.2 \times 10^3/\text{T})$ (a) $k_f = k_r K_{\text{eq}}$ See reverse reaction	$\pm 0.4$ (a)
4,18	$\text{O}_2 + \text{H} \rightarrow \text{HO} + \text{O}$			
4,18M	$\text{O}_2 + \text{H} + \text{M} \rightarrow \text{HO}_2 + \text{M}$ (f)			
20,M	$\text{HO}_2 + \text{M} \rightarrow \text{O}_2 + \text{H} + \text{M}$ (r)			
	†This Survey			
	$k_f = 6.7 \times 10^{-33} \exp(290/\text{T}) \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$	203-404		$\pm 0.7$ (a)
	M = Ar or He, $k(300) = 1.8 \times 10^{-32}$			
	Rel. M efficiencies: Ar(1.0), He(1.0), $\text{N}_2(3.1)$ , $\text{O}_2(3.1)$ , $\text{H}_2\text{O}(25)$			

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>2</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
			(a) This evaluation is based on the 300 K data reviewed by Baulch, et al (1972) and the recent studies by Hikida, et al (1971), Kurylo (1972) and Wong and Davis (1974).	
	Baulch, et al (1972) eval.	300-2000	$k_f = 4.1 \times 10^{-33} \exp(500/T)$ M = Ar $\pm 2$ Rel. M eff.: Ar(1.0), He(1.0), N <sub>2</sub> (1.3), O <sub>2</sub> (1.3), H <sub>2</sub> O(21) $k_r = k_f/K_{eq} = 3.5 \times 10^{-9} \exp(-23000/T)$ M = Ar $\pm 0.2$ $k_f = 2.35 \times 10^{-32}$ M = Ar $1.64 \times 10^{-32}$ M = Ar $0.75 \times 10^{-32}$ M = He Rel. M efficiencies: He(1.0), Ar(0.8) $1.4(\pm 0.14) \times 10^{-23} \exp(+700(\pm 50)/T)$ , M = Ar Rel. M eff.: at 297K: Ar(1.0), He(0.97), H <sub>2</sub> (1.28).	
	Bishop, Dorfman (1970)	300		
	Hikida, Eyre, Dorfman (1971)	300		
	Ahumada, Michael, Osborne (1972)	300		
	Allen, Moortgat (1973)	115-300		
	Kurylo (1972)	203-404	$6.7 \times 10^{-33} \exp(235/T)$ Rel. M eff.: Ar(1.0), He(1.0), N <sub>2</sub> (3.4)	
	Wong, Davis (1974)	220-360	$6.8 \times 10^{-33} \exp(340/T)$ Rel. M efficiencies: Ar(1.0), He(0.93) N <sub>2</sub> (2.8), H <sub>2</sub> (3.0), CH <sub>4</sub> (21.5)	
	Westenberg, deHaas (1972a)	300	$1.9 \times 10^{-32}$ M = Ar or He	(a)
	Slack (1977)	980-1176	$(9.1 \pm 1.6) \times 10^{-33}$ M = N <sub>2</sub>	(b)
		964-1075	$(6.1 \pm 1.1) \times 10^{-23}$ M = Ar	(a)
		200-2000	$1.85 \times 10^{-28} T^{-1.42}$ M = N <sub>2</sub>	(b)
		200-2200	$5.8 \times 10^{-30} T^{-1}$ M = Ar	(b)
			(a) reanalysis of literature	
			(b) recommendation based on this work	
			and data from literature	
			see reverse reaction	
			no recommendation	
			no recommendation	
			no recommendation	
		300	$< 2.1 \times 10^{-20}$ , F/R > 5000K	estimated
			see reverse reaction	
		295	$< 10^{-13}$	
	Cupitt, Glass (1974)			
	O <sub>2</sub> + Cl + M → ClO + M (f)			
	ClO + M → Cl + O <sub>2</sub> + M (r)			
	† NASA (1977) eval	200-300	$k_r = 5.8 \times 10^{-9} \exp(-3580/T)$	(+2, -1) (a)
			(a) $k_r = k_f/K_{eq}$	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of log k
4,46a	†Nicholas, Norrish (1968)	293	$k_f = 1.7 \times 10^{-33} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$	
	Clyne and Coxon (1968)	300	$k_f = < 5.5 \times 10^{-33}$	
4,47	Stedman (1968)	200-300	$k_f = 5.5 \times 10^{-34}$ (b) As quoted by Clyne and Coxon (1968).	(b)
	$\phi_2 + \text{CN} \rightarrow \text{NC}\phi + \phi$	275-398	$5.3 \times 10^{-11} \exp(-500 \pm 170/T)$	
	Albers, et al (1975)	297	$k/k(\phi + \text{CH}\phi) = 2.74 \pm 0.21 \times 10^{-2}$	
4,49	Peeters, Mahnen (1973)	300	yielding $k = 5.7 \pm 1.2 \times 10^{-12}$	estimated
	$\phi_2 + \text{CH}_3 \rightarrow \text{CH}_2\phi + \text{H}\phi$	1400-1800	$1.7 \times 10^{-13}$ $5 \times 10^{-11}$	
	Washida, Bayes (1976)	259-341	$2.9 \times 10^{-13} \exp(-940/T)$	
	Basco, et al (1972)	295	$3 \times 10^{-16}$	estimated (a)
4,49M	$\phi_2 + \text{CH}_3(^4\text{M}) \rightarrow \text{CH}_3\phi_2(^4\text{M})$		a. Based on negative result and sensitivity limit. Other measurements at higher T suggest an appreciable activation energy.	
	Laufer, Bass (1975)	298	$10^{12} \frac{x}{k} \quad \frac{P(\text{N}_2)}{\text{torr}}$ 0.45 0.7 1.2 1.7 5.0 1.9 2.2 5.1 2.6 1.8 6	(a) (b) (b) (c) (b) (b) (c) (b) (c) (b) (c) (b) (c)
	Washida, Bayes (1976)	300	$1.9 \times 10^{-31} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1} \quad \eta = \text{N}_2$	
	Hochanadel, et al (1977)	295	$2.2 \times 10^{-12}$	
	Basco, et al (1972)	295	$5.1 \times 10^{-13}$	
	van den Bergh, Callear (1971)	295	$2.6 \times 10^{-31} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1} \quad \eta = \text{N}_2$ $1.8 \times 10^{-12}$ $6 \times 10^{-31} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1} \quad \eta = \text{C}_3\text{H}_8$ (a) Values are based on $k(\text{CH}_3 + \text{CH}_3) =$ $9.5 \times 10^{-11}$ , given in Bass, Laufer (1973). (b) 2nd order high pressure limit. (c) 3rd order low pressure limit. see $\text{CH}_3\phi + \phi_2$	preliminary preliminary (a) (b) (c)
4,50	$\phi_2 + \text{CH}_3\phi \rightarrow \dots$			
5,M	$\phi_2(^1\Delta) + \text{M} \rightarrow \phi_2 + \text{M}$	285-322	$2.2 \times 10^{-18} (T/300)^{0.8} \quad \eta = \phi_2$	$\pm 0.1$
	†Hampson, et al (1973) review	77	$1.1 \times 10^{-18} \quad \eta = \phi_2$	(a)
	Huestis, et al (1974)		(a) Liquid phase. When combined with gas phase data of Findlay and Snelling (1971a) (summarized in Hampson, et al (1973) $k = 2.2 \times 10^{-18} (T/300)^{0.5}$	



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
	*Hampson, et al (1973) review	300	$< 2 \times 10^{-20}$ M = N <sub>2</sub>	
	Collins, et al (1973)	300	$1.4 \times 10^{-19}$ M = N <sub>2</sub>	±0.0e
	*Penzhorn, et al (1974)	300	$3.9 \times 10^{-20}$ M = SO <sub>2</sub>	
	Penzhorn, et al (1975)	300	$2.1 \times 10^{-19}$ M = H <sub>2</sub> S	
		300	$k \times 10^{18}$ M	
			0.4 CF <sub>2</sub> Cl <sub>2</sub>	
			<0.08 CCl <sub>4</sub>	
			1.01 CH <sub>3</sub> Cl	
			0.87 CH <sub>2</sub> Cl <sub>2</sub>	
			0.92 CHCl <sub>3</sub>	
	Breckenridge, Miller (1972)	300	$< 2 \times 10^{-15}$ M = SO <sub>2</sub>	(a)
	Fisher, McCarty (1966)	300	$< 3 \times 10^{-16}$ M = CO	(a)
			(a) total rate, quenching and reaction	
5,7	$\theta_2(^1\Delta) + \theta_3 \rightarrow 2\theta_2 + \theta$			
	Clark, Jones, Wayne (1970)	195-439	$6.6 \times 10^{-13}$ exp(-1560/T)	±0.24
	Findlay, Snelling (1971)	283-321	$4.5 \times 10^{-11}$ exp(-2830/T)	±0.11
	Becker, et al (1972)	296-360	$6.0 \times 10^{-11}$ exp(-2850/T)	±0.17
	Collins, et al (1973)	300	$4.4 \times 10^{-15}$	±0.11
	Schofield (1972) review	283-321	$4.5 \times 10^{-11}$ exp(-2830/T)	±0.11
	$\theta_2(^1\Delta) + N \rightarrow N\theta + \theta$			probably
5,8	Schmidt, Schiff (1973)	300	$k < k(N + \theta_2 \rightarrow N\theta + \theta)$	
	Westenberg, et al (1970)	195-300	$k < k(N + \theta_2 \rightarrow N\theta + \theta)$	
	Clark, Wayne (1970)	195-431	$k = 2 \times 10^{-14}$ exp(-600/T)	(a)
			(a) probably refers to physical deactivation, not chemical reaction; see above refs	
5,9	$\theta_2(^1\Delta) + N\theta \rightarrow \theta_2 + N\theta(v > 0)$			
	*Becker, et al (1971)	300	$4.5 \times 10^{-17}$	±0.15
	Yaron, et al (1976)	300	$2.5 \times 10^{-17}$	±0.1
	Giachardi, et al (1976)	293	$(4.5 \pm 1) \times 10^{-17}$	
	Øgrýzlo, Thrush (1973)	300	Nθ(v = 4) observed	
	$\theta_2(^1\Delta) + H \rightarrow$ products			
5,18	Schmidt, Schiff (1973)	300	$(2.5 \pm 0.5) \times 10^{-14}$	(a)
			(a) expt could not distinguish between chemical rxn and physical quenching	
5,27	Westenberg, et al (1970)	300	rxn to give Nθ + θ as products not observed	
	$\theta_2(^1\Delta) + SO \rightarrow \theta_2 + SO(^1\Delta)$			
6,M	Breckenridge, Miller (1972)	300	$3.5 \pm 0.36 \times 10^{-13}$	
	$\theta_2(^1\Sigma) + M \rightarrow \theta_2 + M$			
	*Hampson, et al (1973) review	300	$1.5 \times 10^{-16}$ M = O <sub>2</sub>	±0.12
			$2.0 \times 10^{-15}$ M = N <sub>2</sub>	±0.1
			$4 \times 10^{-12}$ M = H <sub>2</sub> O	±0.18



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$		Notes and Reliability of log k
			$\sigma$	$\sigma^2$	
7, hv	$\sigma_3 + hv \rightarrow \sigma(^3P) + \sigma_2(^3\Sigma_g^-)$ *Hampson, et al (1973) review Johnston (1973)		$\sigma = 1, 450 < \lambda < 750 \text{ nm} = \text{Chappuis bands}$ $\sigma = 0, 250 < \lambda < 350 \text{ nm}$ $k = 3.39 \times 10^{-14} \text{ s}^{-1} \text{ (Daylight)}$ (450 < $\lambda$ < 750 nm = Chappuis bands)		
7, hv	$\sigma_3 + hv \rightarrow \sigma(^3P) + \sigma_2(^1\Delta \text{ or } ^1\Sigma)$ *This survey		$\sigma = 0, \lambda < 303 \text{ nm}$ Total quantum yield ( $\sigma^3P + \sigma^1D$ ) equals unity for 303 < $\lambda$ < 350 nm		
7, hv	$\sigma_3 + hv \rightarrow \sigma(^1D) + \sigma_2(^1\Delta)$ *Lin, DeMore (1973)				
7, hv					
7, hv					
7, M	Hampson, et al (1973) review $\sigma_3 + hv \rightarrow \sigma(^1D) + \sigma_2(^1\Sigma_g^+)$ *Hampson, et al (1973) review $\sigma_3 + M \rightarrow \sigma + \sigma_2 + M$ $\sigma_3 + N \rightarrow NO + \sigma_2$ *NASA (1977) eval				
7, 8					
7, 9	Baulch, et al (1973) review $\sigma_3 + NO \rightarrow NO_2 + \sigma_2(f)$ $\sigma_2 + NO_2 \rightarrow NO + \sigma_3(r)$ *NASA (1977) eval	200-300	$2 \times 10^{-11} \exp(-1070/T)$ (a) Accepts room temp. rec. of Baulch "A factor" chosen; $P_{act}$ derived		$\pm 0.3 \text{ at } 230K$
4, 10		300	$5.7 \times 10^{-13}$		$\pm 0.2$
7, 9		200-300			
4, 10					
7, 9	Birks, et al (1976) Hampson, et al (1973) review Stedman, Niki (1973) Baulch, et al (1973) review	203-361 198-330 298 200-350 200-350 298 300 290	$k_f = 2.1 \times 10^{-12} \exp(-1450/T)$ (a) based on results in Birks et al (1976). Lower "A factor" derived from reconsideration of primary data $k_f = 2.34 \pm 0.23 \times 10^{-12} \exp(-1450 \pm 50/T)$ $k_f = 9 \times 10^{-13} \exp(-1200/T) \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ $k_f = 1.73 \pm 0.1 \times 10^{-14}$ $k_f = 1.5 \times 10^{-12} \exp(-1330/T)$ $k_f = k_f/K_{eq} = 2.8 \times 10^{-12} \exp(-25400/T)$ $k_f = 1.41 \times 10^{-14}$ $k_f = 1.81 \pm 0.13 \times 10^{-14}$ $k_f = 1.70 \times 10^{-14}$		$\pm 0.15 \text{ (a)}$ $\pm 0.11$ $\pm 0.2$ $\pm 0.2$ $\pm 0.2$
7, 9	Chormley, et al (1973) Bemand, et al (1974) Becker, Schurath, Seitz (1974)				

No.	Reaction/Reference + = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{-molecule-sec}$	Notes and Reliability of log k
7.9	$\theta_3 \cdot N_{\theta} - Nd_2(^2B_{1,2}) + d_2(1)$ $- Nd_2(^2A_1) + d_2(2)$ $- \theta_3 \cdot N_{\theta}(3)$ Kurylo, et al (1975) Bar-Ziv, Moy, Gordon (1978) Hui, Cool (1978)	153-373 158-437 138-410	<p style="padding-left: 40px;">see discussion</p> <p style="padding-left: 40px;">see discussion</p> $K_1 = 1.0 \times 10^{-12} \exp(-1450/T)$ $K_2 = 3.8 \times 10^{-13} \exp(-520/T)$ $K_3 = 1.1 \times 10^{-13}$ at 333 K and at 143 K with minimum value of $0.62 \times 10^{-13}$ at approx. 230K	
7.10	$d_3 \cdot Nd_2 - Nd_3 \cdot d_2(f)$ $d_2 \cdot Nd_3 - d_3 \cdot Nd_2(r)$ +-NASA (1977) eval	220-340	$K_f = 1.2 \times 10^{-13} \exp(-2450/T)$	+0.1 (a)
7.11	Davis, Prusaczyk, Dwyer, Kim (1974)	260-343	$K_f = 9.76 \pm 0.54 \times 10^{-14} \exp(-2427 \pm 140/T)$	
	Graham, Johnston (1974)	231-298	$K_f = 1.34 \pm 0.11 \times 10^{-13} \exp(-2466 \pm 30/T)$	
	Hule, Herron (1974a)	259-362	$K_f = 1.57 \pm 0.41 \times 10^{-13} \exp(-2509 \pm 76/T)$	
	Baulch, et al (1973) review	286-302	$K_f = 9.8 \times 10^{-12} \exp(-3500/T)$	+0.7 (b)
		300	$K_r = 7 \times 10^{-34}$ based on $K_f$	+0.3
	Wu, Morris, Niki (1973)	299	$K_f = 4.4 \times 10^{-17}$	+0.06 (c)
	Gormley, et al (1973)	298	$K_f = 3.2 \times 10^{-17}$	
	Stedman, Niki (1973)	298	$K_f = 6.5 \pm 0.8 \times 10^{-17}$	+0.07
	Becker, Schurath, Seitz (1974)	289	$K_f = 3.24 \times 10^{-17}$	
	(a) Derived by combining results from the three 1974 temperature dependent studies			
	(b) Based on Johnston, Yost (1949)			
	(c) Revision, based on later experiments, of value reported by Stedman and Niki (1973)			

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of log k
7,18	$\text{O}_3 + \text{H} \rightarrow \text{HO} + \text{O}_2$ *Clyne, Monkhous (1977)	298-638	$1.0 \pm 0.2 \times 10^{-10} \exp(-516 \pm 50/T)$ (a) NASA (1977) evaluation selected preliminary value reported by these authors, $k = 1.2 \times 10^{-10} \exp(-550/T)$ with $\Delta \log k$ increased to $\pm 0.3$ at 230K.	$\pm 0.2$ (a)
	Lee, et al (1978) Phillips, Schiff (1962)	219-360 300	$1.3 \pm 0.3 \times 10^{-10} \exp(-449 \pm 58/T)$ $2.6 \pm 0.5 \times 10^{-11}$ (b) Quoted by Hampson, et al (1973) review and Baulch, et al (1976) review	(h)
7,19	$\text{O}_3 + \text{HO} \rightarrow \text{HO}_2 + \text{O}_2$ +NASA (1977) eval	200-300	$1.5 \times 10^{-12} \exp(-1000/T)$ (a) *Adjusted from recommendation in Hampson et al (1973) review because of error in derivation of $\text{O}_3$ conc. from absorption meas.	$\pm 0.3$ (a)
	Hampson, et al (1973) review Anderson, Kaufman (1973) Baulch, et al (1976) review DeMore (1973) DeMore (1975)	220-450 220-450 300 300	$1.6 \times 10^{-12} \exp(-1000/T)$ $1.3 \times 10^{-12} \exp(-956/T)$ $6.5 \times 10^{-14}$ $8 \times 10^{-14}$	$\pm 0.3$ $\pm 0.7$
	Davis (1976) Kurylo (1973) $\text{O}_3 + \text{HO}(v_1 > 0) \rightarrow \text{products}$ Coltharp, Worley, Potter (1971)	271-233 300 298	$k/k(\text{OH} + \text{O}_3) = 16.8 \exp(-1230/T)$ at 700 torr $\text{CO}_2$ $7.5 \pm 0.3 \times 10^{-14}$ $6.5 \times 10^{-14}$	$\pm 0.06$
7,19	Streit, Johnston (1976)	300	$y_1(\text{HO})$ $\frac{10^{12} \times k(300 \text{ K})}{\text{CWP}}$ $\frac{\text{SI}}{\text{SI}}$ 2 1.9 $\pm$ 1.1 3 2.4 $\pm$ 0.9 4 2.8 $\pm$ 0.8 5 3.4 $\pm$ 0.7 6 5.3 $\pm$ 0.6 7 6.5 $\pm$ 0.5 8 6.7 $\pm$ 0.5 9 7.7 $\pm$ 0.7 11 $\pm$ 0.4	
7,20	$\text{O}_3 + \text{HO}_2 \rightarrow \text{HO} + 2\text{O}_2$ +NASA (1977) eval. Simonaitis, Heicklen (1973b) DeMore, Tschuikow-Roux (1974) Lloyd (1974) review Baulch, et al (1976) review	200-300 225-298 273-232 200-500 300	$7.3 \times 10^{-14} \exp(-1275/T)$ $k/(k_{\text{ref}})^{0.5} = 1.9 \times 10^{-8} \exp(-1000/T)$ $k/(k_{\text{ref}})^{0.5} = 1.1 \times 10^{-7} \exp(-1550 \pm 50/T)$ $1.7 \times 10^{-13} \exp(-1400/T)$ $1.5 \times 10^{-15}$ (a) $k_{\text{ref}} = k(\text{HO}_2 + \text{HO}_2 \rightarrow \text{H}_2\text{O}_2 + \text{O}_2)$ Preferred value averages temp. dependence of these studies and is based on $k_{\text{ref}}$ value given in this table.	$\pm 0.2$ (a) (a) $\pm 0.3$ $\pm 0.2$

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
7,27	$\phi_3 + S\phi \rightarrow \phi_2 + S\phi_2$ Schofield (1973) review	223-303	$2.5 \times 10^{-12} \exp(-1050/T)$ no recommendation	$\pm 0.5$
7,28	Baulch et al (1976) review $\phi_3 + S\phi_2 \rightarrow S\phi_3 + \phi_2$			
7,32	Daubendiek, Calvert (1974) Davis, Prusaczyk, Dwyer, Kim (1974) $\phi_3 + H_2S \rightarrow$ products	300 300	$k < 8 \times 10^{-24}$ $k < 2 \times 10^{-22}$	
7,35	Glavas, Tohy (1975) Becker, Inocencio, Schurath (1975) $\phi_3 + Cl \rightarrow Cl\phi + \phi_2$	298-343 300	$7 \times 10^{-14} \exp(-3400/T)$ $< 2 \times 10^{-20}$	$\pm 0.4$
7,35Br	+Watson (1977) review Clyne, Nip (1976) Kurylo, Braun (1976) Watson, et al (1976) Zahniser, et al (1976) $\phi_3 + Br \rightarrow Br\phi + \phi_2$	205-298 221-629 213-298 220-350 210-360	$2.7 \times 10^{-11} \exp(-257/T)$ $5.18 \times 10^{-11} \exp(-418/T)$ $2.72 \times 10^{-11} \exp(-298/T)$ $3.08 \times 10^{-11} \exp(-250/T)$ $2.17 \times 10^{-11} \exp(-171/T)$	$\pm 0.12$ at 230K
7,36	+Leu, DeMore (1977) $\phi_3 + Cl\phi \rightarrow \phi Cl\phi + \phi_2$ (a) $\rightarrow Cl\phi\phi + \phi_2$ (b) +NASA (1977) eval	224-422 200-300	$3.34 \times 10^{-11} \exp(-978/T)$ $k_a = 1 \times 10^{-12} \exp(-4000/T)$ $k_b = 1 \times 10^{-12} \exp(-4000/T)$ $< 5 \times 10^{-15}$	$\pm 0.2$ at 230K $\pm 0.6, -2.0$ at 230K (a)
7,36Br	Watson (1977) review	300	(a) Based on unpublished 300K results of Lin and DeMore. Branching ratio unknown - assumed equal to unity.	
7,45	$\phi_3 + Br\phi \rightarrow$ products +Clyne, Cruse (1970) $\phi_3 + C\phi \rightarrow C\phi_2 + \phi_2$	293	$k < 1 \times 10^{-14}$	upper limit only
7,48	Arlin, Warneck (1972) $\phi_3 + CH_2\phi \rightarrow$ products	296	$< 4 \times 10^{-25}$	
7,49	Braslavsky, Heicklen (1976) $\phi_3 + CH_3 \rightarrow CH_3\phi + \phi_2$ (a) $\rightarrow CH_2\phi + H\phi_2$ (b) Simonalitis, Heicklen (1975)	300 221-298	$< 2 \times 10^{-24}$ $(k_a + k_b)/k_{ref} = 12 \exp(-525/T)$ (a) $k_{ref}$ is 2nd order high pressure limit for $CH_3 + \phi_2 (+M) \rightarrow CH_3\phi_2 (+M)$	$\pm 0.3$ upper limit (a)
7,51	$\phi_3 + CH_3\phi_2 \rightarrow$ products	300	$< 2.4 \times 10^{-17}$	
7,54	Simonalitis, Heicklen (1975) $\phi_3 + CH_3\phi\phi \rightarrow$	298-325	$6.8 \times 10^{-13} \exp(-5315/T)$	
7,56	Hastie, et al (1976) $\phi_3 + CH_4 \rightarrow$ products Dillemath, et al (1960)	310-340	$2.7 \times 10^{-13} \exp(-7700/T)$	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
7,57	$\text{C}_3 + \text{C}_2\text{H}_4 \rightarrow \text{products}$ *Herron, Huie (1974) Demerjian, et al (1974) review DeMore (1969) Stedman, et al (1973) Davis (1974) Becker, Schurath, Seitz (1974) Japar, Wu, Niki (1974) Toby, et al (1976) $\text{C}_3 + \text{C}_3\text{H}_6 \rightarrow \text{products}$ *Herron, Huie (1974) Demerjian, et al (1974) review DeMore (1969) Stedman, et al (1973) Becker, Schurath, Seitz (1974) Japar, Wu, Niki (1974)	235-362 300 178-233 300 300 280-360 298 303	$9.0 \times 10^{-15} \exp(-2560/T)$ $2.7 \times 10^{-18}$ $3.2 \times 10^{-15} \exp(-2400/T)$ $1.55 \pm 0.15 \times 10^{-18}$ $1.6 \pm 0.2 \times 10^{-18}$ $1.2 \times 10^{-14} \exp(-2490 \pm 100/T)$ $1.9 \pm 0.1 \times 10^{-18}$ $1.7 \pm 0.1 \times 10^{-18}$	$\pm 0.1$
7,59	$\text{C}_3 + \text{allene} \rightarrow \text{products}$ Toby, Toby (1975)	275-362 300 183, 153, 300 300 280-360 298	$6.1 \times 10^{-15} \exp(-1900/T)$ $1.0 \times 10^{-17}$ $1.6 \times 10^{-15} \exp(-1600/T)$ $1.25 \pm 0.1 \times 10^{-17}$ $1.1 \times 10^{-14} \exp(-1970 \pm 100/T)$ $1.30 \times 10^{-17}$ (a) Includes 300 K point of Hanst et al (1958).	$\pm 0.1$ (a)
7,59	$\text{C}_3 + 1\text{-butene} \rightarrow \text{products}$ Huie, Herron (1975) Japar, Wu, Niki (1974) $\text{C}_3 + \text{cis-2-butene} \rightarrow \text{products}$ Huie, Herron (1975) Japar, Wu, Niki (1974) $\text{C}_3 + \text{trans-2-butene} \rightarrow \text{products}$ Huie, Herron (1975) Japar, Wu, Niki (1974) $\text{C}_3 + 1,3\text{-butadiene} \rightarrow \text{products}$ Toby, Toby (1975) Becker, Schurath, Seitz (1974) $\text{C}_3 + \text{cis-C}_2\text{Cl}_2\text{H}_2 \rightarrow \text{products}$ Blume et al (1976) $\text{C}_3 + \text{trans-C}_2\text{Cl}_2\text{H}_2 \rightarrow \text{products}$ Blume et al (1976) $\text{N} + \text{N} + \text{M} \rightarrow \text{N}_2 + \text{M}$ *Baulch, et al (1973) review	225-363 298 225-363 298 225-363 298 273-343 280-360 296 296 100-600	$2.9 \pm 0.2 \times 10^{-15} \exp(-1686 \pm 20/T)$ $1.23 \times 10^{-17}$ $3.1 \pm 0.7 \times 10^{-15} \exp(-956 \pm 54/T)$ $1.61 \times 10^{-16}$ $6.0 \pm 1.0 \times 10^{-15} \exp(-1051 \pm 43/T)$ $2.60 \times 10^{-16}$ $1 \times 10^{-13} \exp(-2900/T)$ $5.4 \times 10^{-14} \exp(-2680/T)$ $3.7 \times 10^{-20}$ $2.5 \times 10^{-19}$ $8.3 \times 10^{-34} \exp(+500/T) \text{cm}^6 \text{molecule}^{-2} \text{s}^{-1}$ $\text{M} = \text{N}_2$ where $-d[\text{N}]/dt = 2k[\text{N}]^2[\text{M}]$ $7.6 \times 10^{-32} \text{T}^{-1/2} \text{M} = \text{N}_2$	$\pm 0.7$ (a) $\pm 0.5$ $\pm 0.2$ (200-600K) <0.2 (T<200K)
7,59	Taylor (1975) review	2000-10000		



No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
8,9 1,12	N + Nθ → N <sub>2</sub> + θ (f) θ + N <sub>2</sub> → N + Nθ (r) +Clyne, McDermid (1975) Baulch, et al (1973) review	298-670 300-5000 2000-5000	3.0 x 10 <sup>-32</sup> T <sup>-1/2</sup> M = θ <sub>2</sub> , θ, Nθ 6.5 x 10 <sup>-27</sup> T <sup>-3/2</sup> M = N k <sub>f</sub> = 8.2 x 10 <sup>-11</sup> exp(-410/T) ±0.2 at 230K k <sub>f</sub> = 2.7 x 10 <sup>-11</sup> ±0.1 (a) k <sub>r</sub> = 1.3 x 10 <sup>-10</sup> exp(-39000/T) ±0.7 (a) Uncertainty increases to ±0.3 for T> 2000. k <sub>f</sub> based on 300 K data and k <sub>r</sub> (T> 2000).	
8,10	Taylor (1975) review N + Nθ <sub>2</sub> → N <sub>2</sub> θ + θ +NASA (1977) eval. Clyne, McDermid (1975)	2000-10000 200-300 298	k <sub>f</sub> = 2.7 x 10 <sup>-11</sup> 2 x 10 <sup>-11</sup> exp(-800/T) ±0.3 at 230K (a,b) 1.4 x 10 <sup>-12</sup> (a) Accepts room temp. result of Clyne + McDermid "A factor" chosen; Fact derived (b) Believed to be sole reaction channel	
8,19	N + Hθ → Nθ + H *Baulch, et al (1973) review	300	5.3 x 10 <sup>-11</sup> (a) Based on k/k(θ + Hθ → θ <sub>2</sub> + H) = 1.4	
8,27	N + Sθ → products Baulch, et al (1976) review		no recommendation	
8,29	N + Sθ <sub>3</sub> → products Baulch, et al (1976) review		no recommendation	
8,37	N + θClθ → Nθ + Clθ Watson (1977) review	298	k < 6 x 10 <sup>-13</sup> preliminary value	
9,hv	Nθ + hv Hampson, et al (1973a) review		Recommended values of oscillator strengths from which absorption cross sections can be derived are given for β, γ, and ε bands. Recommended values of absorption cross section 106 < λ < 135 nm. No recommended quantum yield values. Oscillator strength for δ bands, confirmed by Mandelman and Carrington (1974)	
9,M	*Bethke (1959) Nθ + M → N + θ + M Baulch, et al (1973) review	4200-6700	Insufficient data for a reliable recommendation. Use, with caution: 6.6 x 10 <sup>-4</sup> T <sup>-1.5</sup> exp(-75.5 x 10 <sup>3</sup> /T) cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> (M = Ar, θ <sub>2</sub> , N <sub>2</sub> ) k(M = Nθ, N, θ)/k(Ar) ~ 1:8	
9,9	Nθ + Nθ → N + Nθ <sub>2</sub> Baulch, et al (1973) review		Endothermic. Unimportant compared to Nθ + Nθ → N <sub>2</sub> θ + θ see reverse reaction	
9,9	Nθ + Nθ → N <sub>2</sub> θ + θ			



No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
9,9,4	N $\theta$ + N $\theta$ + $\theta_2$ $\rightarrow$ N $\theta_2$ + N $\theta_2$		see $\theta_2$ + N $\theta$ + N $\theta$ $\rightarrow$ N $\theta_2$ + N $\theta_2$	
9,10,22	N $\theta$ + N $\theta_2$ + H $\theta_2$ $\rightarrow$ 2HN $\theta_2$ (f) 2HN $\theta_2$ $\rightarrow$ N $\theta$ + N $\theta_2$ + H $\theta_2$ (r) *Chan, et al (1976)	296	$k_f = 6.0 \times 10^{-38}$ cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> $k_r = 9.5 \times 10^{-19}$ cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> Reviewed earlier work, probably heterogeneous	
9,11	Hampson, et al (1973) review N $\theta$ + N $\theta_3$ $\rightarrow$ 2N $\theta_2$ Graham, Johnston (1978) Baulch, et al (1973) review Johnston (1966) review	300 300 300 300	(1.9 $\pm$ 0.4) $\times 10^{-11}$ $\sim 2 \times 10^{-11}$ 8.3 $\times 10^{-12}$ Temperature effect is probably zero or very small.	$\pm 0.7$ estimated
9,15	N $\theta$ + NH $\rightarrow$ products Hansen, et al (1976) Gordon, et al (1971) Mulvihill, Phillips (1975)	300 300 1500	4.7 $\times 10^{-11}$ 3.8 $\times 10^{-11}$ N $\theta_2$ + $\theta$ + H identified as products of major rxn channel	$\pm 0.1$
9,16	N $\theta$ + NH $\theta_2$ $\rightarrow$ N $\theta_2$ + H $\theta_2$ * *Hancock, et al (1975) Lesclaux, et al (1975)	298 300	2.1 $\times 10^{-11}$ 1.8 $\times 10^{-11}$ (a) $\gamma^{-1.25}$ dependence observed 300-500K	(a)
9,18M	Gehring, et al (1973) Gordon, et al (1971) N $\theta$ + H + M $\rightarrow$ HN $\theta$ + M *This survey Baulch, et al (1973) review	300 300 220-400 230-700	8 $\times 10^{-12}$ 2.7 $\times 10^{-11}$ 2.1 $\times 10^{-32}$ exp(300/T) 1.5 $\times 10^{-32}$ exp(300/T) cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> M = H $\theta_2$	$\pm 0.1$ $\pm 0.2$
9,19	Atkinson, Cvetanovic (1973) Allen, Moortgat (1973) Thrush (1973) review N $\theta$ + H $\theta$ $\rightarrow$ N $\theta_2$ + H Baulch, et al (1973) review	286-390 180-300 230-700 298-633	2.5 $\times 10^{-32}$ exp(270/T) M = H $\theta_2$ 5.6( $\pm 0.6$ ) $\times 10^{-23}$ exp( $\pm 375(\pm 65)/T$ ), M = Ar 2 $\times 10^{-32}$ exp( $\pm 300/T$ ), M = H $\theta_2$ 5.2 $\times 10^{-12}$ exp( $-15.1 \times 10^3/T$ ) based on k = 5.8 $\times 10^{-10}$ exp( $-740/T$ ) for reverse reaction (a) corrected expression	$\pm 0.2$ (298K) (a) increasing to $\pm 0.3$ (623K)
9,19M	N $\theta$ + H $\theta$ (+M) $\rightarrow$ HN $\theta_2$ (+M) Hampson, et al (1973a) review	217 217 222 227 237 251 265 273-395	Elevation/km 15 20 25 30 35 40 45 2.2 $\times 10^{-32}$ exp( $\pm 1110/T$ ) cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> M=He $\pm 0.2$ at 300K (3rd order low pressure limit) k(T,M) 2.4 $\times 10^{-12}$ Abs. value of log k $\pm 0.5$ 1.5 $\times 10^{-12}$ Relative value of k 8.9 $\times 10^{-13}$ at different altitudes 4.8 $\times 10^{-13}$ reliable to $\pm 20\%$ 2.6 $\times 10^{-13}$ M = N $\theta_2$ 1.1 $\times 10^{-13}$ 5.9 $\times 10^{-14}$	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of log k
9.20		300	$2 \times 10^{-12} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ (2d order high pressure limit) $5.8 \pm 1.2 \times 10^{-31} \times (295/T)^{2.4} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ $M = \text{N}_2$ (3rd order low pressure limit) Relative efficiencies: $\text{N}_2(1.0)$ , $\text{Ar}(0.58)$ , $\text{He}(0.57)$	$\pm 0.7$
	Anderson, et al (1974)	295, 439		
	Howard, Evenson (1974)	296	$7.8 \pm 1.2 \times 10^{-31} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ $M = \text{N}_2$ (3rd order low pressure limit) Relative efficiencies: $\text{N}_2(1.0)$ , $\text{Ar}(0.55)$ , $\text{He}(0.51)$	
	Cox (1974a)	294	$k/k(\text{H}\theta + \text{C}\theta) = 40 \pm 7$ $k = 1.2 \times 10^{-11} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ (a) 2nd order rate constant, $[M] = 1 \text{ atm } \text{N}_2 + \text{O}_2$ ; $k_{\text{ref}} = 3 \times 10^{-13}$ , this survey	(a)
	Harris, Wayne (1975)	300	$7 \pm 2 \times 10^{-31} \text{ cm}^6\text{molecule}^{-2}\text{s}^{-1}$ , $M = \text{Ar}$ $15 \pm 5 \times 10^{-31}$ , $M = \text{N}_2$ (3rd order low pressure limits)	
	Gordon, Mulac (1975)	435	$7.5 \pm 0.3 \times 10^{-12} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ (2nd order rate constant at 1 atm. $\text{H}_2\theta$ vapor)	
	Atkinson, Hansen, Pitts (1975a)	298	$4.94 \pm 0.5 \times 10^{-13}$ $M = 25 \text{ Torr } \text{N}_2$	(b)
	Cox, Derwent, Holt (1976)	298	(b) also data for $M = \text{Ar}$ at 25-655 torr $1.17 \times 10^{-11}$ $M = 1 \text{ atm } \text{N}_2 + \text{O}_2$	(c)
	Overend, et al (1976)	295	(c) relative to $k(\text{H}\theta + \text{H}_2) = 7 \times 10^{-15}$ $0.956 \pm 0.05 \times 10^{-12}$ $M = 25 \text{ torr } \text{N}_2$	
	Sie, et al (1976b)	298	$6.89 \pm 0.68 \times 10^{-12}$ $M = 770 \text{ torr } \text{N}_2$ $k/k_{\text{ref}} = 16.1$ at 95 torr and = 22 at 408 and 768 torr total pressure, mostly $\text{H}_2$ where ref rxn is $\text{H}\theta + \text{C}\theta \rightarrow \text{C}\theta_2 + \text{H}$	$\pm 0.4$ at 230K
9.20M	$\text{N}\theta + \text{H}\theta_2 \rightarrow \text{N}\theta_2 + \text{H}\theta$	296	$(8.1 \pm 1.5) \times 10^{-12}$	
	†Howard, Evenson (1977)	298-670	$2.0 \times 10^{-11} \exp(-1200/T)$	(a)
	Hack, et al (1975)	245-328	$k/(k_{\text{ref}})^{0.5} = 6.4 \times 10^{-6} \exp(-700/T)$	
	Simonaitis, Heicklen (1978)		(a) ref rxn: $2\text{H}\theta_2 \rightarrow \text{H}_2\theta_2 + \text{O}_2$	
	Cox, Derwent (1975)	296	$1.2 \times 10^{-12}$	$\pm 0.5$
9.21	Payne, Stief, Davis (1973)	300	$3 \times 10^{-13}$	$\pm 0.5$
	Glanzer, Troe (1975)	1350-1700	$7.5 \times 10^{-12}$	
	$\text{N}\theta + \text{H}\theta_2(+M) \rightarrow \text{H}\theta_3(+M)$			
	Cox, Derwent (1975)	296	$1.4 \times 10^{-13}$ at 1 atm $\text{N}_2 + \text{O}_2$	
9.21	Simonaitis, Heicklen (1976a)	296	$< 2 \times 10^{-15}$	
	$\text{N}\theta + \text{H}_2 \rightarrow \text{H}\theta + \text{H}$			
	Baulch, et al (1973) review	2000	$5.3 \times 10^{-18} \text{ cm}^3\text{molecule}^{-1}\text{s}^{-1}$ (a) No data. Value based on reverse rate	$\pm 0.2$ (a)

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
9,22	N $\theta$ + H <sub>2</sub> $\theta$ $\rightarrow$ HN $\theta$ + H $\theta$ Baulch, et al (1973) review	2000	4 x 10 <sup>-18</sup> (a) No data. Value based on reverse rate. (corrected)	$\pm 0.2$ (a)
9,23	N $\theta$ + H <sub>2</sub> $\theta_2$ $\rightarrow$ H $\theta$ + HN $\theta_2$ *Hampson, et al (1973) review	300 500 300	< 5 x 10 <sup>-20</sup> ~ 2 x 10 <sup>-20</sup> < 5 x 10 <sup>-20</sup>	
9,35M	Gray, et al (1972) N $\theta$ + Cl + M $\rightarrow$ ClN $\theta$ + M +*NASA (1977) eval	200-300	1.7 x 10 <sup>-32</sup> exp(+530/T) M = N <sub>2</sub> (a) See Watson (1977) review	$\pm 0.6$ at 230K (a)
9,36	N $\theta$ + Cl $\theta$ $\rightarrow$ N $\theta_2$ + Cl +*NASA (1977) eval Watson (1977) review Zahniser, Kaufman (1977)	200-300 220-298 230-258	1.0 x 10 <sup>-11</sup> exp(200 $\pm$ 100/T) 8 x 10 <sup>-12</sup> exp(250/T) k/k <sub>ref</sub> = (0.52 $\pm$ 0.05) exp(373 $\pm$ 23/T) (a) k <sub>ref</sub> = k(Cl + $\theta_3$ )	$\pm 0.3$ at 230K $\pm 0.1$ (a)
9,36Br	N $\theta$ + Br $\theta$ $\rightarrow$ N $\theta_2$ + Br +*NASA (1977) eval Clyne, Watson (1975) N $\theta$ + $\theta$ Cl $\theta$ $\rightarrow$ N $\theta_2$ + Cl $\theta$	200-300 298	2.1 x 10 <sup>-11</sup> 2.2 x 10 <sup>-11</sup>	$\pm 0.4$ at 230K $\pm 0.1$
9,37	+*NASA (1977) eval Bemand, Clyne, Watson (1973)	200-300 300	2.5 x 10 <sup>-12</sup> exp(-600/T) 3.4 x 10 <sup>-13</sup> (a) Based on room temp. value of Bemand, Clyne, Watson; temp dependence estimated	$\pm 0.6$ at 230K (a)
9,49M	N $\theta$ + CH <sub>3</sub> (+M) $\rightarrow$ CH <sub>3</sub> N $\theta$ (+M) Laufer, Bass (1975)	298	$10^{11} \frac{x}{k}$ .3 .45 1.1 4 x 10 <sup>-12</sup> 1.7 x 10 <sup>-11</sup> (a) Values are based on k(CH <sub>3</sub> + CH <sub>3</sub> ) = 9.5 x 10 <sup>-11</sup> given in Bass, Laufer (1973). (b) 2nd order high pressure limit. see CH <sub>3</sub> + N $\theta$ $\rightarrow$ . . .	(a) (b) (b)
9,50	N $\theta$ + CH <sub>3</sub> $\theta$ $\rightarrow$ . . .	298		
9,51	N $\theta$ + CH <sub>3</sub> $\theta_2$ $\rightarrow$ CH <sub>3</sub> $\theta_2$ N $\theta$ (a) $\rightarrow$ CH <sub>2</sub> $\theta$ + HN $\theta$ (b) $\rightarrow$ CH <sub>3</sub> $\theta$ + N $\theta_2$ (c) +Demerjian, et al (1974) review Cox, et al (1976)	300 298 296 298	k <sub>c</sub> = 3.3 x 10 <sup>-12</sup> exp(-500/T) k <sub>c</sub> $\geq$ 1.2 x 10 <sup>-12</sup> (a) based on value of k(CH <sub>3</sub> $\theta_2$ + CH <sub>3</sub> $\theta_2$ ) Reaction proceeds exclusively through channel (c) k <sub>c</sub> /k = 0.80 $\pm$ 0.15 (k = k <sub>a</sub> + k <sub>b</sub> + k <sub>c</sub> )	estimate lower limit (a)
	Pate, Finlayson, Pitts (1974) Simonaitis, Helcklen (1974)	296 298		

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of log k
10, hv	$\text{N}_2 + \text{hv} \rightarrow \text{N} + \text{O}$ *Bates (1974)		$\lambda/\text{nm}$ 295-358 398 400 405 410 $\phi(\text{O}(^3\text{P}))$ 1.0 - 0.0008 ( $\lambda = 275$ ) 0.90 ± 0.05 0.76 ± 0.04 0.37 ± 0.02 0.14 ± 0.01 (a) Based on analysis of data in Jones and Bayes (1973) Extinction coefficient values tabulated for $\lambda = 185\text{-}410\text{nm}$ at 0.125nm intervals. Also gives values for $\text{N}_2\text{O}_4$ see reverse reaction	(a)
10, M	$\text{N}_2 + \text{M} \rightarrow \text{N} + \text{O} + \text{M}$		No recommendation	
10, 10	$\text{N}_2 + \text{N}_2 \rightarrow \text{N}_4$ Baulch, et al (1973) review			
10, 10	$\text{N}_2 + \text{N}_2 \rightarrow \text{N}_4 + \text{O}_2$ Baulch, et al (1973) review			
10, 10M	$\text{N}_2 + \text{N}_2 + \text{M} \rightarrow \text{N}_4 + \text{M}$ (f) $\text{N}_2\text{O}_4 + \text{M} \rightarrow \text{N}_2 + \text{N}_2\text{O}_2 + \text{M}$ (r) Baulch, et al (1973) review	600-2000	$3.3 \times 10^{-12} \exp(-13540/T)$ $k_1[\text{N}_2\text{O}_2]^2 = -(1/2)d[\text{N}_2]/dt$ $k_f = 3.0 \times 10^{-35} \exp(1040/T)\text{cm}^6\text{molecule}^{-2}\text{s}^{-1}$ (a) corrected expression; $k_f = k_r k_{eq}$ $k_r = 4.2 \times 10^{-7} \exp(-5550/T)$ $k_r = 3.3 \times 10^{-7} \exp(-5540/T)$	±0.15 (500-1000K) increasing to 0.3 at 2000K $M = \text{N}_2$ ±0.15 $M = \text{N}_2$ ±0.15
10, 11	$\text{N}_2 + \text{N}_2 \rightarrow \text{N}_4$ $\text{O}_2 + \text{N}_2 \rightarrow \text{N}_2\text{O}_2 + \text{O}_2$ *Baulch, et al (1973) review	300-850 300-500 338-396	$k_f = 2.3 \times 10^{-13} \exp(-1000/T)$ $k_r = k_f/k_{eq} = 8 \times 10^{-41} \exp(400/T)\text{cm}^6\text{molecule}^{-2}\text{s}^{-1}$ $k_f = 2.5 \times 10^{-14} \exp(-1230/T)$ (a) Based on $\text{N}_2\text{O}_5 + \text{M} \rightarrow \text{N}_2\text{O}_4 + \text{M}$ and $2\text{N}_2\text{O}_2 \rightarrow 2\text{N}_2 + \text{O}_2$	±0.4 (a) ±0.4
10, 11M	$\text{N}_2 + \text{N}_2 + \text{M} \rightarrow \text{N}_4 + \text{M}$ This survey			
14, M	$\text{N}_2 + \text{N}_2 + \text{M} \rightarrow \text{N}_4 + \text{M}$ (f) $\text{N}_2\text{O}_5 + \text{M} \rightarrow \text{N}_2\text{O}_4 + \text{M}$ (r)	217 217 222 227 237 251 265	Elevation/km $k(T, M)$ $\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$ 15 20 25 30 35 40 45 1.9 x 10 <sup>-12</sup> 1.1 x 10 <sup>-12</sup> 6.6 x 10 <sup>-13</sup> 4.5 x 10 <sup>-13</sup> 2.7 x 10 <sup>-13</sup> 1.5 x 10 <sup>-13</sup> 7.1 x 10 <sup>-14</sup>	18.60 18.27 17.93 17.58 17.26 16.92 16.60



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of log k
	Graham, Johnston (1978)	300	$k_f = 3.8 \times 10^{-12} \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ (2d order high pressure limit)	$\pm 0.4$ (a)
		300-340	$k_r = 2.2 \times 10^{-5} \exp(-9700/T) \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	$\pm 0.3$ (a)
		273-300	$k_r = 5.7 \times 10^{14} \exp(-10600/T) \text{ s}^{-1}$ (limiting first order expression)	$\pm 0.4$ (a)
		298-329	$K_{eq} = 1.2 \times 10^{-27} \exp(-11180/T) \text{ cm}^3 \text{molecule}^{-1}$	
		273-300	$k_f(M) = 1.48 \times 10^{-13} \exp(-861/T) \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	(b)
			$k_r(M) = 1.24 \times 10^{14} \exp(-10317/T) \text{ s}^{-1}$	(c)
			(a) $k_f = K_{eq} k_r$ , $M = \text{N}_2, \text{O}_2, \text{NO}$	
			(b) 2d order rate const at 1 atm	
			(c) 1st order rate const at 1 atm	
		300	$k > k(\text{NO} + \text{NH}_2)$	
10,16	$\text{NO}_2 + \text{NH}_2 \rightarrow \text{products}$ Jayanty, et al (1975d)		$6.7 \times 10^{-12} \exp(-13900/T)$	
10,17	$\text{NO}_2 + \text{NH}_3 \rightarrow \text{NH}_2 + \text{HNO}_2$ Bedford, Thomas (1972)	615-660		
10,18	$\text{NO}_2 + \text{H} \rightarrow \text{HO} + \text{NO}$ Wagner, et al (1976a)			
	Clyne, Monkhouse (1977)	243-461	$7.1 \times 10^{-10} \exp(-505 \pm 84/T)$	$\pm 0.1$
	Bemand, Clyne (1977)	298-653	$4.80 \times 10^{-10} \exp(-400 \pm 70/T)$	$\pm 0.1$
10,19M	$\text{NO}_2 + \text{HO} + \text{M} \rightarrow \text{HNO}_3 + \text{M}(\text{f})$	298	$1.13 \pm 0.22 \times 10^{-10}$	
26,M	$\text{HNO}_3 + \text{M} \rightarrow \text{HO} + \text{NO}_2 + \text{M}(\text{r})$ †NASA (1977) eval	200-350	$\log_{10}(k_f) = -A/T/(B + T) - 0.5 \log_{10}(T/280)$ $A = A_1 + A_2 Z + A_3 Z^2 + A_4 Z^3$ $B = B_1 + B_2 Z + B_3 Z^2$ $A_1 = 31.62273$ $A_2 = -0.258304$ $A_3 = -0.0889287$ $A_4 = 2.520173 \times 10^{-3}$ $B_1 = -327.372$ $B_2 = 44.5586$ $B_3 = -1.38092$	
			where $Z = \log_{10}[\text{N}_2]$ and is applicable only for the ranges 200<T/K<350 and $16.3 < \log_{10}[\text{N}_2]/\text{molecule} \cdot \text{cm}^{-3} < 15.5$ , with an estimated reliability in log k of $\pm 0.10$ (reliability analogous to log). Air is approximately 6 percent less efficient than nitrogen as a third body; i.e., $k$ for ( $M = \text{Air}$ ) = $0.94 k$ for $M = \text{N}_2$ For normal tropospheric conditions of 1 atm air and T = 300K, above expression yields value of $k_f = 1.1 \times 10^{-11}$	
	Anastasi, Smith (1976)	296	$k_f = 2.6 \times 10^{-30} M = \text{N}_2$ Rel. Efficiencies: $\text{N}_2(1.0)$ , $\text{He}(0.34)$ , $\text{Ar}(0.42)$ , $\text{O}_2(0.68)$ , $\text{SF}_6(2.5)$	
		220-550	$n = -2.6$ for temp dependence of form $T^n$ , $M = \text{N}_2$ values given for bimolecular rate constant over	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
	Baulch, et al (1973) review	273-400	ranges T = 220-550K and [N <sub>2</sub> ] = 3 x 10 <sup>17</sup> to 1.6 x 10 <sup>19</sup> molecule cm <sup>-3</sup> k <sub>f</sub> = 3.6 x 10 <sup>-32</sup> exp(+1100/T)cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> M = He ±0.2 k <sub>r</sub> = 2.7 x 10 <sup>-9</sup> exp(-15400/T)cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> M = Ar ±0.4 k <sub>f</sub> = 8 x 10 <sup>-12</sup> cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> (2d order high pressure limit) k <sub>f</sub> = 2.3 ± 0.5 x 10 <sup>-30</sup> x (295/T) <sup>2.5</sup> cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> M = N <sub>2</sub> ±0.4 Relative efficiencies: N <sub>2</sub> (1.0), Ar(0.43), He(0.43) k <sub>f</sub> = 2.9 ± 0.4 x 10 <sup>-30</sup> M = N <sub>2</sub> k <sub>f</sub> = 5.3 x 10 <sup>-12</sup> cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> (2nd order rate constant at 1 atm. H <sub>2</sub> O vapor) 15 ± 5 x 10 <sup>-31</sup> cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> , M = Ar 26 ± 10 x 10 <sup>-31</sup> , M = N <sub>2</sub> k <sub>f</sub> = (1.0 ± 0.1) x 10 <sup>-30</sup> M = Ar Values given for bimolecular rate constant over pressure range 25-646 torr Ar	
10,20M	N <sub>2</sub> + H <sub>2</sub> O + M → H <sub>2</sub> O <sub>2</sub> N <sub>2</sub> + M †NASA (1977) eval	200-300	tentatively, use expression given for rxn 10,36M N <sub>2</sub> O + Cl <sub>2</sub> + M → N <sub>2</sub> O <sub>2</sub> Cl <sub>2</sub> + M with twice the assigned uncertainty	
	Simonaitis, Heicklen (1978)	245 295	k[M]/k(H <sub>2</sub> O + N <sub>2</sub> ) = 0.61 (a) k[M]/k(H <sub>2</sub> O + N <sub>2</sub> ) = 0.036 (a) (a) M = 700 torr H <sub>2</sub> . Rxn to give H <sub>2</sub> ON <sub>2</sub> + O <sub>2</sub> is negligible	(a) (a)
	Howard (1977)	300	(2.09 ± 0.52) x 10 <sup>-31</sup> M = N <sub>2</sub> Pel. eff: N <sub>2</sub> (1.0), He(0.48); O <sub>2</sub> (0.72); N <sub>2</sub> O(3.2) k(H <sub>2</sub> O + N <sub>2</sub> → H <sub>2</sub> ON <sub>2</sub> + O <sub>2</sub> ) < 3 x 10 <sup>-15</sup> k[M] = 1.2 x 10 <sup>-13</sup> for [M] = 1 atm product probably H <sub>2</sub> O <sub>2</sub> N <sub>2</sub> not H <sub>2</sub> ON <sub>2</sub> + O <sub>2</sub> as assumed by authors	
10,28	N <sub>2</sub> O + S <sub>2</sub> → S <sub>2</sub> O <sub>3</sub> + N <sub>2</sub>			
10,35M	Davis (1974) N <sub>2</sub> O + Cl + M → N <sub>2</sub> O <sub>2</sub> Cl + M	300	k < 2 x 10 <sup>-24</sup>	
10,36M	†Clyne, White (1974) N <sub>2</sub> O + Cl <sub>2</sub> + M → Cl <sub>2</sub> ON <sub>2</sub> + M †NASA (1977) eval	298 200-300	7.2 x 10 <sup>-31</sup> cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> 3.3 x 10 <sup>-23</sup> T <sup>-3.34</sup> 1 ± 8.7 x 10 <sup>-9</sup> T <sup>-0.6</sup> [M] <sup>0.5</sup> (a) Expression developed by Zahniser, Chang, Kaufman (1977) (4.40 ± 0.66) x 10 <sup>-33</sup> exp(1087/T) M = N <sub>2</sub>	±0.6 at 230K ±0.3, -0.1 at 230K (a)
	Birks, et al (1977)	250-356		



No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Notes and Reliability of log k
	Leu, Lin, DeMore (1977)	297	$9.57 \pm 0.28 \times 10^{-32}$ $(3.69 \pm 0.24) \times 10^{-33} \exp(1150/T)$	M = He M = N <sub>2</sub>
	Zahniser, Chang, Kaufman (1977)	248-417 251-365 251-365 297	$(2.66 \pm 0.35) \times 10^{-33} \exp(1140/T)$ $(3.54 \pm 0.06) \times 10^{-33} \exp(950/T)$ $5.53 \times 10^{-24} T^{-3.15}$ $1.52 \times 10^{-31}$	M = He M = He M = He M = N <sub>2</sub>
	N <sub>2</sub> + CH <sub>3</sub> → CH <sub>3</sub> δ + Nδ		(a) Two forms of temp dependence are given: $\exp(B/T)$ and $T^{-n}$	
10,49	Glanzer, Troe (1973) review	300-1400	$3.3 \times 10^{-11}$ (a) Based on $k/k(\text{CH}_3 + \text{Nδ}) = 3.3$ at room temperature and on shock wave pyrolysis of CH <sub>3</sub> Nδ <sub>2</sub>	±0.25 (a)
10,49M	Nδ <sub>2</sub> + CH <sub>3</sub> (→M) → CH <sub>3</sub> Nδ <sub>2</sub> (→M)			
	Glanzer, Troe (1973) review	295	$2.8 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (a) Based on high pressure limit rate ratio $k/k(\text{CH}_3 + \text{Nδ}) = 1.7$	±0.1 (a)
10,49M	Nδ <sub>2</sub> + CH <sub>3</sub> + M → CH <sub>3</sub> Nδ <sub>2</sub> + M			
	Glanzer, Troe (1973) review	500-1400	$6.9 \times 10^{-31} \text{ cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$ , M = Ar (a) $k_f = k_r^{\text{eq}}$ see CH <sub>3</sub> δ + Nδ <sub>2</sub> → . . .	(a)
10,50 10,51	Nδ <sub>2</sub> + CH <sub>3</sub> δ → . . . Nδ <sub>2</sub> + CH <sub>3</sub> δ <sub>2</sub> → CH <sub>3</sub> δ <sub>2</sub> Nδ <sub>2</sub> (a) → CH <sub>2</sub> δ + HδNδ <sub>2</sub> (b) → CH <sub>3</sub> δ + Nδ <sub>3</sub> (c)	298	$k_a/k = 0.75 \pm 0.05$ $k_b/k = 0.25 \pm 0.1$ $k_c/k < 0.1$	
	*Heicklen (1973) review			
11,1v	Nδ <sub>3</sub> + hν → Nδ <sub>2</sub> + δ (1) → Nδ + δ <sub>2</sub> (2)	300	table of absorption cross sections for wavelength region 400-700nm $J_1 = 0.099 \text{ s}^{-1}$ $J_2 = 0.04 \text{ s}^{-1}$ (a) This table also given in Molina's review (1977). Values supersede results in Johnston, Graham (1974) (b) Specific photodissociation rates for troposphere with overhead sun. see reverse reaction	(a) (b) (b)
11,M	Nδ <sub>3</sub> + M → Nδ <sub>2</sub> + δ + M			
11,11	Nδ <sub>3</sub> + Nδ <sub>3</sub> → 2Nδ <sub>2</sub> + δ <sub>2</sub> Graham, Johnston (1978) Baulch, et al (1973) review	298-329 293-309	$8.5 \times 10^{-13} \exp(-2450/T)$ $5 \times 10^{-12} \exp(-3000/T)$ (a) $-d[\text{Nδ}_3]/dt = 2k[\text{Nδ}_3]^2$ . Based on rate of δ <sub>3</sub> decomp. in presence of N <sub>2</sub> δ <sub>5</sub>	±0.5 (a)

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
11,22	N <sub>2</sub> O + H <sub>2</sub> O → HNO <sub>3</sub> + HO Baulch, et al (1973) review	300	2.3 x 10 <sup>-26</sup> (a) k <sub>f</sub> = k <sub>r</sub> K <sub>eq</sub>	±0.3 (a)
11,28	N <sub>2</sub> O + SO <sub>2</sub> → NO <sub>2</sub> + SO <sub>3</sub> Daubendiek, Calvert (1975) Davis (1976)	300 300	k < 7 x 10 <sup>-21</sup> k < 1 x 10 <sup>-21</sup>	preliminary
12,M	N <sub>2</sub> + M → N + N + M *Baulch, et al (1973) review	6000-15000	6.1 x 10 <sup>-3</sup> T <sup>-1.6</sup> exp(-113,200/T)cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> M = N <sub>2</sub>	±0.5
12,19	N <sub>2</sub> + HO → N <sub>2</sub> O + H *Baulch, et al (1973) review	700-2500	5.4 x 10 <sup>-12</sup> exp(-40.4 x 10 <sup>3</sup> /T) (a) k <sub>f</sub> = k <sub>r</sub> K <sub>eq</sub> (corrected expression)	±0.2 (a)
13,hv	N <sub>2</sub> O + hv → products Johnston, Selwyn (1975)	294	figures of absorption cross section vs λ for λ = 210-328nm. Negligible absorption for λ > 260nm	
13,M	N <sub>2</sub> O + M → N <sub>2</sub> + O + M N <sub>2</sub> O + H → N <sub>2</sub> + HO Selwyn, et al (1977)	194-302	Absorption cross section values tabulated for λ = 173-240nm at 1 nm intervals for T = 194, 225, 243, 265 and 302K see reverse reaction	
13,18	N <sub>2</sub> O + M → N <sub>2</sub> + O + M N <sub>2</sub> O + H → N <sub>2</sub> + HO Albers, et al (1975)	718-1111 700-2500 773	(3.6 ± 1.2) x 10 <sup>-10</sup> exp(-8710 ± 350/T) 1.26 x 10 <sup>-10</sup> exp(-7600/T) 4.3 x 10 <sup>-15</sup>	±0.2 ±0.15
13,19	N <sub>2</sub> O + HO → products Chang, Kaufman (1977) Bierman, et al (1976) Atkinson, Perry, Pitts (1976)	480 298 298 443 440	< 4 x 10 <sup>-16</sup> 3.8 ± 1.2 x 10 <sup>-17</sup> < 2 x 10 <sup>-16</sup> < 2 x 10 <sup>-16</sup> < 2 x 10 <sup>-14</sup>	upper limit only
13,36	Gordon, Mulac (1975) N <sub>2</sub> O + ClO → products +NASA (1977) eval	200-300	k ≤ 1 x 10 <sup>-12</sup> exp(-4260/T) (a) "A factor" chosen. Fact fitted to upper limit at T = 587K reported in Watson's review (1977)	±2 at 230K (a)
13,45	N <sub>2</sub> O + CO → products Milks, Matula (1973)	1169-1655	3.5 x 10 <sup>-13</sup> exp(-8650/T)	(a)
14,hv	N <sub>2</sub> O <sub>5</sub> + hv → . . . Molina (1977) review		table of absorption cross sections for wavelength range 206-380nm (a) Gives values for λ = 205-310nm from Graham, Johnston (1977) which supersede results in Johnston, Graham (1974). Also gives values for λ = 320-380nm from Jones, Wulf (1937)	

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of $\log k$
<p>Murphy (1969)</p> <p>Hampson, et al (1973a) review</p> <p>Daubendiek, Calvert (1974)</p> <p><math>\text{N}_2\text{O}_5 (+M) \rightarrow \text{NO}_2 + \text{NO}_3 (+M)</math></p> <p>*This survey</p>				
14,M		217	$0.5 \times 10^{-15}$ $\lambda = 280 \text{ nm}$	
		217	Meas. by Holmes, Daniels (1934) are consistent	
		222	Survey of photochemistry	
		227	Figure giving molar extinction coefficient	
		237	vs $\lambda$ . $250 < \lambda/\text{nm} < 350$	
		251	Elevation/km $k/\text{s}^{-1}$	
		265	15 $5.0 \times 10^{-7}$	$\pm 0.5$
			20 $1.8 \times 10^{-7}$	
			25 $3.0 \times 10^{-7}$	
			30 $5.5 \times 10^{-7}$	
			35 $1.9 \times 10^{-6}$	
			40 $1.6 \times 10^{-5}$	
			45 $4.4 \times 10^{-5}$	
			Extrapolated from data taken between 273 and 353 K.	
			See reverse reaction	
14,M	$\text{N}_2\text{O}_5 + M \rightarrow \text{NO}_2 + \text{NO}_3 + M$	300	$k < 1 \times 10^{-20}$	
14,22	$\text{N}_2\text{O}_5 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3$	298	$< 1.3 \times 10^{-20}$	
	Hampson, et al (1973) review			
	Morris, Niki (1973)			
14,28	$\text{N}_2\text{O}_5 + \text{SO}_2 \rightarrow \text{products}$	300	$k < 4 \times 10^{-23}$	
	Daubendiek, Calvert (1974)	300	$k < 8 \times 10^{-24}$	
	Davis (1974)			
16,18M	$\text{NH}_2 + \text{H} + M \rightarrow \text{NH}_3 + M$	2000-3000	$1.2 \times 10^{-33} \exp(11200/T) \text{ cm}^6 \text{ molecule}^{-2}\text{s}^{-1}$ , $M \rightarrow \text{Ar} \pm 0.3$ (a)	
	Baulch, et al (1973) review		(a) $k_f \rightarrow K_{eq} k_r$	
			See reverse reaction	
16,19	$\text{NH}_2 + \text{H}_2 \rightarrow \text{NH}_3 + \text{H}$	800	$< 10^{-16}$ $k_f = K_{eq} k_r$	approximate
16,21	$\text{NH}_2 + \text{H}_2 \rightarrow \text{NH}_3 + \text{H}$		no data, no recommendation	
16,22	$\text{NH}_2 + \text{H}_2\text{O} \rightarrow \text{NH}_3 + \text{H}_2\text{O}$			
17,M	$\text{NH}_3 + M \rightarrow \text{NH}_2 + \text{H} + M$	2000-3000	$k = 1.5 \times 10^{-8} \exp(-42400/T) (M = \text{Ar})$	$\pm 0.7$ (a)
	Baulch, et al (1973) review		(a) To be used when $P(M) < 4 \text{ atm.}$	
17,18	$\text{NH}_3 + \text{H} \rightarrow \text{NH}_2 + \text{H}_2$	800	$< 10^{-16}$	approximate
17,19	$\text{NH}_3 + \text{H}_2 \rightarrow \text{NH}_2 + \text{H}_2\text{O}$			
	Baulch, et al (1973) review			
	$\text{NH}_3 + \text{H}_2 \rightarrow \text{NH}_2 + \text{H}_2\text{O}$			
	*Smith, Zellner (1975)	228-472	$2.3 \times 10^{-12} \exp(-800/T)$	$\pm 0.1$
	Perry, Atkinson, Pitts (1976a)	298-427	$2.93 \times 10^{-12} \exp(-850 \pm 150/T)$	preliminary
	Davis (1977)	300	$(1.4 \pm 0.1) \times 10^{-13}$	
	Cox, Derwent, Holt (1975)	300	$(1.2 \pm 0.4) \times 10^{-13}$	
	Hack, et al (1974)	298-669	$5.3 \pm 0.8 \times 10^{-12} \exp(-920/T)$	
	Stuhl (1973a)	298	$1.5 \times 10^{-13}$	$\pm 0.1$
	Kurylo (1973)	298	$4.1 \times 10^{-14}$	$\pm 0.06$
	Gordon, Mulac (1975)	418	$4.2 \pm 0.5 \times 10^{-13}$	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of $\log k$
17,36	$\text{NH}_3 + \text{ClO} \rightarrow \text{products}$ Walker (1972)	670	$k < 5 \times 10^{-16}$ (a) As quoted by Watson (1977) review.	(a)
17a,18	$\text{N}_2\text{H}_4 + \text{H} \rightarrow \text{H}_2 + \text{N}_2\text{H}_3$ Stief, Payne (1976)	228-400	$(9.87 \pm 1.17) \times 10^{-12} \exp(-1200 \pm 50/T)$	
18,18M	$\text{H} + \text{H} + \text{M} \rightarrow \text{H}_2 + \text{M} \text{ (f)}$			
21,M	$\text{H}_2 + \text{M} \rightarrow \text{H} + \text{H} + \text{M} \text{ (r)}$ Baulch, et al (1972) review	300 1700-5000	$k_f = 8.3 \times 10^{-23} \text{ cm}^6 \text{molecule}^{-2} \text{s}^{-1} \quad \text{M} = \text{H}_2$ $= 1.8 \times 10^{-30} \text{ T}^{-1} \quad \text{M} = \text{Ar}$ Rel. efficiencies: $\text{H}_2(1.0)$ , $\text{Ar}(0.25)$ $-d[\text{H}]/dt = 2k[\text{H}]^2[\text{M}]$ $k_r = 3.7 \times 10^{-10} \exp(-48300/T) \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ $\text{M} = \text{Ar}$ see reverse reaction	$\pm 0.2$ $\pm 0.3$ $\pm 0.3$
18,19	$\text{H} + \text{HO} \rightarrow \text{H}_2 + \text{O}$			
18,19M	$\text{H} + \text{HO} + \text{M} \rightarrow \text{H}_2\text{O} + \text{M} \text{ (f)}$			
22,M	$\text{H}_2\text{O} + \text{M} \rightarrow \text{H} + \text{HO} + \text{M} \text{ (r)}$ *Baulch, et al (1972) review	1000-3000 2000-6000	$k_f = 6.1 \times 10^{-26} \text{ T}^{-2} \text{ cm}^6 \text{molecule}^{-2} \text{s}^{-1}, \text{M} = \text{N}_2$ $k_r = 5.8 \times 10^{-9} \exp(-52900/T) \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ $\text{M} = \text{N}_2$ $k_f = 4.3 \times 10^{-25} \text{ T}^{-2.6} \quad \text{M} = \text{He}$ Rel eff: $\text{He}(1.0)$ , $\text{Ar}(1.5)$ , $\text{N}_2(3.2)$	$\pm 0.3$ $\pm 0.2$ $\pm 0.3$
18,20	Zellner, Erler, Field (1977)	230-300		
18,20	$\text{H} + \text{HO}_2 \rightarrow \text{HO} + \text{HO} \text{ (1f)}$	290-800	$k_{1f} = 4.2 \times 10^{-10} \exp(-950/T)$	$\pm 0.3 \text{ (a)}$
19,19	$\text{HO} + \text{HO} \rightarrow \text{H} + \text{HO}_2 \text{ (1r)}$		$k_{1r} = k_{1f}/K_{eq} = 2.0 \times 10^{-11} \exp(-20200/T)$	$\pm 0.3$
18,20	$\text{H} + \text{HO}_2 \rightarrow \text{H}_2 + \text{O}_2 \text{ (2)}$	290-800	$k_2 = 4.2 \times 10^{-11} \exp(-750/T)$	$\pm 0.4$
18,20	$\text{H} + \text{HO}_2 \rightarrow \text{H}_2\text{O} + \text{O} \text{ (3)}$ Baulch, et al (1972) review		$k_{2r} = k_{2f}/K_{eq} = 9.1 \times 10^{-11} \exp(-29100/T)$ $k_3 = \text{no recommendation}$ $k_{1f}$ and $k_2 = \text{same as Baulch, et al}$	$\pm 0.4$
	Lloyd (1974) review	300-1000	$k_3 = 8.3 \times 10^{-11} \exp(-500/T)$	$> \pm 0.7 \text{ (a)}$ $> \pm 0.5 \text{ (b)}$
	Moortgat, Allen (1973)	297	$k_{1f} = 8.3 \times 10^{-12}$ $k_2 = 1 \times 10^{-11}$ $k_3 = 6.7 \times 10^{-12}$ (a) Reaction 1 and rate ratios control values for reactions 2 and 3. (b) Temperature coefficient estimated. (c) Secondary reaction in $\text{H} + \text{O}_2 + \text{M}$ systems, fit of concentration vs. time plots.	? (c) ? (c) ? (c)

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>2</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
18,22	H + H <sub>2</sub> O → H <sub>2</sub> + HO Baulch, et al (1972) review	300-2500	1.5 x 10 <sup>-10</sup> exp(-10250/T)	±0.06 at 300K increasing to ±0.2 at high temp.
18,23	H + H <sub>2</sub> O <sub>2</sub> → H <sub>2</sub> + HO <sub>2</sub> (a) → HO + H <sub>2</sub> O (b) *Klemm, Payne, Stief (1975) Gorse, Volman (1974)	283-353 300	k <sub>a</sub> + k <sub>b</sub> = 5.2 ± 2.0 x 10 <sup>-12</sup> exp(-1400 ± 140/T) k <sub>a</sub> = 3.1 ± 0.8 x 10 <sup>-15</sup> k <sub>b</sub> = 5.7 ± 1.4 x 10 <sup>-15</sup>	
	Hampson, et al (1973)	300-800	k <sub>a</sub> = 2.8 x 10 <sup>-12</sup> exp(-1900/T) k <sub>b</sub> no recommendation (a) Accepts recommendations of Baulch, et al (1972). k <sub>b</sub> /k <sub>a</sub> = 1.3	(a) (a)
18,24	Meagher, Heicklen (1975) H + HNO → H <sub>2</sub> + NO Hampson, et al (1973) review Baulch, et al (1973) review	298 211-703 2000 300 2000	> 5 x 10 <sup>-14</sup> 7 x 10 <sup>-12</sup> 10 <sup>-13</sup> < k < 10 <sup>-12</sup> 8 x 10 <sup>-12</sup>	±0.3 ±0.3 ±0.2
18,25	H + HNO <sub>2</sub> → products Hampson, et al (1973) review Baulch, et al (1973) review		No data No recommendation	
18,26	H + HNO <sub>3</sub> → products Hampson, et al (1973) review Chapman, Wayne (1974) Baulch, et al (1973) review	300 300 300	< 1 x 10 <sup>-13</sup> < 2 x 10 <sup>-15</sup> < 10 <sup>-13</sup>	approximate upper limit
18,28M	H + SO <sub>2</sub> + M → SO <sub>3</sub> + M Baulch, et al (1976) review	1650-2120	1.4 x 10 <sup>-32</sup>	±0.2
18,31 21,26a	H + HS → H <sub>2</sub> + S (f) H <sub>2</sub> + S → H + HS (r) Baulch, et al (1976) review	298	k <sub>f</sub> = 2.5 x 10 <sup>-11</sup> k <sub>r</sub> = 2.2 x 10 <sup>-25</sup> (a) k <sub>r</sub> = k <sub>f</sub> /K <sub>eq</sub> k <sub>f</sub> = 2.5 ± 0.8 x 10 <sup>-11</sup>	±0.2 ±0.2 (a)
18,32	Cupitt, Glass (1974) H + H <sub>2</sub> S → H <sub>2</sub> + HS Kurylo, Peterson, Braun (1971)	298 190-464	1.29 x 10 <sup>-11</sup> exp(860/T) (a) Selected by Baulch, et al (1976) review	(a)
18,34c	H + COS → CO + HS Baulch, et al (1976) review	300	2.2 x 10 <sup>-14</sup>	±0.2
18,37	H + ClO → HO + Cl *Watson (1977) review	298	5.7 ± 1.2 x 10 <sup>-11</sup>	



No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
18,39	H + HCl → H <sub>2</sub> + Cl Ambidge, et al (1976a)	298-521	7.8 x 10 <sup>-12</sup> exp(-1600±84/T)	
18,42	H + N <sub>2</sub> O → HCl + NO Spencer, Glass (1975)	295	4.3 x 10 <sup>-14</sup>	
18,44	H + N <sub>2</sub> O → HCl + NO Wagner, et al (1976a)	255-461	7.6 x 10 <sup>-11</sup> exp(-457 ± 72/T)	
	H + Cl <sub>2</sub> → HCl + Cl *This survey	250-700	1.46 x 10 <sup>-10</sup> exp(-593/T)	±0.1 (a)
18,45M	H + CO + M → HCO + M Baulch, et al (1976) review	300-730	(a) Based on data in Remand, Clyne and Wagner, et al (1.41 ± 0.24) x 10 <sup>-10</sup> exp(-575 ± 65/T)	
18,46	H + CO <sub>2</sub> → HCO + CO Wagner, et al (1976)	252-458	(1.44 ± 0.28) x 10 <sup>-10</sup> exp(-600 ± 70/T)	
18,48	H + CH <sub>2</sub> O → H <sub>2</sub> + CH <sub>2</sub> Ambidge, et al (1976)	292-434	(7.6 ± 2.2) x 10 <sup>-11</sup> exp(-714 ± 100/T)	
18,53	H + CH <sub>3</sub> COOH → CH <sub>3</sub> O + H <sub>2</sub> O (a) Ridley, et al (1972) Westenberg, deHaas (1972b) H + CH <sub>3</sub> COOH → CH <sub>3</sub> O + H <sub>2</sub> O (b) → CH <sub>3</sub> O <sub>2</sub> + H <sub>2</sub> (c) → CH <sub>2</sub> COH + H <sub>2</sub> (c) Slemr, Warneck (1977)	298-773 297 297-652 250-358	2.0 x 10 <sup>-33</sup> exp(-850/T), M = H <sub>2</sub> see reverse reaction 5.4 x 10 <sup>-14</sup> 2.2 x 10 <sup>-11</sup> exp(-1890/T) k = (2.8 ± 0.9) x 10 <sup>-13</sup> exp(-930 ± 95/T) k <sub>a</sub> /k = 0.43 ± 0.07 k <sub>b</sub> /k = 0.52 ± 0.07 k <sub>c</sub> /k = 0.05	±0.12 ±0.04 overall rate const
18,54	H + CH <sub>3</sub> ONH → CH <sub>3</sub> OH + NO (a) → CH <sub>2</sub> ONH + H <sub>2</sub> (b) → CH <sub>3</sub> O + HNO (c) Moortgat, et al (1977)	223-398	k = (4.3 ± 0.9) x 10 <sup>-13</sup> exp(-950 ± 55/T) k <sub>a</sub> /k = 0.47 ± 0.05 (k <sub>b</sub> + k <sub>c</sub> )/k = 0.53 ± 0.05	overall rate const
18,64	H + CH <sub>3</sub> Cl → HCl + CH <sub>3</sub> Westenberg, deHaas (1975a)	500-800	6.2 x 10 <sup>-11</sup> exp(-4650/T)	
19,M	H <sub>2</sub> + M → H + H + M Baulch, et al (1972) review		no recommendation (E/R > 50000 K) see reverse reaction	
19,19	H <sub>2</sub> + H <sub>2</sub> → H + H <sub>2</sub> H <sub>2</sub> + H <sub>2</sub> → H <sub>2</sub> + H <sub>2</sub> Baulch, et al (1972) review		no recommendation	
19,19	H <sub>2</sub> + H <sub>2</sub> → H <sub>2</sub> + H <sub>2</sub> (f) H <sub>2</sub> + H <sub>2</sub> → H <sub>2</sub> + H <sub>2</sub> (r) *Baulch, et al (1972) review	300-2000 300-2000 1500-2000	k <sub>f</sub> = 1.0 x 10 <sup>-11</sup> exp(-550/T) k <sub>r</sub> = 1.1 x 10 <sup>-10</sup> exp(-9240/T) k <sub>f</sub> = 5.1 x 10 <sup>-11</sup> exp(-3500/T)	±0.2 ±0.2
1,22	H <sub>2</sub> + H <sub>2</sub> → H <sub>2</sub> + H <sub>2</sub> (r) Rawlins, Gardiner (1974) Westenberg, deHaas (1973a)	300	k <sub>r</sub> = 2.3 x 10 <sup>-12</sup>	±0.06



No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
19,19M 23,M	McKenzie, et al (1973)	298	k <sub>f</sub> = 2.1 ± 0.5 x 10 <sup>-12</sup>	
	Clyne, Down (1974)	300	k <sub>f</sub> = 1.4 ± 0.2 x 10 <sup>-12</sup>	
		300	k <sub>f</sub> = 1.7 ± 0.6 x 10 <sup>-12</sup>	(a)
	Trainor, von Rosenberg (1974)	300	k <sub>f</sub> = 2.1 ± 0.2 x 10 <sup>-12</sup>	
	Wilson (1972) review	300	k <sub>f</sub> = 2.57 x 10 <sup>-12</sup>	+0.1, -0.5
	Hθ + Hθ + N → H <sub>2</sub> θ <sub>2</sub> + M (f)		(a) Evaluation based on authors' work and other recent data	
	H <sub>2</sub> θ <sub>2</sub> + M → Hθ + Hθ + M (r)		-d[θH]/dt = 2k[θH] <sup>2</sup> [M]	
	+NASA (1977) eval	200-300	k <sub>f</sub> = 1.25 x 10 <sup>-32</sup> exp(900/T)	(a)
	Trainor, von Rosenberg (1974)	300	(a) Accepts room temp value of Trainor, von Rosenberg. E/R value chosen same as for comb. of θH with Nθ	
	Baulch, et al (1972) review	700-1500	k <sub>f</sub> = 2.5 ± 0.3 x 10 <sup>-31</sup> M = N <sub>2</sub> k <sub>f</sub> = 2.5 x 10 <sup>-33</sup> exp(+2550/T)cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> M = N <sub>2</sub>	±0.1 700<T<950K
19,20	Hθ + Hθ <sub>2</sub> → H <sub>2</sub> θ + θ <sub>2</sub>		k <sub>r</sub> = 2.0 x 10 <sup>-7</sup> exp(-22900/T)cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	±0.3 at 1500K
	+NASA (1977) eval	200-300	M = N <sub>2</sub> ; k <sub>f</sub> = k <sub>r</sub> K <sub>eq</sub>	
	Burrows, Harris, Thrush (1977)	293	3 x 10 <sup>-11</sup> 5.1 x 10 <sup>-11</sup>	0.5 at 230K (a)
	Lloyd (1974) review	300-1000	(a) Based on k(Hθ + H <sub>2</sub> θ <sub>2</sub> ) = 7.9 x 10 <sup>-13</sup>	
	Hochanadel, et al (1972)	300	8.7 x 10 <sup>-11</sup> exp(-500/T)	±0.2
19,21	DeMore, Tschuikow-Roux (1974)	300	2 x 10 <sup>-10</sup> 1.1 x 10 <sup>-10</sup>	±0.5 (a)
	Hθ + H <sub>2</sub> → H <sub>2</sub> θ + H		(a) Adjusted value = based on reported ratio and values of k(θ <sub>3</sub> + Hθ) and k(Hθ <sub>2</sub> + Hθ <sub>2</sub> ) in this survey.	
	+Baulch, et al (1972) review	300-2500	3.6 x 10 <sup>-11</sup> exp(-2590/T) cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	±0.08 at 700K ±0.2 at high temp.
	Wilson (1972) review	300-2000	3.8 x 10 <sup>-11</sup> exp(-2600/T)	±0.3
	Stuhl, Niki (1972)	298	7.1 x 10 <sup>-15</sup>	±0.06
	Westenberg (1973a)	298	7.6 x 10 <sup>-15</sup>	±0.04
		352	1.8 x 10 <sup>-14</sup>	
		403	3.2 x 10 <sup>-14</sup>	
		518	1.4 x 10 <sup>-13</sup>	
		628	3.6 x 10 <sup>-13</sup>	
		745	6.6 x 10 <sup>-13</sup>	
	Smith, Zellner (1974a)	210-460	1.8 x 10 <sup>-11</sup> exp(-2330/T)	
	Gardiner, et al (1974)	1350-1600	8.7 x 10 <sup>-11</sup> exp(-3250/T)	
	Greiner (1969)	300-500	6.8 x 10 <sup>-12</sup> exp(-2020/T)	±0.15
	Atkinson, Hansen, Pitts (1975)	297-434	5.9 x 10 <sup>-12</sup> exp(-2008 ± 150/T)	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^2\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of $\log k$
19,21	Overend, et al (1975)	295	$(5.80 \pm 0.26) \times 10^{-15}$	
19,22	Trainor, von Rosenberg (1975)	300	$5.3 \times 10^{-15}$	
19,23	Vandooren, et al (1975)	600-1300	$1.2 \times 10^{-11} \exp(-2200/T)$	
	$\text{H}\theta + \text{D}_2 \rightarrow \text{HD}\theta + \text{D}$			
	Smith, Zellner (1974a)	210-460	$1.25 \times 10^{-11} \exp(-2550/T)$	
	$\text{H}\theta + \text{H}_2\theta \rightarrow \text{H}_2\theta_2 + \text{H}$		no recommendation	$\pm 0.3$ at 230K (a)
	Baulch, et al (1972) review			
	$\text{H}\theta + \text{H}_2\theta_2 \rightarrow \text{H}_2\theta + \text{H}\theta_2$			
	†NASA (1977) eval	200-300	$1 \times 10^{-11} \exp(-750/T)$	
	Hack, et al (1975)	298-670	$8.0 \times 10^{-12} \exp(-670/T)$	
	Greiner (1968)	300-458	$4.1 \times 10^{-13} T^{1/2} \exp(-600/T)$	
	Hampson (1973) quoting	300-800	$1.7 \times 10^{-11} \exp(-910/T)$	$\pm 0.2$
	Baulch, et al (1972) review			
	Gorse, Volman (1972)	300	$k/k(\text{H}\theta + \text{C}\theta) = 8.13$	
	Meagher, Heicklen (1974)	298	$k/k(\text{H}\theta + \text{C}\theta) = 4.1 \pm 0.6$	
19,24	$\text{H}\theta + \text{HN}\theta \rightarrow \text{H}_2\theta + \text{N}\theta$		(a) Based on results of Hack, et al (1975) and Greiner (1968)	
19,25	Hampson, et al (1973) review	1600-2100	$7 \times 10^{-11}$	$\pm 0.7$
	Baulch, et al (1973) review	2000	$6 \times 10^{-11}$	$\pm 0.2$
	$\text{H}\theta + \text{HN}\theta_2 \rightarrow \text{H}_2\theta + \text{N}\theta_2$			
	Cox, Derwent, Holt (1976)	296	$(6.6 \pm 0.3) \times 10^{-12}$	
			ref. rxn is $\text{H}\theta + \text{H}_2 \rightarrow \text{H}_2\theta + \text{H}$ with $k_{\text{ref}} = 7 \times 10^{-15}$	
19,26	Cox (1974a)	294	$k/k(\text{H}\theta + \text{C}\theta) = 15 \pm 1$ at 1 atm air	$\pm 0.1$ at 230K (a)
	$\text{H}\theta + \text{HN}\theta_3 \rightarrow \text{H}_2\theta + \text{N}\theta_3$			
	†Smith, Zellner (1975)	240-406	$8 \pm 2 \times 10^{-14}$	
	Margitan, et al (1975)	270-470	$8.9 \pm 1.3 \times 10^{-14}$	
	Hampson, et al (1973) review	300-650	$6 \times 10^{-13} \exp(-400/T)$	
	Johnston (1974) review	300-700	$(1.5 \text{ to } 2.0) \times 10^{-13}$	
	Baulch, et al (1975) review	300	$1.3 \times 10^{-13}$	
19,28M	$\text{H}\theta + \text{S}\theta_2 + \text{M} \rightarrow \text{HS}\theta_3 + \text{M}$		(a) Value recommended by authors for $T \leq 298\text{K}$ .	
	Davis (1976)	300	$10^{13} \frac{\text{K}}{\text{K}}$	(a)
			$0.87 \pm 0.06$	
		150	$1.6 \pm 0.1$	
		500	$2.7 \pm 0.2$	
			$10^{13} \frac{\text{K}}{\text{K}}$	
		50	$1.37 \pm 0.06$	
		150	$2.50 \pm 0.3$	
		500	$3.71 \pm 0.3$	
			$10^{13} \frac{\text{K}}{\text{K}}$	
		5	$0.08 \pm 0.08$	
		10	$1.43 \pm 0.14$	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of log k
			$2.44 \pm 0.3$	
			20	
			760	
			(b) Extrapolated value.	
	Harris, Wayne (1975)	300	$4.5 \pm 1.5 \times 10^{-31} \text{ cm}^6 \text{molecule}^{-2} \text{s}^{-1}$ , $M = \text{Ar}$	
			$7.2 \pm 2.6 \times 10^{-31}$ , $M = \text{N}_2$	
	Cox (1974a)	294	$k/k(\text{H}\theta + \text{C}\theta) = 4 \pm 0.5$	(c)
			(c) 2nd order rate constant, $[M] = 1 \text{ atm}$ . $\text{N}_2 + \theta_2$	
			see entry for $k(\text{H}\theta + \text{C}\theta)$	
	Gordon, Mulac (1975)	435	$1.8 \times 10^{-12} \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	
			(2d order rate constant at 1 atm. $\text{H}_2\theta$ vapor)	
	Castleman, Tang (1977)	297	$1.6 \times 10^{-31} \text{ cm}^6 \text{molecule}^{-2} \text{s}^{-1}$ , $M = \text{N}_2$	
			$k/k(\text{H}\theta + \text{C}\theta) = 4.3$	(d)
			(d) Measured ratio of 2d order rate constants at 1 atm $\text{N}_2$ .	
			Values for k given in paper assumed $k_{\text{ref}} = 1.5 \times 10^{-12}$	
	Atkinson, Perry, Pitts (1976)	298	over entire pressure range. See entry for $k(\text{H}\theta + \text{C}\theta)$	
			$10^{13} \frac{k}{k_{\text{ref}}}$	(e)
			1.35	
			2.16	
			3.10	
			4.38	
			5.87	
			6.55	
			(e) Effective 2nd order rate constant in $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	
19,32	$\text{H}\theta + \text{H}_2\text{S} \rightarrow \text{H}_2\theta + \text{HS}$			
	Baulch, et al (1976) review	300-500	$1.05 \times 10^{-11} \exp(-200/T)$	$\pm 0.15$
	Perry, Atkinson, Pitts (1976a)	297-427	$(5.2 \pm 0.5) \times 10^{-12}$	
	Stuhl (1974)	298	$3.1 \pm 0.5 \times 10^{-12}$	
			see reverse reaction	
19,35	$\text{H}\theta + \text{Cl} \rightarrow \theta + \text{HCl}$			
19,39	$\text{H}\theta + \text{HCl} \rightarrow \text{H}_2\theta + \text{Cl}$			
	†Watson (1977) review	220-300	$3.0 \times 10^{-12} \exp(-425/T)$	$\pm 0.1$ at 230K
	Smith, Zellner (1974a)	210-460	$4.1 \times 10^{-12} \exp(-530/T)$	
	Zahniser, et al (1974)	224-460	$2.0 \pm 0.1 \times 10^{-12} \exp(-312 \pm 10/T)$	
	Takacs, Glass (1973)	295	$6.4 \pm 1.5 \times 10^{-13}$	$\pm 0.1$
	Davis (1974)	300	$6.5 \pm 0.4 \times 10^{-13}$	
19,39	$\text{H}\theta + \text{DCI} \rightarrow \text{HD}\theta + \text{Cl}$			
	Smith, Zellner (1974a)	210-460	$4.7 \times 10^{-12} \exp(-780/T)$	
19,39Br	$\text{H}\theta + \text{HBr} \rightarrow \text{H}_2\theta + \text{Br}$			
	†Takacs, Glass (1973)	300	$5.1 \times 10^{-12}$	$\pm 0.36$ at 230K
19,43a	$\text{H}\theta + \text{N}\theta_3\text{Cl} \rightarrow \text{products}$			
	†Zahniser, et al (1977)	246-387	$(1.19 \pm 0.10) \times 10^{-12} \exp(-333 \pm 22/T)$	$\pm 0.36$ at 230K
19,45	Ravishankara, et al (1977)	245	$3.7 \times 10^{-13}$	
18,46	$\text{H}\theta + \text{C}\theta \rightarrow \text{C}\theta_2 + \text{H}$ (f)			
	$\text{H} + \text{C}\theta_2 \rightarrow \text{H}\theta + \text{C}\theta$ (r)			
	†NASA (1977) eval	200-400	$k_f = 1.4 \times 10^{-13}$	low pressure value
				$\pm 0.1$

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of log k
*Baulch, et al (1976) review		250-2500	$\log k_f = -12.95 + 3.94 \times 10^{-4} \times T$ same as given in Baulch, Drysdale (1974) recommendation based on low pressure studies	$\pm 0.1$
		1000-3000	$k_r = k_f/K_{eq} = 2.5 \times 10^{-10} \exp(-13300/T)$	$\pm 0.1$
		300	$k_f = 3.0 \times 10^{-13}$ (a) Tentative rec. 1 atmosphere air have shown an apparent increase in k in the presence of $\text{SF}_6$ , $\text{H}_2$ , $\text{N}_2/\theta_2$ but not He or Ar.	$\pm 0.1$ (a)
			Biermann, et al (1978) have identified a combined effect of inert gas and $\theta_2$ . Trace impurities of $\theta_2$ have a significant effect and 1 torr is sufficient to produce the limiting rate constant in the presence of 1 atm of $\text{N}_2$ .	
Butler, et al (1977)		298	<u>Experiments bearing on the "pressure effect"</u> Mole fractions	
			$k_f/k_{ref}$	$\text{N}_2 - \text{C}_2 - \text{C}_2\theta$
			0.094	100
			0.10	0.49/0.33/0.15
			0.17	0.50/0.34/0.15
			0.15	0.73/0.17/0.08
			0.16	0.50/0.34/0.15
			0.18	0.73/0.17/0.07
				0.74/0.17/0.08
			Ref. rxn is $\text{H}\theta + \text{isobutane} \rightarrow \text{products}$ , with 6-28% correction for side rxn. Ratios calculated from reported data using authors' value for $k_{ref} = 1.59 \times 10^{-12}$ which was determined in the 100 torr expt. relative to $k(\text{H}\theta + \text{C}\theta) = 1.5 \times 10^{-13}$	
Chan, et al (1977)		298	$k_f/k_{ref}$	P/torr
			0.059	100 in "synthetic air"
			0.127	700
Sie, Simonaits, Heicklen (1976a)		217-298 298	Ref. rxn is $\text{H}\theta + \text{isobutane} \rightarrow \text{products}$	
			$k_f/k_{ref}$	P/torr
			0.2exp(1700/T)	high pressure limit in $\text{H}_2$
			14	20 ( $\text{H}_2$ )
			21	83 ( $\text{H}_2$ )
			42	296 ( $\text{H}_2$ )
			49	702 ( $\text{H}_2$ )
			17	605 ( $\text{He}$ ) + 2 $\theta$ ( $\text{H}_2$ )
			33	550 ( $\text{SF}_6$ ) + 2 $\theta$ ( $\text{H}_2$ )
			49	605 ( $\text{SF}_6$ ) + 2 $\theta$ ( $\text{H}_2$ )
Cox, Derwent, Holt (1976)		296	Ref. rxn is $\text{H}\theta + \text{H}_2 \rightarrow \text{H}_2\theta + \text{H}$	
			$k_f/k_{ref}$	P/torr
Atkinson, Perry, Pitts (1976)		299	38.6	700 $\text{N}_2/\theta_2 = 2$
			Ref. rxn. is $\text{H}\theta + \text{H}_2 \rightarrow \text{H}_2\theta + \text{H}$	
			(1.54 $\pm$ 0.16) $\times 10^{-13}$ P = 25-654 torr Ar	

No.	Reaction/Reference * - Preferred Value † - NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$		Notes and Reliability of log k
			$10^{13} \frac{x}{k}$	$\frac{P(\text{SF}_6)}{\text{torr}}$	

19.48	Perry, Atkinson, Pitts (1977a)	299	1.53 1.93 2.40 3.09 3.42	25 76 208 403 603	
	Everend, Paraskevopoulos (1977)	296	(2.03 $\pm$ 0.08) $\times 10^{-13}$ at 50 torr He (3.24 $\pm$ 0.20) $\times 10^{-13}$ at 200, 350 torr SF <sub>6</sub>		
	Gordon, Mulac (1975)	298	1.50 $\times 10^{-13}$ at 710 torr Ar, 10 torr H <sub>2</sub> <sup>o</sup>		
	Trainor, von Rosenberg (1975)	300	Experiments either at low pressure or for which pressure effects were not studied.		
	Vandooren, et al (1975)	400-800	1.25 $\times 10^{-13}$		
	Blordi, et al (1975)	1000-1800	1.33 $\times 10^{-13}$		
	Steinert, Zellner (1975)	1350-1750	3.85 $\times 10^{-12}$ exp(-2850/T)		
	Wilson (1972) review	300-900	7.8 $\times 10^{-13}$		
	Stuhl, Niki (1972)	300-2000	log k = -12.93 + 4.0 $\times 10^{-4}$ x T		$\pm 0.3$
	Westenberg, deHaas (1973a)	300	5.1 $\times 10^{-13}$ exp(-300/T)		$\pm 0.06$
		298	1.35 $\times 10^{-13}$		$\pm 0.04$
		396	1.33 $\times 10^{-13}$		
		523	1.38 $\times 10^{-13}$		
		707	1.44 $\times 10^{-13}$		
		915	1.69 $\times 10^{-13}$		
	Davis, Fischer, Schlif (1974)	220-373	2.17 $\times 10^{-13}$		
	Greiner (1969)	300-500	2.15 $\pm$ 0.15 $\times 10^{-13}$ exp(-80 $\pm$ 40/T)		$\pm 0.04$
	Smith, Zellner (1973)	300	2.1 $\times 10^{-13}$ exp(-115/T)		$\pm 0.1$
			1.45 $\times 10^{-13}$		(h)
			(b) Data reported 210 < T < 460 K. Slight positive temperature dependence, possibly curved.		
	Peeters, Mahnen (1973)	1600-1900	4.7 $\times 10^{-13}$		
	Gardiner, et al (1973)	1500-2000	6.7 $\times 10^{-12}$ exp(-4000/T)		
	Howard, Evenson (1974)	296	1.56 $\pm$ 0.2 $\times 10^{-13}$		
	H <sub>2</sub> + CH <sub>3</sub> <sup>o</sup> $\rightarrow$ H <sub>2</sub> <sup>o</sup> + CH <sub>3</sub> <sup>o</sup>				
	†NASA (1977) eval	200-300	3 $\times 10^{-11}$ exp(-250/T)		$\pm 0.4$ at 230K (a)
19.52	*Morris, Niki (1971)	300	1.4 $\times 10^{-11}$		$\pm 0.1$
	Wilson (1972) review	300-1600	8 $\times 10^{-11}$		$\pm 1$
	Peeters, Mahnen (1973)	1400-1800	3.8 $\times 10^{-11}$		
			(a) "A factor" chosen; E <sub>act</sub> fitted to room temp value of Morris, Niki		
	H <sub>2</sub> + CH <sub>3</sub> <sup>o</sup> H $\rightarrow$ products				
19.54	Campbell, et al (1976)	292	9.5 $\times 10^{-13}$		
	H <sub>2</sub> + CH <sub>3</sub> NO <sub>2</sub> $\rightarrow$ products				
19.54	Campbell, Goodman (1975b)	292	9.2 $\times 10^{-13}$		
	H <sub>2</sub> + CH <sub>3</sub> ONO $\rightarrow$ products				
	Campbell, Goodman (1975b)	292	1.3 $\times 10^{-12}$		



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
19,56	H <sub>2</sub> + CH <sub>4</sub> → CH <sub>3</sub> + H <sub>2</sub> θ			
	†Davis, Fischer, Schliff (1974)	240-373	2.36 ± 0.21 x 10 <sup>-12</sup> exp(-1710 ± 88/T)	±0.04
	Margitan, et al (1974)	290-440	3.83 x 10 <sup>-12</sup> exp(-1840 ± 20/T)	±0.06
	Wilson (1972) review	300-2000	4.7 x 10 <sup>-11</sup> exp(-2500/T)	±0.7
	Overend, et al (1975)	300	6.51 ± 0.26 x 10 <sup>-15</sup>	
	Peeters, Mahnen (1973)	1100-1900	5 x 10 <sup>-11</sup> exp(-3000/T)	
	Greiner (1970)	300-500	5.5 x 10 <sup>-12</sup> exp(-1900/T)	±0.1
	Gordon, Mulac (1975)	381	2.6 x 10 <sup>-14</sup> (1 atm H <sub>2</sub> θ vapor)	
		416	5.5 x 10 <sup>-14</sup> (1 atm H <sub>2</sub> θ vapor)	
		Zellner, Steinert (1976)	300-900	5.76 x 10 <sup>-21</sup> T <sup>3.08</sup> exp(-1010/T)
19,56a	Howard, Evenson (1976)	296	(9.5 ± 1.4) x 10 <sup>-15</sup>	
	Cox, Derwent, Holt (1976)	296	(7.3 ± 0.9) x 10 <sup>-15</sup>	
			ref rxn is H <sub>2</sub> + H <sub>2</sub> → H <sub>2</sub> θ + H with k <sub>ref</sub> = 7 x 10 <sup>-15</sup>	
	H <sub>2</sub> + C <sub>2</sub> H <sub>2</sub> → products			
	Smith, Zellner (1973)	210-460	2.0 x 10 <sup>-12</sup> exp(-250/T)	
	Pastrana, Carr (1974)	300	2.0 ± 0.6 x 10 <sup>-13</sup>	
	Davis, et al (1975)	300	1.65 ± 0.15 x 10 <sup>-13</sup>	
	H <sub>2</sub> + C <sub>2</sub> H <sub>4</sub> → products			
	Atkinson, Perry, Pitts (1977)	299-425	2.2 x 10 <sup>-12</sup> exp(385 ± 150/T)	(a)
			(a) High pressure values (225-660 torr Ar) Values also given for 25 and 75 torr pressure dependent values given over range 1 to 7 torr He 1.0 ± 0.2 x 10 <sup>-11</sup>	(b)
19,57	Howard (1976)	296		
	Overend, Paraskevopoulos (1977)	296		
			(b) Authors' suggested limiting high pressure value based on expts at 400 torr SF <sub>6</sub> and CF <sub>4</sub> and analysis of dependence of k on [H <sub>2</sub> θ] at low values of [H <sub>2</sub> θ]	
	Stuhl (1973c)	298	3 x 10 <sup>-12</sup>	±0.12
	Smith, Zellner (1973)	210-460	7.5 x 10 <sup>-12</sup> exp(-110/T)	
	Davis, et al (1975)	300	10 <sup>-12</sup> x k P(He)/torr	
			2.24 2.78 3.63 4.06 4.72 5.33	3 5 10 20 100 300
		300	3.64 x 10 <sup>-12</sup> at 3 torr N <sub>2</sub>	
		300	1.8 x 10 <sup>-12</sup>	
		300	1.7 ± 0.5 x 10 <sup>-12</sup>	
	300-500	1.26 x 10 <sup>-12</sup> exp(454/T) at 100 torr He		
	381	6.2 x 10 <sup>-12</sup> (1 atm H <sub>2</sub> θ vapor)		
	416	7.3 x 10 <sup>-12</sup> (1 atm H <sub>2</sub> θ vapor)		

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^2\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of $\log k$
19,58	$\text{H}\theta + \text{C}_2\text{H}_6 \rightarrow \text{products}$ Overend, et al (1975) Greiner (1970)	300 300-500 296	$2.64 \pm 0.17 \times 10^{-13}$ $1.86 \times 10^{-11} \exp(-1230/T)$ $(2.9 \pm 0.6) \times 10^{-13}$	
19,59	$\text{H}\theta + \text{C}_3\text{H}_8 \rightarrow \text{products}$ Atkinson, Pitts (1975b) Stuhl (1973c) Morris, et al (1971) Bradley, et al (1973) Gorse, Volman (1974)	297-425 298 300 300 300	$4.1 \times 10^{-12} \exp(540 \pm 150/T)$ $1.45 \times 10^{-11}$ $1.7 \times 10^{-11}$ $5.0 \pm 1.7 \times 10^{-12}$ $k/k(\text{C}\theta + \text{H}\theta) = 89.3$ (low pressure) $k = 1.2 \times 10^{-11}$	$\pm 0.06$     (a)
19,60	$\text{H}\theta + \text{n-C}_4\text{H}_{10} \rightarrow \text{products}$ Davis (1976) Gordon, Mulac (1975) Stuhl (1973b) Perry, Atkinson, Pitts (1976b) Greiner (1970) Gorse, Volman (1974)	300 381 416 300 297-420 300-500 300	$k(\text{C}\theta + \text{H}\theta) = 1.4 \times 10^{-13}$ , this survey $1.53 \pm 0.06 \times 10^{-11}$ $1.4 \times 10^{-11}$ (1 atm $\text{H}_2\theta$ vapor) $2.0 \times 10^{-11}$ (1 atm $\text{H}_2\theta$ vapor)  $2.35 \times 10^{-12}$ $1.76 \times 10^{-11} \exp(-560/T)$ $1.41 \times 10^{-11} \exp(-524/T)$ $k/k(\text{C}\theta + \text{H}\theta) = 19.4$ (low pressure) $k = 2.7 \times 10^{-12}$	          (a)
19,60	$\text{H}\theta + \text{isobutane} \rightarrow \text{products}$ Greiner (1970) Gorse, Volman (1974)	300-500 300	$k(\text{C}\theta + \text{H}\theta) = 1.4 \times 10^{-13}$ , this survey  $8.7 \times 10^{-12} \exp(-387/T)$ $k/k(\text{H}\theta + \text{C}\theta) = 23.5$ (low pressure) $k = 3.3 \times 10^{-12}$	   (a)
19,60	$\text{H}\theta + \text{alkane} \rightarrow \text{H}_2\theta + \text{alkyl radical}$ Greiner (1970)	300	$k(\text{C}\theta + \text{H}\theta) = 1.4 \times 10^{-13}$ , this survey $1.6 \times 10^{-12}$ (b) Determined in expt. at 100 torr relative to $k(\text{H}\theta + \text{C}\theta) = 1.5 \times 10^{-13}$	(b)
19,60a	$\text{H}\theta + \text{C}_6\text{H}_6 \rightarrow \text{products}$ Perry, Atkinson, Pitts (1977)	298 380-470 298	$k = [1.0 \exp(-820/T) N_p + 2.3 \exp(-430/T) N_s + 2.1 \exp(+95/T) N_t] \times 10^{-12}$ Where $N_p$ , $N_s$ , and $N_t$ are the number of primary secondary, and tertiary hydrogen atoms respectively. Do not use formula for $\text{CH}_4$ and $\text{C}_2\text{H}_6$ .  $(1.20 \pm 0.15) \times 10^{-12}$ $k_1 = 4 \times 10^{-11} \exp(-2000/T)$ $k_1/k = 0.05$	total rxn; $P = 100$ torr Ar (a) (b)
		298	(a) $k_1$ refers to abstraction rxn (b) Fraction of rxn proceeding by abstraction; expression for $k_1$ extrapolated to 298K  $(1.24 \pm 0.12) \times 10^{-12}$ total rxn; $P = 50-600$ torr Ar	

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$		Notes and Reliability of $\log k$
19,60a	Davis, Bollinger, Fischer (1975)	298	$0.85 \times 10^{-12}$	$P = 3 \text{ torr He}$	total rxn
			$1.36 \times 10^{-12}$	$P = 20 \text{ torr He}$	total rxn
			$1.59 \times 10^{-12}$	$P = 100 \text{ torr He}$	total rxn
	$\text{H}\theta + \text{C}_6\text{H}_5\text{CH}_3 \rightarrow \text{products}$ Perry, Atkinson, Pitts (1977)	298 380-470 298	$(6.40 \pm 0.64) \times 10^{-12}$ $k_1 = 5 \times 10^{-12} \exp(-450/T)$ $k_1/k = 0.16$ (a) $k_1$ refers to H atom abstraction (b) Fraction of rxn proceeding by abstraction; expression for $k_1$ extrapolated to 298K	total rxn; $P = 200 \text{ torr Ar}$ (a) (b)	
19,61	Hansen, Atkinson, Pitts (1975)	298	$(5.78 \pm 0.58) \times 10^{-12}$	total rxn; $P = 100-600 \text{ torr Ar}$	total rxn
	Davis, Bollinger, Fischer (1975)	298	$3.60 \times 10^{-12}$	$P = 3 \text{ torr He}$	total rxn
			$5.00 \times 10^{-12}$	$P = 20 \text{ torr He}$	total rxn
	$\text{H}\theta + \text{CF}_2\text{Cl}_2 \rightarrow \text{products}$ +NASA (1977) eval Atkinson, Hansen, Pitts (1975) Howard, Evenson (1976a) Chang, Kaufman (1977)	200-300 297-424 296 480	$k < 1 \times 10^{-12} \exp(-3560/T)$ $k < 1 \times 10^{-15}$ $k < 4 \times 10^{-16}$ $k < 6 \times 10^{-16}$ (a) upper limit only (b) Expression based on upper limit at 480K reported by Chang and Kaufman. "A factor" chosen; E/R value derived	(a,b) (a) (a) (a)	
19,62	$\text{H}\theta + \text{CFCl}_3 \rightarrow \text{products}$ +NASA (1977) eval Atkinson, Hansen, Pitts (1975) Howard, Evenson (1976a) Chang, Kaufman (1977)	200-300 297-424 296 480	$k < 1 \times 10^{-12} \exp(-3650/T)$ $k < 1 \times 10^{-15}$ $k < 5 \times 10^{-16}$ $k < 5 \times 10^{-16}$ (a) upper limit only (b) Expression based on upper limit at 480K reported by Chang and Kaufman. "A factor" chosen; E/R value derived	(a,b) (a) (a) (a)	
19,63	$\text{H}\theta + \text{CCl}_4 \rightarrow \text{products}$ Howard, Evenson (1976a)	296	$k < 4 \times 10^{-15}$	upper limit only	
19,64	$\text{H}\theta + \text{CH}_2\text{Cl} \rightarrow \text{H}_2\theta + \text{CH}_2\text{Cl}$ +NASA (1977) eval Davis, et al (1976) Perry, Atkinson, Pitts (1976) Howard, Evenson (1976a)	200-300 250-350 298-423 296	$2.2 \times 10^{-12} \exp(-1142/T)$ $(1.84 \pm 0.18) \times 10^{-12} \exp(-1098 \pm 35/T)$ $4.1 \times 10^{-12} \exp(-1359 \pm 150/T)$ $(3.6 \pm 0.8) \times 10^{-14}$	$\pm 0.2 \text{ at } 230\text{K}$	
	$\text{H}\theta + \text{CH}_2\text{Cl}_2 \rightarrow \text{H}_2\theta + \text{CHCl}_2$ +NASA (1977) eval Davis, et al (1976) Perry, Atkinson, Pitts (1976) Howard, Evenson (1976)	200-300 245-375 298 296	$5.2 \times 10^{-12} \exp(-1094/T)$ $(4.27 \pm 0.63) \times 10^{-12} \exp(-1094 \pm 81/T)$ $(1.45 \pm 0.20) \times 10^{-13}$ $(1.55 \pm 0.14) \times 10^{-13}$	$\pm 0.2 \text{ at } 230\text{K}$	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
19,64	H $\theta$ + CHCl <sub>3</sub> → H <sub>2</sub> $\theta$ + CCl <sub>3</sub> †Davis, et al (1976)	245-375	(4.69 ± 0.71) x 10 <sup>-12</sup> exp(-1134 ± 108/T)	±0.2 at 230K
19,65	Howard, Evenson (1976a) H $\theta$ + CHFCl <sub>2</sub> → H <sub>2</sub> $\theta$ + CFCl <sub>2</sub> †NASA (1977) eval	296	1.01 ± 0.15 x 10 <sup>-13</sup>	
19,65	Perry, Atkinson, Pitts (1976) Chang, Kaufman (1977) Watson, et al (1977) Howard, Evenson (1976a) H $\theta$ + CHF <sub>2</sub> Cl → H <sub>2</sub> $\theta$ + CF <sub>2</sub> Cl	200-300 298-422 241-396 245-375 296	1.3 x 10 <sup>-12</sup> exp(-1127/T) 1.75 x 10 <sup>-12</sup> exp(-1253 ± 150/T) (1.16 ± 0.17) x 10 <sup>-12</sup> exp(-1073 ± 40/T) (1.87 ± 0.2) x 10 <sup>-12</sup> exp(-1245 ± 26/T) 2.6 x 10 <sup>-14</sup>	±0.2 at 230K
19,65	†NASA (1977) eval Atkinson, Hansen, Pitts (1975) Chang, Kaufman (1977) Watson, et al (1977) Howard, Evenson (1976a) H $\theta$ + CH <sub>2</sub> FCl → H <sub>2</sub> $\theta$ + CHFCl	200-300 297-434 253-427 250-350 296	1.2 x 10 <sup>-12</sup> exp(-1660/T) 1.21 x 10 <sup>-12</sup> exp(-1636 ± 150/T) (1.20 ± 0.16) x 10 <sup>-12</sup> exp(-1657 ± 39/T) (9.25 ± 1.0) x 10 <sup>-13</sup> exp(-1575 ± 71/T) 3.4 x 10 <sup>-15</sup>	±0.2
19,65	†Watson, et al (1977) Howard, Evenson (1976a) H $\theta$ + CH <sub>3</sub> F → products	245-375 296	(2.84 ± 0.3) x 10 <sup>-12</sup> exp(-1259 ± 50/T) 3.7 ± 0.6 x 10 <sup>-14</sup>	
19,65	Howard, Evenson (1976a) H $\theta$ + CH <sub>2</sub> F <sub>2</sub> → products	296	16 ± 2 x 10 <sup>-15</sup>	
19,65	Howard, Evenson (1976a) H $\theta$ + CHF <sub>3</sub> → products	296	7.8 ± 1.2 x 10 <sup>-15</sup>	
19,65	Howard, Evenson (1976a) H $\theta$ + CF <sub>4</sub> → products	296	2 x 10 <sup>-16</sup>	
19,65	Howard, Evenson (1976a) H $\theta$ + CF <sub>3</sub> Cl → products	296	k < 4 x 10 <sup>-16</sup>	upper limit only
19,65	Howard, Evenson (1976a) H $\theta$ + CH <sub>3</sub> Br → H <sub>2</sub> $\theta$ + CH <sub>2</sub> Br	296	k < 7 x 10 <sup>-16</sup>	upper limit only
19,65	†Davis, et al (1976) Howard, Evenson (1976a) H $\theta$ + CH <sub>3</sub> CCl <sub>3</sub> → products	245-350 296	(7.93 ± 0.79) x 10 <sup>-13</sup> exp(-889 ± 98/T) 35 ± 8 x 10 <sup>-13</sup>	±0.2 at 230K
19,65	†NASA (1977) eval Watson, et al (1977) Chang, Kaufman (1977) Howard, Evenson (1976b) H $\theta$ + C <sub>2</sub> HCl <sub>3</sub> → products	200-300 260-375 275-405 296	3.5 x 10 <sup>-12</sup> exp(-1562/T) (3.72 ± 0.4) x 10 <sup>-12</sup> exp(-1627 ± 50/T) (1.95 ± 0.24) x 10 <sup>-12</sup> exp(-1333 ± 37/T) 15 ± 3 x 10 <sup>-15</sup>	±0.36 at 230K
19,65	†NASA (1977) eval Chang, Kaufman (1977) Davis (1977) Howard (1976)	200-300 234-420 300 296	2.3 x 10 <sup>-12</sup> (5.32 ± 0.71) x 10 <sup>-13</sup> exp(+445 ± 41/T) 2.35 ± 0.25 x 10 <sup>-12</sup> 2.0 ± 0.4 x 10 <sup>-12</sup>	±0.36 preliminary



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of log k
19,65	$\text{H}\theta + \text{C}_2\text{Cl}_4 \rightarrow \text{products}$ †*Chang, Kaufman (1977) Davis (1977)	297-420 260-375 296	$(9.44 \pm 1.34) \times 10^{-12} \exp(-1199 \pm 55/T)$ $(1.06 \pm 0.5) \times 10^{-11} \exp(-1300 \pm 150/T)$ $1.70 \times 0.34 \times 10^{-13}$	$\pm 0.36$ at 230K preliminary
19,65	$\text{H}\theta + \text{CH}_3\text{CF}_2\text{Cl} \rightarrow \text{products}$ Howard (1976)			
19,65	Watson, et al (1977) Howard, Evenson (1976b) $\text{H}\theta + \text{CF}_2\text{ClCFCl}_2 \rightarrow \text{products}$	273-375 296	$(1.15 \pm 0.15) \times 10^{-12} \exp(-1748 \pm 30/T)$ $2.8 \pm 0.4 \times 10^{-15}$	
20,M	Watson, et al (1977) $\text{H}\theta_2 + \text{M} \rightarrow \text{H} + \theta_2 + \text{M}$	298	$k < 3 \times 10^{-16}$ see reverse reaction	upper limit only
20,M	$\text{H}\theta_2 + \text{M} \rightarrow \text{H}\theta + \theta + \text{M}$		see reverse reaction	
20,20	$\text{H}\theta_2 + \text{H}\theta_2 \rightarrow \text{H}_2\theta_2 + \theta_2$ †*Hamilton, Lil (1977)	298	$-d[\text{H}\theta_2]/dt = 2k[\text{H}\theta_2]^2$ $2.5 \times 10^{-12}$ (a) Superseded result reported in Hamilton (1975)	$\pm 0.3$ (a)
20,21	Baulch, et al (1972) review Lloyd (1974) review $\text{H}\theta_2 + \text{H}_2 \rightarrow \text{H}_2\theta_2 + \text{H}$	300 300-1000	$3.3 \times 10^{-12}$ $1.7 \times 10^{-11} \exp(-500/T)$ Temperature coefficient	$\pm 0.3$ at 300K larger at higher T
20,22	$\text{H}\theta_2 + \text{H}_2\theta \rightarrow \text{H}_2\theta_2 + \text{H}\theta$ Baulch, et al (1972) review Lloyd (1974) review	300-800 300-1000	$1.2 \times 10^{-12} \exp(-9400/T)$ same recommendation. $k_f = k_r K_{\text{eq}}$	$\pm 0.3$
20,28	$\text{H}\theta_2 + \text{S}\theta_2 \rightarrow \text{S}\theta_3 + \text{H}\theta$ *Payne, Stief, Davis (1973)	300-800 300-1000	$4.7 \times 10^{-11} \exp(-16,500/T)$ same recommendation $k_f = k_r K_{\text{eq}}$	$\pm 0.2$
20,35	$\text{H}\theta_2 + \text{Cl} \rightarrow \text{HCl} + \theta_2$ †*Leu, DeKore (1976)	300	$9 \times 10^{-16} \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ (a) Relative rate measurement. Value based on $k(\text{H}\theta_2 + \text{H}\theta_2)$ this survey.	$\pm 0.3$ (a)
20,35Br	$\text{H}\theta_2 + \text{Br} \rightarrow \text{HBr} + \theta_2$ †*NASA (1977) eval	200-300	$3 \times 10^{-11}$ (a) estimate - no data	$\pm 0.6$ at 230K $\pm 1.5$ at 230K (a)
20,36	$\text{H}\theta_2 + \text{Cl}\theta \rightarrow \text{H}\theta\text{Cl} + \theta_2$ †*NASA (1977) eval	200-300	$2 \times 10^{-13}$ (a) No data; value assumed to be intermediate between $k(\text{H}\theta_2 + \text{H}\theta_2)$ and $k(\text{Cl}\theta + \text{Cl}\theta)$	$\pm 1$ at 230K (a)
20,45	$\text{H}\theta_2 + \text{C}\theta \rightarrow \text{C}\theta_2 + \text{H}\theta$ *This survey Baulch, et al (1976) review	300 700-1000	$2.5 \times 10^{-10} \exp(-11900/T)$	$\pm 0.5$ a



No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	Notes and Reliability of log k	
				log k	
20,48	Lloyd (1974) eval.	300-1000	$1.7 \times 10^{-13} \exp(-5000/T)$	b	
	Westenberg, deHaas (1972a)	300	$\sim 1 \times 10^{-12}$	c,d	
	Davis, Payne, Stief (1973)	300	$< 10^{-20}$	e	
	Wyrach, et al (1974)	300	$< 3 \times 10^{-18}$	e	
	Simonaitis, Heicklen (1973a)	373-473	$< 5 \times 10^{-18}$	e	
20,51	Volman, Gorse (1972)	330	$< 10^{-15}$	f	
	Baldwin, Walker, Webster (1970)	713-773	$1 \times 10^{-10} \exp(-11500/T)$	e	
	Vardanyan, Dangyan, Sachyan (1972)	878-952	$2.2 \times 10^{-10} \exp(-11500/T)$	e	
NOTE: a. The low value of k selected in this evaluation is based on the evaluation by Lloyd and the measurements by Davis, Payne, Stief and Simonaitis, Heicklen.					
b. Based on Davis, Payne and Stief (1972) and high T results. Temperature coef. probably maximum value.					
c. Indirect measurement using a low-pressure discharge flow system.					
d. Relative rate measurement = reference reaction: $\text{H} + \text{H}\theta_2 \rightarrow \text{H}\theta + \text{H}\theta$					
e. Relative rate measurement = reference reaction: $\text{H}\theta_2 + \text{H}\theta_2 \rightarrow \text{H}_2\theta_2 + \theta_2$					
f. Relative rate measurement = reference reaction: $\text{H}\theta + \text{C}\theta \rightarrow \text{C}\theta_2 + \text{H}$					
20,57	$\text{H}\theta_2 + \text{CH}_2\theta \rightarrow \text{H}_2\theta_2 + \text{CH}\theta$	300-800	$1.7 \times 10^{-12} \exp(-4000/T)$		$> \pm 1$ at 300K, $\pm 0.2$ at 800K
	*Lloyd (1974) review	773	$1.6 \times 10^{-15}$	(a)	
(a) Derived by computer fit to data. Based on $k(\text{H}\theta_2 + \text{H}\theta_2 \rightarrow \text{H}_2\theta_2 + \theta_2) = 3.0 \times 10^{-12} \text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$					
20,58	$\text{H}\theta_2 + \text{CH}_3\theta_2 \rightarrow \text{CH}_3\theta_2\text{H} + \theta_2$	300	$6.7 \times 10^{-14}$		estimate
	Demerjian et al (1972) review	773	$2.5 \times 10^{-17}$	(a)	
(a) Relative to $k(\text{H}\theta_2 + \text{HCH}\theta) = 1.6 \times 10^{-15}$					
20,59	$\text{H}\theta_2 + \text{C}_2\text{H}_4 \rightarrow \text{addition products}$	300	$\sim 1.7 \times 10^{-17}$		$> \pm 1$ (a)
	Lloyd (1974) review	300-1000		(a) Suggestion. Data are irreconcilable.	
20,60	$\text{H}\theta_2 + \text{C}_2\text{H}_6 \rightarrow \text{H}_2\theta_2 + \text{C}_2\text{H}_5$	300-1000	$\sim 1.7 \times 10^{-12} \exp(-7000/T)$		$\pm 1$ (a)
	Lloyd (1974) review	300-1000		(a) Relative rate data versus $2\text{H}\theta_2 \rightarrow \text{H}_2\theta_2 + \theta_2$ and $\text{H}\theta_2 + \text{C}\theta \rightarrow \text{H}\theta + \text{C}\theta_2$ . Temp. coef. estimated.	
20,61	$\text{H}\theta_2 + \text{C}_3\text{H}_8 \rightarrow \text{H}_2\theta_2 + 1-\text{C}_3\text{H}_7$	300-1000	$< 3.3 \times 10^{-13} \exp(-5300/T)$		$\pm 1$ (a)
	Lloyd (1974) review	300-1000			

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of log k																																													
20,60	$\text{H}\theta_2 + n\text{-C}_4\text{H}_{10} \rightarrow \text{H}_2\theta_2 + s\text{-C}_4\text{H}_9$ (primarily) Lloyd (1974) review	300-1000	(a) Based on upper limit meas. Temp. coef. estimated. $8.7 \times 10^{-13} \exp(-5300/T)$ (a) Based on upper limit meas. Temp. coef. estimated.	$\pm 1$ (a)																																													
20,60	$\text{H}\theta_2 + i\text{-C}_4\text{H}_{10} \rightarrow \text{H}_2\theta_2 + t\text{-C}_4\text{H}_9$ Lloyd (1974) review	300-1000	$1.7 \times 10^{-13} \exp(-3500/T)$ (a) Based on upper limit meas. Temp. coef. estimated.	$\pm 1$ (a)																																													
20,60	$\text{H}\theta_2 + \text{alkane} \rightarrow \text{H}_2\theta_2 + \text{alkyl radical}$ Walker (1977)	300-800	$k = 8 \times 10^{-14} [N_p \exp(-7520/T) + N_s \exp(-6330/T)] + N_t \exp(-4990/T)$ where $N_p$ , $N_s$ , and $N_t$ are the number of primary, secondary, and tertiary hydrogen atoms respectively see reverse reaction																																														
21,M 21,35	$\text{H}_2 + \text{M} \rightarrow \text{H} + \text{H} + \text{M}$ $\text{H}_2 + \text{Cl} \rightarrow \text{HCl} + \text{H}$ † Watson (1977) eval Watson, et al (1977) Lee, et al (1977)	200-300 213-350 200-500	$3.5 \times 10^{-11} \exp(-2290/T)$ $(5.5 \pm 0.5) \times 10^{-11} \exp(-2375 \pm 100/T)$ $(2.66 \pm 0.42) \times 10^{-11} \exp(-2230 \pm 60/T)$	$\pm 0.12$ at 230K																																													
21,36	$\text{H}_2 + \text{Cl}\theta \rightarrow \text{products}$ † NASA (1977) eval	200-300	$k \leq 1 \times 10^{-12} \exp(-4800/T)$ (a) "A factor" chosen. $F_{\text{act}}$ fitted to upper limit at $T = 670\text{K}$ reported in Watson's review (1977)	$\pm 2$ at 270K (a)																																													
22,29	$\text{H}_2\theta + \text{S}\theta_3 \rightarrow \text{products}$ Castleman, et al (1975)	298	$9.1 \pm 2.9 \times 10^{-13}$																																														
23,hv	$\text{H}_2\theta_2 + h\nu \rightarrow \text{H}\theta + \text{H}\theta$ DeMore (1977)		<table> <tr> <th><math>\Delta/nm</math></th><th><math>10^{20} \frac{x}{\sigma}</math></th><th><math>\Delta/nm</math></th><th><math>10^{20} \frac{x}{\sigma}</math></th><th>(a)</th></tr> <tr><td>200</td><td>48.3</td><td>280</td><td>2.1</td><td></td></tr> <tr><td>210</td><td>36.3</td><td>290</td><td>1.14</td><td></td></tr> <tr><td>220</td><td>26.4</td><td>300</td><td>0.66</td><td></td></tr> <tr><td>230</td><td>18.7</td><td>310</td><td>0.38</td><td></td></tr> <tr><td>240</td><td>12.9</td><td>320</td><td>0.21</td><td></td></tr> <tr><td>250</td><td>8.9</td><td>330</td><td>0.13</td><td></td></tr> <tr><td>260</td><td>5.5</td><td>340</td><td>0.08</td><td></td></tr> <tr><td>270</td><td>3.4</td><td>350</td><td>0.04</td><td></td></tr> </table> <p>(a) <math>\sigma</math> is the absorption cross section in units of <math>\text{cm}^2\text{molecule}^{-1}</math> (base e). These values by DeMore (private communication) are in good agreement (within 10%) with unpublished results by Molina, et al and with results of Holt, et al (1948) over common ranges of measurements</p>	$\Delta/nm$	$10^{20} \frac{x}{\sigma}$	$\Delta/nm$	$10^{20} \frac{x}{\sigma}$	(a)	200	48.3	280	2.1		210	36.3	290	1.14		220	26.4	300	0.66		230	18.7	310	0.38		240	12.9	320	0.21		250	8.9	330	0.13		260	5.5	340	0.08		270	3.4	350	0.04		
$\Delta/nm$	$10^{20} \frac{x}{\sigma}$	$\Delta/nm$	$10^{20} \frac{x}{\sigma}$	(a)																																													
200	48.3	280	2.1																																														
210	36.3	290	1.14																																														
220	26.4	300	0.66																																														
230	18.7	310	0.38																																														
240	12.9	320	0.21																																														
250	8.9	330	0.13																																														
260	5.5	340	0.08																																														
270	3.4	350	0.04																																														
<p>Hampson (1973) review</p> <p>Quantum yields (primary)</p> <p><math>\phi(-\text{H}_2\theta_2) = 1.0 \quad \lambda &gt; 200\text{nm}</math></p>																																																	

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of log k
23, M	$\text{H}_2\text{O}_2 + \text{M} \rightarrow \text{H}_2\text{O} + \text{HO} + \text{M}$		see reverse equation	
23, 35	$\text{H}_2\text{O}_2 + \text{Cl} \rightarrow \text{HCl} + \text{HO}_2$	265-400	$1.7 \times 10^{-12} \exp(-384/T)$	$\pm 0.36$ at 230K
23, 35Br	+*Watson (1977) review $\text{H}_2\text{O}_2 + \text{Br} \rightarrow \text{HBr} + \text{HO}_2$	200-300	$5 \times 10^{-12} \exp(-1570/T)$	$\pm 0.6$ at 270K
24, M	+*NASA (1977) eval $\text{HN}_3 + \text{M} \rightarrow \text{H} + \text{N}_2 + \text{M}$	230-700	$5 \times 10^{-8} \exp(-24,500/T)$ $\text{M} = \text{H}_2$ (a) No data. Value based on reverse rate	$\pm 0.2$ (a)
24, 24	$\text{HN}_3 + \text{HN}_3 \rightarrow \text{H}_2 + \text{N}_2 + \text{N}_2$	300	$4 \times 10^{-15}$	$\pm 0.3$
25, hv	*Baulch, et al (1973) review $\text{HN}_3 + \text{hv} \rightarrow \text{H} + \text{N}_2$ (a) $\rightarrow \text{H} + \text{N}_2$ (b) Molina (1977) review	300	table of absorption cross sections for wavelength range 200-400nm. (a) Gives corrected values which supersede results in Johnston, Graham (1974). Also gives results of Cox and Derwent (1976)	(a)
	Cox, Derwent (1976)	300	table of absorption cross sections for $\lambda = 200-400\text{nm}$	
	Cox (1974b)		$\phi_a \geq 0.97$	(a)
26, hv	$\text{HN}_3 + \text{hv} \rightarrow \text{H} + \text{N}_2$ *Johnston (1974) review	217 217 222 227 237 251 265	Elevation/km $k(\text{noon})/\text{s}^{-1}$ $k(24 \text{ hr. av.})/\text{s}^{-1}$ 15 $5.1 \times 10^{-7}$ $1.5 \times 10^{-7}$ 20 $7.7 \times 10^{-7}$ $2.3 \times 10^{-7}$ 25 $2.8 \times 10^{-6}$ $6.8 \times 10^{-7}$ 30 $1.4 \times 10^{-5}$ $3.5 \times 10^{-6}$ 35 $4.1 \times 10^{-5}$ $1.2 \times 10^{-5}$ 40 $7.6 \times 10^{-5}$ $2.7 \times 10^{-5}$ 45 $1.1 \times 10^{-4}$ $4.2 \times 10^{-5}$ Rates are for 45° latitude solar equinox standard ozone background	$\pm 0.15$
	*Johnston, Graham (1973)		Table of absorption coefficients for $\lambda = 190-325\text{nm}$	
	Johnston, et al (1974)		$\phi = 1$ $200 < \lambda < 315\text{nm}$	(a)
	Blaume (1973)		(a) Products probably those shown above. Absorption cross section data for for $185 < \lambda < 325 \text{ nm}$ in good agreement with that of Johnston and Graham (1973). figure showing extinction coefficient vs $\lambda$ for $\lambda = 120-170\text{nm}$ see reverse reaction	
26, M	$\text{HN}_3 + \text{M} \rightarrow \text{H} + \text{N}_2 + \text{M}$	200-300	$1.0 \times 10^{-11} \exp(-2170/T)$	$\pm 0.6$ at 230K (a)
26, 35	$\text{HN}_3 + \text{Cl} \rightarrow \text{HCl} + \text{N}_2$ +*NASA (1977) eval Leu, DeMore (1976)	295	$(6.8 \pm 3.4) \times 10^{-15}$ (a) "A factor" chosen. Eact derived to fit room temp. result of Leu, DeMore.	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
226a, 34b	S + CS <sub>2</sub> → S <sub>2</sub> + CS	298	6.5 x 10 <sup>-13</sup>	±0.2
26a, 34c	Baulch, et al (1976) review S + COS → CS + S <sub>2</sub>	230-2600	2.8 x 10 <sup>-12</sup> exp(-2050/T)	±0.5
27, 27	Baulch, et al (1976) review S $\dot{O}$ + S $\dot{O}$ → S $\dot{O}_2$ + S or (S $\dot{O}$ ) <sub>2</sub>	300 1000 300	< 3 x 10 <sup>-15</sup> < 2 x 10 <sup>-13</sup> 8.3 ± 6.7 x 10 <sup>-16</sup>	
27, 29	Schofield (1973) review Chung, Calvert, Bottenheim (1975) S $\dot{O}$ + S $\dot{O}_3$ → 2S $\dot{O}_2$	300	2 ± 1.2 x 10 <sup>-15</sup>	
28, 49M	Chung, Calvert, Bottenheim (1975) S $\dot{O}_2$ + CH <sub>3</sub> (•M) → CH <sub>3</sub> S $\dot{O}_2$ (•M)	300	3 x 10 <sup>-13</sup> cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	±0.1
31, 31	James, et al (1973) HS + HS → H <sub>2</sub> S + S	295	1.2 x 10 <sup>-11</sup>	±0.2
35, 35M	Baulch, et al (1976) review Cl + Cl + M → Cl <sub>2</sub> + M	200-500	6 x 10 <sup>-34</sup> exp(900 ± 250/T) cm <sup>6</sup> molecule <sup>-2</sup> s <sup>-1</sup> M = Ar	±0.1
35, 37	*Watson (1977) review Cl + OCl $\dot{O}$ → 2Cl $\dot{O}$	300-588	(5.9 ± 0.9) x 10 <sup>-11</sup> exp(0 ± 120/T)	
35, 37	*Watson (1977) review Cl + Cl $\dot{O}$ → Cl <sub>2</sub> + $\dot{O}_2$ (a) → 2Cl $\dot{O}$ (b)		k <sub>a</sub> = 1.6 x 10 <sup>-10</sup> k <sub>b</sub> = 1.1 x 10 <sup>-11</sup> k <sub>a</sub> = 1.56 x 10 <sup>-10</sup> k <sub>a</sub> /k <sub>b</sub> = 108 k <sub>a</sub> /k <sub>b</sub> = 15	
35, 44a	*NASA (1977) eval Johnston, et al (1969)	298	6.8 x 10 <sup>-13</sup>	
35, 44a	Nicholas, Norrish (1968) Cl + Cl <sub>2</sub> $\dot{O}$ → Cl <sub>2</sub> + Cl $\dot{O}$	258	3 ± 0.5 x 10 <sup>-11</sup>	
35, 42	*Watson (1977) review Cl + NOCl → NO + Cl <sub>2</sub>	300	k >> 3 x 10 <sup>-14</sup>	
35, 43	*Watson (1977) review Cl + NO <sub>2</sub> Cl → NO <sub>2</sub> + Cl <sub>2</sub>	258	1.68 x 10 <sup>-12</sup> exp(-607/T)	±0.3
35, 43a	Watson (1977) review Cl + NO <sub>3</sub> Cl → products	298	7.3 x 10 <sup>-12</sup> exp(-1269/T)	(a)
35, 56	*Kurylo, Manning (1977) Cl + CB <sub>4</sub> → HCl + CH <sub>3</sub>	224-273	(7.93 ± 1.53) x 10 <sup>-12</sup> exp(-1273 ± 51/T)	(b)
35, 56	*Watson (1977) review Manning, Kurylo (1977)	200-300	(7.94 ± 0.7) x 10 <sup>-12</sup> exp(-1260 ± 35/T)	
35, 56	Watson, et al (1976)	218-401	5.44 x 10 <sup>-19</sup> T <sup>2.50</sup> exp(-608/T)	
35, 56	Whytock, et al (1977)	200-299	(6.51 ± 0.79) x 10 <sup>-12</sup> exp(-1229 ± 27/T)	
35, 56	Zahniser, et al (1978)	200-500	8.6 x 10 <sup>-18</sup> T <sup>2.11</sup> exp(-795/T)	
35, 56	Davis, et al (1970)	200-300	(8.2 ± 0.6) x 10 <sup>-12</sup> exp(-1320 ± 20/T)	
35, 56		300	1.5 ± 0.1 x 10 <sup>-13</sup>	

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
	Fettis, Knox (1964) Clyne, Walker (1973)	300-686	4 x 10 <sup>-11</sup> exp(-1930/T) 5.1 x 10 <sup>-11</sup> exp(-1790/T) (a) Based on four direct 1976, 1977 studies listed immediately below. Uncertainty in log k(230K) is ±0.12 and -0.36 (b) Value changed subsequent to publication to allow for effect of small C <sub>2</sub> H <sub>6</sub> impurity see Watson's review (1977)	
35,58	Cl + C <sub>2</sub> H <sub>6</sub> → HCl + C <sub>2</sub> H <sub>5</sub> Manning, Kurylo (1977) Cl + hv → Cl + θ *Watson (1977) review	22-322	(7.29 ± 1.23) x 10 <sup>-11</sup> exp(-60 ± 44/T) absorption cross section data 225 < λ < 300 nm	
36,36	Clθ + Clθ → Cl + Clθθ (a) → Cl <sub>2</sub> + θ <sub>2</sub> (b) → θClθ + Cl (c) †*NASA (1977) eval	200-300	k <sub>a</sub> = 1 x 10 <sup>-12</sup> exp(-1238/T) k <sub>b</sub> = 5 x 10 <sup>-13</sup> exp(-1238/T) k <sub>c</sub> = 2 x 10 <sup>-12</sup> exp(-2200/T) (a) -d[Clθ]/dt = k[Clθ] <sup>2</sup> . Low pressure. See Watson's review (1977) for high pressure rec.	±0.2 at 230K (a) ±0.2 at 230K ±1 at 230K
36,36Br	Clθ + Brθ → Br + θClθ (a) → Br + Clθθ (b) †*Watson (1977) review			
36,45	Clθ + Oθ → Oθ <sub>2</sub> + Cl †*NASA (1977) eval	200-300	k <sub>a</sub> = (6.7 ± 1.0) x 10 <sup>-12</sup> k <sub>b</sub> = (6.7 ± 1.0) x 10 <sup>-12</sup>	±0.36 at 230K ±0.36 at 230K
36,56	Clθ + CH <sub>4</sub> → products †*NASA (1977) eval	200-300	k ≤ 1 x 10 <sup>-12</sup> exp(-3700/T) (a) "A factor" chosen. E <sub>act</sub> fitted to upper limit at T = 587K reported in Watson's review (1977)	±2 at 230K (a)
36,57	Clθ + C <sub>2</sub> H <sub>4</sub> → products Walker (1972)	200-300	k ≤ 1 x 10 <sup>-12</sup> exp(-3700/T) (a) "A factor" chosen. E <sub>act</sub> fitted to upper limit at T = 670K reported in Watson's review (1977)	±2 at 230K (a)
36,59	Clθ + C <sub>2</sub> H <sub>2</sub> → products Walker (1972)	670	k < 5 x 10 <sup>-16</sup> (a) As quoted by Watson (1977) review.	(a)
36Br, 36Br	Brθ + Brθ → 2Br + θ <sub>2</sub> †*NASA (1977) eval Clyne, Watson (1975) Clyne, Cruse (1970)	200-300 298 293	2.9 x 10 <sup>-11</sup> exp(-450/T) 6.4 x 10 <sup>-12</sup> 5.3 x 10 <sup>-11</sup> exp(-450/T) (a) As quoted by Watson (1977) review.	±0.36 at 230K



No.

37, M	ClO <sub>2</sub> + M → Cl + O <sub>2</sub> + M Watson (1977) review		Table of values of rate constant as a function of altitude.
37, hv	ClO <sub>2</sub> + hv → ClO + O		absorption cross section data 225 < λ < 280 nm
37, hv	ClO <sub>2</sub> + hv → ClO + O		Abs. cross section data 150 < λ < 480nm
39, hv	HCl + hv → H + Cl Watson (1977) review		absorption cross section data 140 < λ < 220 nm
40, hv	HCl + hv → products Watson (1977) review		Abs. cross section data 290 < λ < 450nm
42, hv	NOCl + hv Molina (1977) review		Table of absorption cross section values for λ = 190-400nm
43, hv	ClNO <sub>2</sub> + hv Molina (1977) review		Absorption cross section data for λ = 190-400nm
43, hv	ClONO <sub>2</sub> + hv Molina (1977)		Absorption cross section data for λ = 235-400nm
43a, hv	NO <sub>3</sub> Cl + hv Rowland, Spencer, Molina (1976)		Table of absorption cross section values for λ = 186-460nm. Also table of solar photodissociation coefficients Data also tabulated in reviews by Watson (1977) and by Molina (1977). Molina's review also discusses quantum yield data
44, hv	Cl <sub>2</sub> + hv → 2Cl Watson (1977) review		absorption cross section data 240 < λ < 450 nm see CH <sub>3</sub> O + CO
45, 50	CO + CH <sub>3</sub> → . . .		
48, hv	CH <sub>2</sub> O + hv → CH <sub>2</sub> O* → H + HCO (1) → H <sub>2</sub> + CO (2) This survey (1978)		

λ/nm	$\frac{\sigma_1}{\sigma_2}$	$\frac{\sigma_1 + \sigma_2}{0.90}$
285	0.61	1
290	0.74	1
295	0.74	1
300	0.74	1
305	0.74	1
310	0.74	1
315	0.74	1
320	0.60	1
325	0.45	1
330	0.31	1

No.	Reaction/Reference * = Preferred Value † = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant $k/\text{cm}^3\text{molecule}^{-1}\text{s}^{-1}$	Notes and Reliability of $\log k$
335			0.17	1
340			0.02	0.64
Tentative recommendation based principally on results reported by Horowitz and Calvert (1978a) and by Moortgat, et al (1978). Also considers results reported in Clark, Moore and Nogar (1977), Horowitz and Calvert (1978b), Houston and Moore (1976), Lewis and Lee (1977), Lewis, Tang and Lee (1976) and Marling (1977).				
Apparent quantum yield values are slightly higher in presence of $\phi_2$ . Effect is assumed here to be due to secondary reactions.				
	Calvert, et al (1972)		$\frac{\Delta/\text{nm}}{10^{20}} \frac{\sigma}{\text{cm}^2}$	$\frac{\Delta/\text{nm}}{10^{20}} \frac{\sigma}{\text{cm}^2}$ (a)
50.4	$\text{CH}_3\phi + \phi_2 \rightarrow \text{CH}_2\phi + \text{H}\phi_2$ (1)		290 3.18	330 2.36
50.9	$\text{CH}_3\phi + \text{N}\phi \rightarrow \text{CH}_3\phi\text{N}\phi$ (2a) $\rightarrow \text{CH}_2\phi + \text{HN}\phi$ (2b)		300 3.25	340 1.97
50.10	$\text{CH}_3\phi + \text{N}\phi_2 \rightarrow \text{CH}_3\phi\text{N}\phi_2$ (3a) $\rightarrow \text{CH}_2\phi + \text{HN}\phi_2$ (3b)		310 3.14	350 0.84
50.45	$\text{CH}_3\phi + \text{C}\phi \rightarrow \text{products}$ (4) *Heicklen (1973)		320 2.34	360 0.18
(a) $\sigma$ is the value of the absorption cross section averaged for 10nm bands. See McQuigg, Calvert (1969) for figure of of absm. coeff. vs wave length. Also see Miller and Lee (1977) for higher resolution spectra from which absorption cross section values can be derived.				
258			$k_1 \sim 3 \times 10^{-18}$ $\sim 1.6 \times 10^{-13} \exp(-3300/T)$	
298			$k_2 \sim 8 \times 10^{-14}$ $k_1/k_2 = 4.7 \times 10^{-5} \pm 20\%$	ratio probably is smaller at 220K
298			$k_{2b}/k_2 = 0.145 \pm 0.015$	
363			$k_2/k_3 = 1.2 \pm 0.1$	
403			$k_{2a}/k_{3a} = 1.1$ $= 1.8$ $= 2.7$	
258			$k_{3a}/k_3 = 0.9 \pm 0.1$	
258			$k_{3b}/k_3 = 0.1 \pm 0.01$	$k_{3a}/k_{3b}$ probably constant $\approx 100-400K$
298-423			$k_4/k_2 = 5 \times 10^{-4}$	
300-400	Demerjian, et al (1974) review		$k_1 = 1.6 \times 10^{-17}$	

No.	Reaction/Reference * = Preferred Value + = NASA (1977) eval	Temp. Range/K	Reaction Rate Constant k/cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup>	Notes and Reliability of log k
$\sim 4.2 \times 10^{-13} \exp(-3000/T)$ adjusted to fit C <sub>4</sub> H <sub>10</sub> /N <sub>2</sub> simulation studies				
51, 51	Batt, McCulloch, Milne (1975)  Barker, et al (1977) 2CH <sub>3</sub> • → 2CH <sub>3</sub> • + • (a) → CH <sub>3</sub> • + CH <sub>2</sub> • + • (b) → CH <sub>3</sub> • + CH <sub>2</sub> • + • (c)  Weaver, et al (1975)	393-473     396-442	$k_{2a} = 1.7 \times 10^{-13}$ $k_{2b}/k_{2a} = 0.17$ $k_{3a} = 3.3 \times 10^{-13}$ $k_{3b}/k_{3a} = 0.1$ $k_{2a} \sim 3 \times 10^{-11}$ $k_{2b} \sim 7 \times 10^{-12}$ $k_1 = 5 \times 10^{-13} \exp(-2000/T)$	±0.4 ±0.5
54, hv	Parkes (1977)  CH <sub>3</sub> • + hv → CH <sub>3</sub> • + N• (a) → isomer (b) → CH <sub>2</sub> • + H• (c) → CH <sub>2</sub> • + H• + N• (d)  *Heicklen (1973) review	300	$k_a/k = 0.43$ $k_b/k = 0.50$ $k_c/k = 0.07$ $k_a = 1.6 \pm 0.4 \times 10^{-13}$ $k_b + k_c = 3.0 \pm 0.8 \times 10^{-13}$	
61, hv	CF <sub>2</sub> Cl <sub>2</sub> + hv Watson (1977) review	298 298 298	$k_a/k = 0.76 \pm 0.02, \lambda = 366\text{nm}$ $k_b/k = 0.24 \pm 0.04, \lambda = 366\text{nm}$ $(k_c + k_d)/k < 0.02, \lambda = 366\text{nm}$ $k = k_a + k_b + k_c + k_d$	
62, hv	CFCl <sub>3</sub> + hv Molina (1977) review Watson (1977) review		Absorption cross section data for $174 < \lambda/\text{nm} < 220$ Low temperature absorption cross section data	
63, hv	CCl <sub>4</sub> + hv Molina (1977) review Watson (1977) review		Tabulated values of absorption cross sections for $174 < \lambda/\text{nm} < 230$ . Low temperature abs. cross section data	
			Absorption cross section data for $174 < \lambda/\text{nm} < 238$ . see Watson's review (1977)	

## REFERENCES

- Ahumada, J. J., Michael, J. V., and Osborne, D. T., "Pressure Dependence and Third Body Effects on The Rate Constants for  $H + O_2$ ,  $H + NO$ , and  $H + CO$ ," J. Chem. Phys. 57, 3736-3745 (1972)
- Albers, E. A., Hoyermann, K., Wagner, H. Gg., and Wolfrum J., "Study of the Reaction of Ammonia with Oxygen Atoms," Symp. Combust. 12th (Combustion Institute, Pittsburgh, 1969) 313-321
- Albers, E. A., Hoyermann, K., Wagner, H. Gg., and Wolfrum J., "Absolute Measurements of Rate Coefficients for the Reactions of H and O Atoms with  $H_2O_2$  and  $H_2O$ ," Symp. Combust. 13th (Combustion Institute, Pittsburgh, 1971) 81-88
- Albers, E. A., Hoyermann, K., Schacke, H., Schmatjko, K. J., Wagner, H. Gg., and Wolfrum, J., "Absolute Rate Coefficients for the Reaction of H=Atoms with  $N_2O$  and Some Reactions of CN Radicals," Symp. Combust. 15th (Combustion Institute, Pittsburgh, 1975) 765-773
- Allen, E. R., and Moortgat, G. K., "Kinetic Studies of the Termolecular Reactions of Hydrogen Atoms with Nitric Oxide and with Molecular Oxygen," Third Int. Symp. Gas Kinetics, Brussels (1973), Abstract D1
- Ambidge, P. F., Bradley, J. N., and Whytock, D. A., "Kinetic Study of the Reaction of Hydrogen Atoms with Molecular Chlorine," J. Chem. Soc. Faraday Trans. I 72 1157-1164 (1976)
- Ambidge, P. F., Bradley, J. N., and Whytock, D. A., "Kinetic Study of the Reaction of Hydrogen Atoms with Hydrogen Chloride," J. Chem. Soc. Faraday Trans. I 72, 2143-2149 (1976a)
- Anastasi, C., and Smith, I. W. M., "Rate Measurements of Reactions of OH by Resonance Absorption Part 5. Rate Constants for  $OH + NO_2 (+ M) \rightarrow HNO_3 (+ M)$  Over a Wide Range of Temperature and Pressure," J. Chem. Soc., Faraday Trans. II 72, 1459-1468 (1976)
- Anderson, J. G., and Kaufman, F., "Kinetics of the Reaction  $OH(v = 0) + O_3 \rightarrow HO_2 + O_2$ ," Chem. Phys. Lett. 15, 483-486 (1973)
- Anderson, J. G., Margitan, J. J., and Kaufman, F., "Gas Phase Recombination of OH with NO and  $NO_2$ ," J. Chem. Phys. 60, 3310-3317 (1974)
- Arin, L. M., and Warneck, P., "Reaction of Ozone with Carbon Monoxide," J. Phys. Chem. 76, 1514-1515 (1972)
- Arnoldi, D., and Wolfrum, J., "The Reaction of Vibrationally Excited HCl with Oxygen and Hydrogen Atoms," Chem. Phys. Lett. 24, 234-238 (1974)
- Atkinson, R., and Cvetanović, R. J., "Determination of the Rates of Hydrogen Atom Reaction with NO by a Modulation Technique," Can. J. Chem. 51, 370-372 (1973)
- Atkinson, R., Hansen, D. A., and Pitts, J. N., Jr., "Rate Constants for the Reaction of the OH Radical with  $H_2$  and NO ( $M = Ar$  and  $N_2$ )," J. Chem. Phys. 62, 3284-3288 (1975a)
- Atkinson, R., Hansen, D. A., and Pitts, J. N., Jr., "Rate Constants for the Reaction of OH Radicals with  $CHF_2Cl$ ,  $CF_2Cl_2$ ,  $CFCl_3$ , and  $H_2$  over the Temperature Range 297-434°K," J. Chem. Phys. 63, 1703-1706 (1975)
- Atkinson, R., Perry, R. A., and Pitts, J. N., Jr., "Kinetics of the Reactions of OH Radicals with CO and  $N_2O$ ," Chem. Phys. Lett. 44, 204-208 (1976)
- Atkinson, R., Perry, R. A., and Pitts, J. N., Jr., "Rate Constants for the Reactions of the OH Radical with  $NO_2$  ( $M = Ar$  and  $N_2$ ) and  $SO_2$  ( $M = Ar$ )," J. Chem. Phys. 65, 306-310 (1976)
- Atkinson, R., Perry, R. A., and Pitts, J. N., Jr., "Rate Constants for the Reaction of OH Radicals with Ethylene Over the Temperature Range 299-425°K," J. Chem. Phys. 66, 1197-1201 (1977)



- Atkinson, R., and Pitts, J. N., Jr., "Temperature Dependence of the Reaction Rate Constants for  $\Theta(^3P)$  Atoms with  $C_2H_4$ ,  $C_3H_6$  and  $NO(M = N_2\Theta)$ , Determined by a Modulation Technique," Chem. Phys. Lett. 27, 467-470 (1974)
- Atkinson, R., and Pitts, J. N., Jr., "Rate Constants for the Reaction of  $\Theta(^3P)$  Atoms with  $SO_2$  ( $M = N_2\Theta$ ) Over the Temperature Range 299-392 K," Chem. Phys. Lett. 29, 28-30 (1974)
- Atkinson, R., and Pitts, J. N., Jr., "Temperature Dependence of the Absolute Rate Constants for the Reaction of  $\Theta(^3P)$  Atoms with a Series of Aromatic Hydrocarbons over the Range 299-392°K," J. Phys. Chem. 79, 295-297 (1975a)
- Atkinson, R., and Pitts, J. N., Jr., "Rate Constants for the Reaction of  $\Theta H$  Radicals with Propylene and the Butenes over the Temperature Range 297-425°K," J. Chem. Phys. 63, 3591-3595 (1975b)
- Balakhnin, V. P., Egorov, V. I., and Intezarova, E. I., "Kinetics Investigation of the Elemental Reactions of Oxygen Atoms in the Gas Phase by EPR. II. The Reaction  $\Theta + HCl = \Theta H + Cl$ ," Kinet. Catal. 12, 258-262 (1971); tr. of: Kinet., Katal. 12, 299-303 (1971)
- Baldwin, R. R., Gethin, A., and Walker, R. W., "Reaction of Hydrogen Atoms with Nitrous Oxide," J. Chem. Soc., Faraday Trans. I 69, 352-358 (1973)
- Baldwin, R. R., Fuller, A. R., Longthorn, D., and Walker, R. W., "Addition of Formaldehyde to Slowly Reacting Hydrogen + Oxygen Mixtures," J. Chem. Soc., Faraday Trans. I 68, 1362-1367 (1972)
- Baldwin, R. R., Walker, R. W., and Webster, S. J., "The Carbon Monoxide-Sensitized Decomposition of Hydrogen Peroxide," Combust. Flame 15, 167-172 (1970)
- Ball, M. J., and Larkin, G. S., "Determination of the Rates of Atomic Reactions by the Discharge-Flow Method," Nature Phys. Sci. 245, 63-64 (1973)
- Bar-Ziv, E., Moy, J., and Gordon, R. J., "Temperature Dependence of the Laser Enhanced Reaction  $NO + \Theta_3(OO1)$  II. Contribution from Reactive and Non-Reactive Channels," J. Chem. Phys. 68, 1013 (1978)
- Barassin, J., and Combourieu, J., "No. 1 - Etude Cinetique des Reactions entre l'Oxygene Atomique et les Derives Chlores du Methane. II. - Reactions  $CH_3Cl + \Theta$ ,  $CHCl_3 + \Theta$ ,  $CCl_4 + \Theta$  et  $CH_4 + \Theta$ : Resultats Experimentaux," Bull. Soc. Chim. Fr. 1-5 (1974)
- Barker, J. R., Benson, S. W., and Golden, D. M., "The Decomposition of Dimethyl Peroxide and the Rate Constant for  $CH_3\Theta + \Theta_2 \rightarrow CH_2\Theta + H\Theta_2$ ," Int. J. Chem. Kinet. 9, 31-53 (1977)
- Basco, N., James, D. G. L., and James, F. C., "A Quantitative Study of Alkyl Radical Reactions by Kinetic Spectroscopy. II. Combination of the Methyl Radical with the Oxygen Molecule," Int. J. Chem. Kinet. 4, 129-149 (1972)
- Basco, N., James, D. G. L., and Suart, R. D., "A Quantitative Study of Alkyl Radical Reactions by Kinetic Spectroscopy. Part I. Mutual Combination of Methyl Radicals and Combination of Methyl Radicals with Nitric Oxide," Int. J. Chem. Kinet. 2, 215-234 (1970)
- Bass, A. M., and Laufer, A. H., "The Methyl Radical Combination Rate Constant as Determined by Kinetic Spectroscopy," Int. J. Chem. Kinet. 5, 1053-1065 (1973)
- Bass, A. M., Ledford, A. E., Jr., and Laufer, A. H., "Extinction Coefficients of  $NO_2$  and  $N_2\Theta_4$ ," J. Res. Nat. Bur. Stand., Sect. A 80, 143-166 (1976)



- Batt, L., and McCulloch, R. D., "Pyrolysis of Dimethyl Peroxide," *Int. J. Chem. Kinet.* **8**, 491-500 (1976)
- Batt, L., McCulloch, R. D., and Milne, R. T., "Thermochemical and Kinetic Studies of Alkyl Nitrites ( $RON\dot{O}$ ) =  $D(R\dot{O}-N\dot{O})$ , The Reactions between  $R\dot{O}$  and  $N\dot{O}$ , and the Decomposition of  $R\dot{O}$ ," *Int. J. Chem. Kinet.*, Symp. No. 1, 441-461 (1975)
- Baulch, D. L., and Drysdale, D. D., "An Evaluation of the Rate Data for the Reaction  $C\dot{O} + \dot{O}H \rightarrow C\dot{O}_2 + H$ ," *Combust. Flame* **23**, 215-225 (1974)
- Baulch, D. L., Drysdale, D. D., Horne, D. G., and Lloyd, A. C., "Evaluated Kinetic Data for High Temperature Reactions, Vol. 1: Homogeneous Gas Phase Reactions of the  $H_2-\dot{O}_2$  System," (Butterworths, London, 1972)
- Baulch, D. L., Drysdale, D. D., and Horne, D. G., "Evaluated Kinetic Data for High Temperature Reactions, Vol. 2: Homogeneous Gas Phase Reactions of the  $H_2-N_2-\dot{O}_2$  System," (Butterworths, London, 1973)
- Baulch, D. L., Drysdale, D. D., Duxbury, J., and Grant, S. J., "Evaluated Kinetic Data for High Temperature Reactions, Vol. 3: Homogeneous Gas Phase Reactions of the  $\dot{O}_2-\dot{O}_3$  System, the  $C\dot{O}-\dot{O}_2-H_2$  System, and of Sulphur-Containing Species," (Butterworths, London, 1976)
- Baulch, D. L., Drysdale, D. D., and Horne, D. G., [unpublished data sheets prepared for this survey] University of Leeds, 1973
- Baulch, D. L., Drysdale, D. D., and Horne, D. G., "Rate Constants for Reactions of the  $H_2-N_2-\dot{O}_2$  System," in *Chemical Kinetics Data Survey V. Sixty-six Contributed Rate and Photochemical Data Evaluations on Ninety-four Reactions*, NBSIR 73-206, D. Garvin, Ed., National Bureau of Standards, Washington, D. C. (1973) pages 49-115
- Bayes, K. D., letter to RFH, March 11, 1974
- Becker, K. H., Groth, W., and Kley, D., "The Rate Constant of the Aeronomic Reaction  $N + \dot{O}_2$ ," *Z. Naturforsch. A* **24**, 1280-1281 (1969)
- Becker, K. H., Groth, W., and Schurath, U., "The Quenching of Metastable  $\dot{O}_2(^1\Delta_g)$  and  $\dot{O}_2(^1\Sigma_g^+)$  Molecules," *Chem. Phys. Lett.* **8**, 259-262 (1971)
- Becker, K. H., Groth, W., and Schurath, U., "Reactions of  $\dot{O}_2(^1\Delta_g)$  with  $\dot{O}$ zone," *Chem. Phys. Letts.* **14**, 489-492 (1972)
- Becker, K. H., Groth, W., and Thran, D., "Mechanism of the Air Afterglow  $N\dot{O} + \dot{O} \rightarrow N\dot{O}_2 + h\nu$ ," *Symp. Combust.* 14th (Combustion Institute, Pittsburgh, 1973) 353-363
- Becker, K. H., Inocencio, M., and Schurath, U., "The Reaction of  $\dot{O}$ zone with Hydrogen Sulfide and its Organic Derivatives," *Int. J. Chem. Kinet.*, Symp. No. 1, 205-220 (1975)
- Becker, K. H., Schurath, U., and Seitz, H., " $\dot{O}$ zone- $\dot{O}$ lefin Reactions in the Gas Phase. 1. Rate Constants and Activation Energies," *Int. J. Chem. Kinet.* **6**, 725-739 (1974)
- Beddard, G. S., Giachardi, D. J., and Wayne, R. P., "The Vacuum Ultra-Violet Absorption Spectrum of Anhydrous Nitric Acid," *J. Photochem.* **3**, 321-323 (1974/75)
- Bedford, G., and Thomas, J. H., "Reaction between Ammonia and Nitrogen Dioxide," *J. Chem. Soc., Faraday Trans. I* **68**, 2163-2170 (1972)
- Bemand, P. P., and Clyne, M. A. A., "Atomic Resonance Fluorescence for Rate Constants of Rapid Bimolecular Reactions. Part 6.-Hydrogen Atom Reactions:  $H + Cl_2$  from 300 to 730K and  $H + N\dot{O}_2$  at 298K," *J. Chem. Soc., Faraday Trans. II* **73**, 394-405 (1977)
- Bemand, P. P., Clyne, M. A. A., and Watson, R. T., "Reactions of Chlorine Oxide Radicals. Part 4.-Rate Constants for the Reaction  $Cl + \dot{O}Cl\dot{O}$ ,

- $\Theta + \text{HClO}$ ,  $\text{H} + \text{HClO}$ ,  $\text{NO} + \text{HClO}$  and  $\Theta + \text{ClO}$ ," J. Chem. Soc., Faraday Trans. I 69, 1356-1374 (1973)
- Bemand, P. P., Clyne, M. A. A., and Watson, R. T., "Atomic Resonance Fluorescence and Mass Spectrometry for Measurements of the Rate Constants for Elementary Reactions:  $\Theta(^3\text{P}_j) + \text{NO}_2 \rightarrow \text{NO} + \Theta_2$  and  $\text{NO} + \Theta_3 \rightarrow \text{NO}_2 + \Theta_2$ ," J. Chem. Soc., Faraday Trans. II 70, 564-576 (1974)
- Benson, S. W., and O'Neal, H. E., "Kinetic Data on Gas Phase Unimolecular Reactions," NBS-NSRDS-21 (1970). (Supt. Doc., U.S. Govt. Printing Office, Washington, D.C. 20402)
- Bethke, G. W., "Oscillator Strengths in the Far Ultraviolet. I. Nitric Oxide," J. Chem. Phys. 31, 662-668 (1959)
- Bevan, P. L., and Johnson, G. R. A., "Kinetics of Ozone Formation in the Pulse Radiolysis of Oxygen Gas," J. Chem. Soc., Faraday Trans. I 69, 216-227 (1973)
- Biaume, F., "Nitric Acid Vapour Absorption Cross-Section Spectrum and Its Photodissociation in the Stratosphere," J. Photochem. 2, 139-149 (1973)
- Bida, G. T., Breckenridge, W. H., and Kolln, W. S., "A Kinetic Study of the Very Fast Reaction:  $\Theta(^3\text{P}) + \text{CS} \rightarrow \text{CO}^+ + \text{S}(^3\text{P})$ ," J. Chem. Phys. 64, 3296-3302 (1976)
- Biermann, H. W., Zetzsch, C., and Stuhl, F., "Rate Constant for the Reaction of OH with  $\text{N}_2\text{O}$  at 298 K," Ber. Bunsenges. Phys. Chem. 80, 909-911 (1976)
- Biermann, H. W., Zetzsch, C., and Stuhl, F., "On the Pressure Dependence of the Reaction of HO with CO," paper presented at 13th Informal Conference on Photochemistry, Clearwater Beach, Florida, Jan. 1978
- Bingham, F. W., Johnson, A. W., and Rice, J. K., "Rate Coefficient for Deactivation of  $\Theta(^1\text{S})$  by  $\text{H}_2\text{O}$ ," J. Chem. Phys. 65, 1663-1665 (1976)
- Biordi, J. C., Lazzara, C. P., and Papp, J. F., "Flame Structure Studies of  $\text{CF}_3\text{Br}$ -Inhibited Methane Flames. II. Kinetics and Mechanisms," Symp. Combust. 15th (Combustion Institute, Pittsburgh, 1975) 917-932
- Birks, J. W., Shoemaker, B., Leck, T. J., and Hinton, D. M., "Studies of Reactions of Importance in the Stratosphere I. Reaction of Nitric Oxide with Ozone," J. Chem. Phys. 65, 5181-5185 (1976)
- Birks, J. W., Shoemaker, B., Leck, T. J., Borders, R. A., and Hart, L. J., "Studies of Reactions of Importance in the Stratosphere. II. Reactions Involving Chlorine Nitrate and Chlorine Dioxide," J. Chem. Phys. 66, 4591-4599 (1977)
- Bishop, W. P., and Dorfman, L. M., "Pulse Radiolysis Studies. XVI. Kinetics of the Reaction of Gaseous Hydrogen Atoms with Molecular Oxygen by Fast Lyman- $\alpha$  Absorption Spectrophotometry," J. Chem. Phys. 52, 3210-3216 (1970)
- Blume, C. W., Hisatsune, I. C., and Heicklen, J., "Gas-Phase Ozonolysis of Cis- and Trans-Dichloroethylene," Int. J. Chem. Kinet. 8, 235-258 (1976)
- Bowman, C. T., "Non-Equilibrium Radical Concentrations in Shock-Initiated Methane Oxidation," Symp. Combust. 15th (Combustion Institute, Pittsburgh, 1975) 869-882
- Bradley, J. N., Hack, W., Hoyeremann, K., and Wagner, H. Gg., "Kinetics of the Reaction of Hydroxyl Radicals with Ethylene and with  $\text{C}_3$  Hydrocarbons," J. Chem. Soc., Faraday Trans. I 69, 1889-1898 (1973)
- Braslavsky, S., and Heicklen, J., "The Gas-Phase Reaction of  $\Theta_3$  with  $\text{H}_2\text{CO}$ ," Int. J. Chem. Kinet. 8, 801-808 (1976)
- Breckenridge, W. H., and Miller, T. A., "Kinetic Study by EPR of the Production and Decay of  $\text{SO}(^1\Delta)$  in the Reaction of  $\Theta_2(^1\Delta_g)$  with  $\text{SO}(^3\Sigma^-)$ ," J. Chem. Phys. 56, 465-474 (1972)

- Brown, R. D. H., Glass, G. P., and Smith, I. W. M., "The Relaxation of HCl(v=1) and DCl (v=1) by  $\Theta$  Atoms Between 196 and 400 K," Chem. Phys. Lett. 32, 517-520 (1975)
- Brown, R. D. H., and Smith, I. W. M., "Absolute Rate Constants for the Reactions  $\Theta(^3P)$  Atoms with HCl and HBr," Int. J. Chem. Kinet. 7, 301-315 (1975)
- Burrows, J. P., Harris, G. W., and Thrush, B. A., "Rates of Reaction of  $H\Theta_2$  with H $\Theta$  and  $\Theta$  studied by Laser Magnetic Resonance," Nature 267, 233-234 (1977)
- Butler, R., Solomon, I. J., and Snelson, A., "Pressure Dependence of the  $C\Theta + \Theta H$  Rate Constant in  $\Theta_2 + N_2$  Mixtures," Chem. Phys. Lett., in press (1977)
- Calvert, J. G., Kerr, J. A., Demerjian, K. L., and McQuigg, R. D., "Photolysis of Formaldehyde as a Hydrogen Atom Source in the Lower Atmosphere," Science 175, 751-752 (1972)
- Campbell, I. M., and Goodman, K., "Reaction of  $\Theta(^3P)$  Atoms with Nitromethane Vapour at 295 K," Chem. Phys. Lett. 34, 105-108 (1975a)
- Campbell, I. M., and Goodman, K., "Rate Constants for Reactions of Hydroxyl Radicals with Nitromethane and Methyl Nitrite Vapours at 292 K," Chem. Phys. Lett. 36, 382-384 (1975b)
- Campbell, I. M., and Gray, C. N., "Rate Constants for  $\Theta(^3P)$  Recombination and Association with  $N(^4S)$ ," Chem. Phys. Lett. 18, 607-609 (1973)
- Campbell, I. M., and Handy, B. J., "Studies of Reactions of Atoms in a Discharge Flow Stirred Reactor Part 1 - The  $\Theta + H_2 + N\Theta$  System," J. Chem. Soc., Faraday Trans. I 71, 2097-2106 (1975)
- Campbell, I. M., McLaughlin, D. F., and Handy, B. J., "Rate Constants for Reactions of Hydroxyl Radicals with Alcohol Vapours at 292 K," Chem. Phys. Lett. 38, 362-364 (1976)
- Castellano, E., and Schumacher, H. J., "La Descomposicion Fotoquimica del Ozono a 3200 A," An. Asoc. Quim. Argent. 63, 9-15 (1975)
- Castleman, A. W., Jr., Davis, R. E., Munkelwitz, H. R., Tang, I. N., and Wood, W. P., "Kinetics of Association Reactions Pertaining to  $H_2S\Theta_4$  Aerosol Formation," Int. J. Chem. Kinet., Symp. No. 1, 629-640 (1975)
- Castleman, A. W., Jr., and Tang, I. N., "Kinetics of the Association Reaction of  $S\Theta_2$  with the Hydroxyl Radical," J. Photochem. 6, 349-354 (1977)
- Chan, W. H., Nordstrom, R. J., Calvert, J. G., and Shaw, J. H., "Kinetic Study of  $H\Theta N\Theta$  Formation and Decay Reactions in Gaseous Mixtures of  $H\Theta N\Theta$ ,  $N\Theta$ ,  $N\Theta_2$ ,  $H_2\Theta$ , and  $N_2$ ," Environ. Sci. Technol. 10, 674-682 (1976); see also article by same authors (which is next entry), Chem. Phys. Lett. 37, 441-446 (1976)
- Chan, W. H., Nordstrom, R. J., Calvert, J. G., and Shaw, J. H., "An IRFTS Spectroscopic Study of the Kinetics and the Mechanism of the Reactions in the Gaseous System,  $H\Theta N\Theta$ ,  $N\Theta$ ,  $N\Theta_2$ ,  $H_2\Theta$ ," Chem. Phys. Lett. 37, 441-446 (1976)
- Chan, W. H., Uselman, W. M., Calvert, J. G., and Shaw, J. H., "The Pressure Dependence of the Rate Constant for the Reaction:  $H\Theta + C\Theta \rightarrow H + C\Theta_2$ ," Chem. Phys. Lett. 45, 240-244 (1977)
- Chang, J. S., and Kaufman, F., "Upper Limits of the Rate Constants for the Reactions of  $CFCl_3$ (F-11),  $CF_2Cl_2$ (F-12), and  $N_2\Theta$  with  $\Theta H$  and Estimates of Corresponding Lower Limit of their Tropospheric Lifetimes," Geophys. Res. Lett. 4, 192-194 (1977)



- Chang, J. S., and Kaufman, F., "Kinetics of the Reactions of Hydroxyl Radicals with Some Halocarbons:  $\text{CHFCl}_2$ ,  $\text{CHF}_2\text{Cl}$ ,  $\text{CH}_3\text{CCl}_3$ ,  $\text{C}_2\text{HCl}_3$ , and  $\text{C}_2\text{Cl}_4$ ," J. Chem. Phys. **66**, 4989-4994 (1977)
- Chapman, C. J., and Wayne, R. P., "The Reaction of Atomic Oxygen and Hydrogen with Nitric Acid," Int. J. Chem. Kinet. **6**, 617-630 (1974)
- Chung, K., Calvert, J. G., and Bottenheim, J. W., "The Photochemistry of Sulfur Dioxide Excited within its First Allowed Band (3130 o) and the 'Forbidden' Band (3700-4000 o)," Int. J. Chem. Kinet. **7**, 161-182 (1975)
- Clark, I. D., Jones, I. T. N., and Wayne, R. P., "The Kinetics of the Reaction between  $\text{O}_2(^1\Delta_g)$  and Ozone," Proc. Roy. Soc. London A **317**, 407-416 (1970)
- Clark, J. H., Moore, C. B., and Nogar, N. S., "The Photochemistry of Formaldehyde: Absolute Quantum Yields, Radical Reactions and NO Reactions," typescript, Dept. of Chemistry, Univ. of California, Berkeley, CA (1977)
- Clark, I. D., and Wayne, R. P., "Kinetics of the Reaction between Atomic Nitrogen and Molecular Oxygen in the Ground ( $^3\Sigma_g^-$ ) and First Excited ( $^1\Delta_g$ ) States," Proc. Roy. Soc. London A **316**, 539-550 (1970)
- Clough, P. N., and Thrush, B. A., "Mechanism of Chemiluminescent Reaction between Nitric Oxide and Ozone," Trans. Faraday Soc. **63**, 915-925 (1967)
- Clyne, M. A. A., and Coxon, J. A., "Kinetics Studies of Oxyhalogen Radical Systems," Proc. Roy. Soc., (London) A **303**, 207-231 (1968)
- Clyne, M. A. A., and Cruse, H. W., "Studies of Ground-State  $^2\text{P}_{3/2}$  Halogen Atoms Using Atomic Resonance Absorption," Trans. Faraday Soc. **67**, 2869-2885 (1971)
- Clyne, M. A. A., and Cruse, H. W., "Atomic Resonance Fluorescence Spectrometry for Rate Constants of Rapid Bimolecular Reactions. Part 1. Reactions  $\text{O} + \text{NO}_2$ ,  $\text{Cl} + \text{ClNO}$ ,  $\text{Br} + \text{ClNO}$ ," J. Chem. Soc., Faraday Trans. II **68**, 1281-1299 (1972)
- Clyne, M. A. A., and Cruse, H. W., "Rates of Elementary Reaction Involving the  $\text{BrO}(\text{X}^2\Pi)$  and  $\text{IO}(\text{X}^2\Pi)$  Radicals. Part 1.-Formation and Decay of the  $\text{BrO}$  Radical," Trans. Faraday Soc. **66**, 2214-2226 (1970)
- Clyne, M. A. A., and Down, S., "Kinetic Behaviour of OH  $\text{X}^2\Pi$  and  $\text{A}^2\Sigma^+$  using Molecular Resonance Fluorescence Spectrometry," J. Chem. Soc. Faraday Trans. II **70**, 253-266 (1974)
- Clyne, M. A. A., and McDermid, I. S., "Mass Spectrometric Determinations of Rates of Elementary Reactions of NO and of  $\text{NO}_2$  with Ground State  $\text{N}^4\text{S}$  Atoms," J. Chem. Soc., Faraday Trans. I **71**, 2189-2208 (1975)
- Clyne, M. A. A., and Monkhouse, P. B., "Atomic Resonance Fluorescence for Rate Constants of Rapid Bimolecular Reactions. Part 5-Hydrogen Atom Reactions;  $\text{H} + \text{NO}_2$  and  $\text{H} + \text{O}_3$ ," J. Chem. Soc., Faraday Trans. II **73**, 298-309 (1977)
- Clyne, M. A. A., Monkhouse, P. B., and Townsend, L. W., "Reactions of  $\text{O}(^3\text{P}_j)$  Atoms with Halogens: The Rate Constants for the Elementary Reactions  $\text{O} + \text{BrCl}$ ,  $\text{O} + \text{Br}_2$ , and  $\text{O} + \text{Cl}_2$ ," Int. J. Chem. Kinet. **8**, 425-449 (1976)
- Clyne, M. A. A., and Nip, W. S., "Reactions of Chlorine Oxide Radicals. Part 6.-The Reaction  $\text{O} + \text{ClO} \rightarrow \text{Cl} + \text{O}_2$  from 220 to 425 K," J. Chem. Soc., Faraday Trans. I **72**, 2211-2217 (1976)
- Clyne, M. A. A., and Nip, W. S., "Study of Elementary Reactions by Atomic Resonance Absorption with a Non-Reversed Source Part 1 - The Reaction  $\text{Cl} + \text{O}_3 \rightarrow \text{ClO} + \text{O}_2$ ," J. Chem. Soc., Faraday Trans. II **72**, 838-847 (1976)

- Clyne, M. A. A., and Walker, R. F., "Absolute Rate Constants for Elementary Reactions in the Chlorination of  $\text{CH}_4$ ,  $\text{CD}_4$ ,  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_2\text{Cl}_2$ ,  $\text{CHCl}_3$ ,  $\text{CDCl}_3$  and  $\text{CBrCl}_3$ ," J. Chem. Soc., Faraday Trans. I **69**, 1547-1567 (1973)
- Clyne, M. A. A., and Watson, R. T., "Kinetics Studies for Diatomic Free Radicals using Mass Spectrometry Part 3 - Elementary Reactions Involving  $\text{BrO X}^2\Pi$  Radicals," J. Chem. Soc., Faraday Trans. I **71**, 336-350 (1975)
- Collins, R. J., Husain, D., and Donovan, R. J., "Kinetic and Spectroscopic Studies of  $\text{O}_2(a^1\Delta_g)$  by Time-Resolved Absorption Spectroscopy in the Vacuum Ultra-violet," J. Chem. Soc., Faraday Trans. II **69**, 145-157 (1973)
- Coltharp, R. N., Worley, S. D., and Potter, A. E., "Reaction Rate of Vibrationally Excited Hydroxyl with Ozone," Appl. Optics **10**, 1786-1789 (1971)
- Colussi, A. J., Singleton, D. L., Irwin, R. S., and Cvetanović, R. J., "Absolute Rates of Oxygen ( $^3P$ ) Atom Reactions with Benzene and Toluene," J. Phys. Chem. **79**, 1900-1903 (1975)
- Cox, R. A., "The Photolysis of Nitrous Acid in the Presence of Carbon Monoxide and Sulphur Dioxide," J. Photochem. **3**, 291-304 (1974)
- Cox, R. A., "The Photolysis of Gaseous Nitrous Acid - A Technique for Obtaining Kinetic Data on Atmospheric Photooxidation Reactions," Int. J. Chem. Kinet., Symp. No. 1, 379-398 (1975)
- Cox, R. A., and Derwent, R. G., "Kinetics of the Reaction of  $\text{HO}_2$  with Nitric Oxide and Nitrogen Dioxide," J. Photochem. **4**, 139-153 (1975)
- Cox, R. A., and Derwent, R. G., "The Ultraviolet Absorption Spectrum of Gaseous Nitrous Acid," J. Photochem. **6**, 23-34 (1976)
- Cox, R. A., Derwent, R. G., and Holt, P. M., "The Photo-oxidation of Ammonia in the Presence of  $\text{NO}$  and  $\text{NO}_2$ ," Chemosphere, No. 4, 201-205 (1975)
- Cox, R. A., Derwent, R. G., and Holt, P. M., "Relative Rate Constants for the Reactions of OH Radicals with  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_6$ ,  $\text{NO}$  and  $\text{HONO}$  at Atmospheric Pressure and 296 K," J. Chem. Soc., Faraday Trans. I **72**, 2031-2043 (1976)
- Cox, R. A., Derwent, R. G., Holt, P. M., and Kerr, J. A., "Photo-oxidation of Methane in the Presence of  $\text{NO}$  and  $\text{NO}_2$ ," J. Chem. Soc. Faraday Trans. I **72**, 2044-2060 (1976)
- Cupitt, L. T., and Glass, G. P., "Reactions of SH with Atomic Oxygen and Hydrogen," Int. J. Chem. Kinet., Symp. No. 1, 39-50 (1975)
- Daubendiek, R. L., and Calvert, J. G., "A Study of the  $\text{N}_2\text{O}_5\text{-SO}_2\text{-O}_3$  Reaction System," Environ. Lett. **8**, 103-116 (1975)
- Davidson, J. A., Schiff, H. I., Streit, G. E., McAfee, J. R., Schmeltekopf, A. I., and Howard, C. J., "Temperature Dependence of  $\text{O}(^1D)$  Rate Constants for Reactions with  $\text{N}_2\text{O}$ ,  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{HCl}$ , and  $\text{NH}_3$ ," J. Chem. Phys. **67**, 5021-5025 (1977)
- Davidson, J. A., and Thrush, B. A., "Reaction of Oxygen Atoms with Methyl and Ethyl Nitrites," J. Chem. Soc., Faraday Trans. I **71**, 2413-2420 (1975)
- Davis, D. D., "Absolute Rate Constants for Elementary Reactions of Atmospheric Importance: Results from the University of Maryland Gas Kinetics Laboratory," Report 3 (Univ. of Maryland, College Park, Md. 20742, 1976)
- Davis, D. D., Braun, W., and Bass, A. M., "Reactions of  $\text{Cl}^2\text{P}_{3/2}$ : Absolute Rate Constants for Reaction with  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_6$ ,  $\text{CH}_2\text{Cl}_2$ ,  $\text{C}_2\text{Cl}_4$ , and  $\text{c-C}_6\text{H}_{12}$ ," Int. J. Chem. Kinet. **2**, 101-114 (1970)
- Davis, D. D., Bollinger, W., and Fischer, S., "A Kinetics Study of the Reaction of the OH Free Radical with Aromatic Compounds. I. Absolute



- Rate Constants for Reaction with Benzene and Toluene at 300°K," J. Phys. Chem. 79, 293-294 (1975)
- Davis, D. D., Fischer, S., and Schiff, R., "Flash Photolysis-Resonance Fluorescence Kinetics Study: Temperature Dependence of the Reactions  $\text{OH} + \text{C}_6\text{H}_6 \rightarrow \text{C}_6\text{H}_5 + \text{H}$  and  $\text{OH} + \text{CH}_4 \rightarrow \text{H}_2\text{O} + \text{CH}_3$ ," J. Chem. Phys. 61, 2213-2219 (1974)
- Davis, D. D., Fischer, S., Schiff, R., Watson, R. T., and Bollinger, W., "A Kinetics Study of the Reaction of OH Radicals with Two  $\text{C}_2$  Hydrocarbons:  $\text{C}_2\text{H}_4$  and  $\text{C}_2\text{H}_2$ ," J. Chem. Phys. 63, 1707-1712 (1975)
- Davis, D. D., Herron, J. T., and Hule, R. E., "Absolute Rate Constants for the Reaction  $\text{O}(^3\text{P}) + \text{N}_2\text{O} \rightarrow \text{NO} + \text{O}_2$  Over the Temperature Range 230-339K," J. Chem. Phys. 58, 530-535 (1973)
- Davis, D. D., Machado, G., Conaway, B., Oh, Y., and Watson, R., "A Temperature Dependent Kinetics Study of the Reaction of OH with  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_2\text{Cl}_2$ ,  $\text{CHCl}_3$ , and  $\text{CH}_3\text{Br}$ ," J. Chem. Phys. 65, 1268-1274 (1976)
- Davis, D. D., Payne, W., and Stief, L., "The Hydroperoxyl Radical in Atmospheric Chemical Dynamics: Reaction with Carbon Monoxide," Science 179, 280-282 (1973)
- Davis, D. D., Prusaczyk, J., Dwyer, M., and Kim, P., "A Stop-Flow Time of Flight Mass Spectrometry Kinetics Study. Reaction of Ozone with Nitrogen Dioxide and Sulfur Dioxide," J. Phys. Chem. 78, 1775-1779 (1974)
- Davis, D. D., Wong, W., and Lephardt, J., "A Laser Flash Photolysis-Resonance Fluorescence Kinetic Study: Reaction of  $\text{O}(^3\text{P})$  with  $\text{O}_3$ ," Chem. Phys. Lett. 22, 273-278 (1973)
- Davis, D. D., Wong, W., and Schiff, R., "A Dye Laser Flash Photolysis Kinetics Study of the Reaction of Ground-State Atomic Oxygen with Hydrogen Peroxide," J. Phys. Chem. 78, 463-464 (1974)
- Demerjian, K. L., Kerr, J. A., and Calvert, J. G., "The Mechanism of Photochemical Smog Formation," Adv. Environ. Sci. Technol. 4, 1-262 (1974)
- J. N. Pitts, Jr., and R. L. Metcalf, editors, Wiley-Interscience
- DeMore, W. B., "Rate Constants for the Reactions of Hydroxyl and Hydroperoxyl with Ozone," Science 180, 735-737 (1973)
- DeMore, W. B., "Arrhenius Constants for the Reactions of Ozone with Ethylene and Acetylene," Int. J. Chem. Kinet. 1, 209-220 (1969)
- DeMore, W. B., "Rate Constant Ratio for the Reactions of OH with  $\text{O}_3$  and  $\text{C}_6\text{H}_6$ ," Int. J. Chem. Kinet., Symp. No. 1, 273-279 (1975)
- DeMore, W. B., and Tschuikow-Roux, E., "Temperature Dependence of the Reactions of OH and  $\text{HO}_2$  with  $\text{O}_3$ ," J. Phys. Chem. 78, 1447-1451 (1974)
- Dillemath, F. J., Skidmore, D. R., and Schubert, C. C., "The Reaction of Ozone with Methane," J. Phys. Chem. 64, 1496-1499 (1960)
- Donovan, R. J., Husain, D., and Kirsch, L. J., "Reactions of Atomic Oxygen. Part 1: The Rate of the Reaction  $\text{O} + \text{O}_2 + \text{M} \rightarrow \text{O}_3 + \text{M}$  ( $\text{M} = \text{He, Ar and Kr}$ )," Trans. Faraday Soc. 66, 2551-2559 (1970)
- Dubinsky, R. N., McKenney, D. J., "Determination of the Rate Constant of the  $\text{O} + \text{H}_2 \rightarrow \text{OH} + \text{H}$  Reaction using Atomic Oxygen Resonance Fluorescence and the Air Afterglow Techniques," Can. J. Chem. 53, 3531-3541 (1975)
- Fettis, G. C., and Knox, J. H., "Rate Constants of Halogen Atom Reactions," G. Porter, Ed., Progress in Reaction Kinet. (Pergamon Press) 2, 1-38 (1964)
- Findlay, F. D., and Snelling, D. R., "Temperature Dependence of the Rate Constant for the Reaction  $\text{O}_2(^1\Delta_g) + \text{O}_3 \rightarrow 2\text{O}_2 + \text{O}$ ," J. Chem. Phys. 54, 2750-2755 (1971)

- Findlay, F. D., and Snelling, D. R., "Collisional Deactivation of  $O_2(^1\Delta_g)$ ," J. Chem. Phys. 55, 545-551 (1971a)
- Fisher, E. R., and McCarty, M., Jr., "Study of the Reaction of Electronically Excited Oxygen Molecules with Carbon Monoxide," J. Chem. Phys. 45, 781-784 (1966)
- Fletcher, I. S., and Husain, D., "The Collisional Quenching of Electronically Excited Oxygen Atoms,  $O(^1D_2)$  by the Gases  $NH_3$ ,  $H_2O_2$ ,  $C_2H_6$ ,  $C_3H_8$ , and  $C(CH_3)_4$  Using Time-Resolved Attenuation of Atomic Resonance Radiation," Can. J. Chem. 54, 1765-1770 (1976a)
- Fletcher, I. S., and Husain, D., "Absolute Reaction Rates of Oxygen( $^1D_2$ ) with Halogenated Paraffins by Atomic Absorption Spectroscopy in the Vacuum Ultraviolet," J. Phys. Chem. 80, 1837-1840 (1976b)
- Foner, S. N., and Hudson, R. L., "Mass Spectrometry of the  $HO_2$  Free Radical," J. Chem. Phys. 36, 2681-2688 (1962)
- Francis, P. D., "The Production of Oxygen Atoms in a Microwave Discharge and the Recombination Kinetics in a Gas Flow System," Brit. J. Appl. Phys. 2, 1717-1730 (1969)
- Gardiner, W. C., Jr., Mallard, W. G., McFarland, M., Morinaga, L., Owen, J. H., Rawlins, W. T., Takeyama, T., and Walker, B. F., "Elementary Reaction Rates from Post-Induction-Period Profiles in Shock-Initiated Combustion," Symp. Combust. 14th (Combustion Institute, Pittsburgh, 1973) 61-72
- Gardiner, W. C., Jr., Mallard, W. G., and Owen, J. H., "Rate Constant of  $OH + H_2 = H_2O + H$  from 1350 to 1600 K," J. Chem. Phys. 60, 2290-2295 (1974)
- Gehring, M., Hoyermann, K., Schacke, H., and Wolfrum, J., "Direct Studies of Some Elementary Steps for the Formation and Destruction of Nitric Oxide in the  $H-N-O$  System," Symp. Combust. 14th (Combustion Institute, Pittsburgh, 1973) 99-105
- Ghormley, J. A., Ellsworth, R. L., and Hochanadel, C. J., "Reaction of Excited Oxygen Atoms with Nitrous Oxide. Rate Constants for Reaction of Ozone with Nitric Oxide and with Nitrogen Dioxide," J. Phys. Chem. 77, 1341-1345 (1973). Erratum: 78, 2698 (1974)
- Giachardi, D. J., Harris, G. W., and Wayne, R. P., "Excited State Formation in the  $H + O_2$  System," Chem. Phys. Lett. 32, 586-588 (1975)
- Giachardi, D. J., Harris, G. W., and Wayne, R. P., "Energy Transfer from Excited  $NO_2^*$  to Molecular Oxygen," J. Chem. Soc., Faraday Trans. II 72, 619-630 (1976)
- Glanzer, K., and Troe, J., [unpublished data sheets prepared for this survey] Ecole Polytechnique Federale de Lausanne (1973)
- Glanzer, K., and Troe, J., " $HO_2$  Formation in Shock Heated  $HN O_3 - NO_2$  Mixtures," Ber. Bunsenges. Phys. Chem. 79, 465-469 (1975)
- Glavas, S., and Toby, S., "The Reaction between Ozone and Hydrogen Sulfide: Kinetics and Effect of Added Gases," Am. Chem. Soc., ACS Symp. Series 17, 122-131 (1975)
- Gordon, R. J., and Lin, M. C., "The Reaction of Nitric Oxide with Vibrationally Excited Ozone," Chem. Phys. Lett. 22, 262-268 (1973)
- Gordon, R. J., and Lin, M. C., "The Reaction of Nitric Oxide with Vibrationally Excited Ozone. II.," J. Chem. Phys. 64, 1058-1064 (1976)
- Gordon, S., and Mulac, W. A., "Reaction of the  $OH(X^2\Pi)$  Radical Produced by the Pulse Radiolysis of Water Vapor," Int. J. Chem. Kinet., Symp. No. 1, 289-299 (1975)

- Gordon, S., Mulac, W., and Nangia, P., "Pulse Radiolysis of Ammonia Gas. II. Rate of Disappearance of the  $\text{NH}_2(\text{X}^2\text{B}_1)$  Radical," J. Phys. Chem. 75, 2087-2093 (1971)
- Gorse, R. A., and Volman, D. H., "Photochemistry of the Gaseous Hydrogen Peroxide-Carbon Monoxide System: Rate Constants for Hydroxyl Radical Reactions with Hydrogen Peroxide and Isobutane by Competitive Kinetics," J. Photochem. 1, 1-10 (1972)
- Gorse, R. A., and Volman, D. H., "Photochemistry of the Gaseous Hydrogen Peroxide-Carbon Monoxide System II: Rate Constants for Hydroxyl Radical Reactions with Hydrocarbons and for Hydrogen Atom Reactions with Hydrogen Peroxide," J. Photochem. 3, 115-122 (1974)
- Graham, R. A., and Johnston, H. S., "Kinetics of the Gas-Phase Reaction between Ozone and Nitrogen Dioxide," J. Chem. Phys. 60, 4628-4629 (1974)
- Graham, R. A., and Johnston, H. S., "The Photochemistry of  $\text{NO}_3$  and the Kinetics of the  $\text{N}_2\text{O}_5$ - $\text{O}_3$  System," J. Phys. Chem. 82, 254-268 (1978)
- Graham, R. E., and Gutman, D., "Measurement of Branching Ratios for the  $\text{O} + \text{CS}_2 \rightarrow \text{OCS} + \text{S}$  Reaction," Photochem. Conference 12th (National Bureau of Standards, Washington, D. C. 20234, 1976) H4-1
- Gray, D., Lissi, E., and Heicklen, J., "The Reaction of Hydrogen Peroxide with Nitrogen Dioxide and Nitric Oxide," J. Phys. Chem. 76, 1919-1924 (1972)
- Greiner, N. R., "Hydroxyl Radical Kinetics by Kinetic Spectroscopy. III. Reactions with  $\text{H}_2\text{O}_2$  in the Range 300-458°K," J. Phys. Chem. 72, 406-410 (1968)
- Greiner, N. R., "Hydroxyl Radical Kinetics by Kinetic Spectroscopy. V. Reactions with  $\text{H}_2$  and  $\text{CO}$  in the Range 300-500°K," J. Chem. Phys. 51, 5049-5051 (1969)
- Greiner, N. R., "Hydroxyl Radical Kinetics by Kinetic Spectroscopy. VI. Reactions with Alkanes in the Range 300-500°K," J. Chem. Phys. 53, 1070-1076 (1970)
- Greiner, N. R., "Hydroxyl Radical Kinetics by Kinetic Spectroscopy. VII. The Reaction with Ethylene in the Range 300-500°K," J. Chem. Phys. 53, 1284-1285 (1970a)
- Hack, W., Hoyer mann, K., and Wagner, H. Gg., "Reaktionen des Hydroxylradikals mit Ammoniak und Hydrazin in der gasphase," Ber. Bunsenges. Physik. Chem. 78, 386-391 (1974)
- Hack, W., Hoyer mann, K., and Wagner, H. Gg., "The Reaction  $\text{NO} + \text{HO}_2 \rightarrow \text{NO}_2 + \text{OH}$  with  $\text{OH} + \text{H}_2\text{O}_2 \rightarrow \text{HO}_2 + \text{H}_2\text{O}$  as an  $\text{HO}_2$ -Source," Int. J. Chem. Kinet., Symp. No. 1, 329-339 (1975)
- Hamilton, E. J., Jr., "Water Vapor Dependence of the Kinetics of the Self-Reaction of  $\text{HO}_2$  in the Gas Phase," J. Chem. Phys. 63, 3682-3683 (1975)
- Hamilton, E. J., Jr., and Lii, R.-R., "The Dependence on  $\text{H}_2\text{O}$  and on  $\text{NH}_3$  of the Kinetics of the Self-Reaction of  $\text{HO}_2$  in the Gas-Phase Formation of  $\text{HO}_2 \cdot \text{H}_2\text{O}$  and  $\text{HO}_2 \cdot \text{NH}_3$  Complexes, Int. J. Chem. Kinet. 9, 875-885 (1977)
- Hampson, R. F., (editor), "Survey of Photochemical and Rate Data for Twenty-Eight Reactions of Interest in Atmospheric Chemistry," J. Phys. Chem. Ref. Data 2, No. 2 pgs. 267-312 (1973). These are revised versions of data sheets originally distributed as NBS Reports 10692 and 10828.
- Hampson, R. F., (editor), "Chemical Kinetics Data Survey VI. Photochemical and Rate Data for Twelve Gas Phase Reactions of Interest for Atmospheric Chemistry," NBSIR 73-207 (1973a)
- Hancock, G., Lange, W., Lenzi, M., and Welge, K. H., "Laser Fluorescence of  $\text{NH}_2$  and Rate Constant Measurement of  $\text{NH}_2 + \text{NO}$ ," Chem Phys. Lett. 33, 168-172 (1975)



- Hansen, D. A., Atkinson, R., and Pitts, J. N., Jr., "Rate Constants for the Reaction of OH Radicals with a Series of Aromatic Hydrocarbons," J. Phys. Chem. 79, 1763-1766 (1975)
- Hansen, I., Hoinghaus, K., Zetzsch, C., and Stuhl, F., "Detection of  $\text{NH}(\text{X}^3\text{E}^-)$  by Resonance Fluorescence in the Pulsed Vacuum UV Photolysis of  $\text{NH}_3$  and Its Application to Reactions of NH Radicals," Chem. Phys. Lett. 42, 370-372 (1976)
- Hanst, P. L., Stephens, E. R., Scott, W. E., and Doerr, R. C., "Atmospheric Ozone-Olefin Reactions," (The Franklin Institute, Philadelphia, Pa. 1958)
- Harker, A., and Johnston, H. S., "Photolysis of Nitrogen Dioxide to Produce Transient O,  $\text{NO}_3$  and  $\text{N}_2\text{O}_5$ ," J. Phys. Chem. 77, 1153-1156 (1973)
- Harris, G. W., and Wayne, R. P., "Reaction of Hydroxyl Radicals with NO,  $\text{NO}_2$  and  $\text{SO}_2$ ," J. Chem. Soc., Faraday Trans. I 71, 610-617 (1975)
- Hastie, D. R., Freeman, C. G., McEwan, M. J., and Schiff, H. I., "The Reactions of Ozone with Methyl and Ethyl Nitrites," Int. J. Chem. Kinet. 8, 307-313 (1976)
- Heicklen, J., "Gas-Phase Reactions of Alkylperoxy and Alkoxy Radicals," Adv. Chem. Ser. (Amer. Chem. Soc., Washington) 76, 23-39 (1968)
- Heicklen, J., "Photochemical and Rate Data for Methyl Nitrite, Methoxy and Methylperoxy," Chemical Kinetics Data Survey V. Sixty-six Contributed Rate and Photochemical Data Evaluations on Ninety-four Reactions, NBSIR 73-206, D. Garvin, Ed., National Bureau of Standards, Washington, D.C. (1973) pages 43-48
- Heidner, R. F., III, and Husain, D., "Electronically Excited Oxygen Atoms,  $\text{O}(2^1\text{D}_2)$ . A Time-Resolved Study of the Collisional Quenching by the Gases  $\text{H}_2$ ,  $\text{D}_2$ , NO,  $\text{N}_2\text{O}$ ,  $\text{NO}_2$ ,  $\text{CH}_4$ , and  $\text{C}_3\text{O}_2$  Using Atomic Absorption Spectroscopy in the Vacuum Ultraviolet," Int. J. Chem. Kinet. 5, 819-831 (1973)
- Heidner, R. F., III, Husain, D., and Wiesenfeld, J. R., "Kinetic Investigation of Electronically Excited Oxygen Atoms,  $\text{O}(2^1\text{D}_2)$ , by Time-resolved Attenuation of Atomic Resonance Radiation in the Vacuum Ultra-violet. Part 2.-Collisional Quenching by the Atmospheric Gases  $\text{N}_2$ ,  $\text{O}_2$ , CO,  $\text{CO}_2$ ,  $\text{H}_2\text{O}$  and  $\text{O}_3$ ," J. Chem. Soc., Faraday Trans. II 69, 927-938 (1973)
- Herron, J. T., and Huie, R. E., "Rate Constants for the Reactions of Ozone with Ethene and Propene, from 235.0 to 362.0 K," J. Phys. Chem. 78, 2085-2088 (1974)
- Herron, J. T., and Huie, R. E., "Rate Constants for the Reactions of Atomic Oxygen ( $\text{O}^3\text{P}$ ) with Organic Compounds in the Gas Phase," J. Phys. Chem. Ref. Data 2, 467-518 (1973)
- Herron, J. T., and Huie, R. E., "Rates of Reaction of Atomic Oxygen. II. Some  $\text{C}_2$  to  $\text{C}_8$  Alkanes," J. Phys. Chem. 73, 3327-3337 (1969)
- Hikida, T., Eyre, J. A., and Dorfman, L. M., "Pulse Radiolysis Studies. XX. Kinetics of Some Addition Reactions of Gaseous Hydrogen Atoms by Fast Lyman- $\alpha$  Absorption Spectrophotometry," J. Chem. Phys. 54, 3422-3428 (1971)
- Hippler, H., and Troe, J., "Hochdruckbereich der Rekombination  $\text{O} + \text{O}_2 \rightarrow \text{O}_3$ ," Ber. Bunsenges. Physik. Chem. 75, 27-32 (1971)
- Hippler, H., Schippert, C., and Troe, J., "Photolysis of  $\text{NO}_2$  and Collisional Energy Transfer in the Reactions  $\text{O} + \text{NO} \rightarrow \text{NO}_2$  and  $\text{O} + \text{NO}_2 \rightarrow \text{NO}_3$ ," Int. J. Chem. Kinet., Symp. No. 1, 27-38 (1975)
- Hochanadel, C. J., Ghormley, J. A., and Ogren, P. J., "Absorption Spectrum and Reaction Kinetics of  $\text{HO}_2$  Radical in the Gas Phase," J. Chem. Phys. 56, 4426-4432 (1972)

- Hochanadel, C. J., Ghormley, J. A., Boyle, J. W., and Ogren, P. J., "Absorption Spectrum and Rates of Formation and Decay of the  $\text{CH}_2\text{O}_2$  Radical," J. Phys. Chem. 81, 3-7 (1977)
- Hogan, L. G., and Burch, D. S., "A Measurement of the Rate Constant for the Reaction  $\text{O} + \text{O}_2 + \text{O}_2 \rightarrow \text{O}_3 + \text{O}_2$ ," J. Chem. Phys. 65, 894-900 (1976)
- Hollinden, G. A., Kurylo, M. J., and Timmons, R. B., "Electron Spin Resonance Study of the Kinetics of the Reaction of  $\text{O}(^3\text{P})$  Atoms with  $\text{H}_2\text{S}$ ," J. Phys. Chem. 74, 588-591 (1970)
- Holmes, H. H., and Daniels, F., "The Photolysis of Nitrogen Oxides:  $\text{N}_2\text{O}_5$ ,  $\text{N}_2\text{O}_4$  and  $\text{NO}_2$ ," J. Amer. Chem. Soc. 56, 630-637 (1934)
- Holt, R. B., McLane, C. K., and Eldenberg, S., "Ultraviolet Absorption Spectrum of Hydrogen Peroxide," J. Chem. Phys. 16, 225-229 (1948).  
Erratum: 16, 638 (1948)
- Horowitz, A., and Calvert, J. G., "The Wavelength Dependence of the Quantum Efficiencies of the Primary Processes in Formaldehyde Photolysis at 25°C," typescript, to be published Int. J. Chem. Kinet. (1978a)
- Horowitz, A., and Calvert, J. G., "The Quantum Efficiency of the Primary Processes in Formaldehyde Photolysis at 3130Å and 25°C," typescript, to be published Int. J. Chem. Kinet. (1978b)
- Houston, P. L., and Moore, C. B., "Formaldehyde Photochemistry: Appearance Rate, Vibrational Relaxation, and Energy Distribution of the CO Product," J. Chem. Phys. 65, 757-770 (1976)
- Howard, C. J., "Rate Constants for the Gas-Phase Reactions of OH Radicals with Ethylene and Halogenated Ethylene Compounds," J. Chem. Phys. 65, 4771-4777 (1976)
- Howard, C. J., "Kinetics of the Reaction of  $\text{HO}_2$  with  $\text{NO}_2$ ," J. Chem. Phys. 67, 5258-5263 (1977)
- Howard, C. J., and Evenson, K. M., "Laser Magnetic Resonance Study of the Gas Phase Reactions of OH with CO, NO, and  $\text{NO}_2$ ," J. Chem. Phys. 61, 1943-1952 (1974)
- Howard, C. J., and Evenson, K. M., "Rate Constants for the Reactions of OH with Ethane and some Halogen Substituted Ethanes at 296 K," J. Chem. Phys. 64, 4303-4306 (1976)
- Howard, C. J., and Evenson, K. M., "Rate Constants for the Reactions OH with  $\text{CH}_4$  and Fluorine, Chlorine, and Bromine Substituted Methanes at 296 K," J. Chem. Phys. 64, 197-202 (1976)
- Howard, C. J., and Evenson, K. M., "Kinetics of the Reaction of  $\text{HO}_2$  with NO," Geophys. Res. Lett. 4, 437-440 (1977)
- Huestis, D. L., Black, G., Edelstein, S. A., and Sharpless, R. L., "Fluorescence and Quenching of  $\text{O}_2(^1\Delta_g)$  and  $[\text{O}_2(^1\Delta_g)]_2$  in Liquid Oxygen," J. Chem. Phys. 60, 4471-4474 (1974)
- Hui, K.-K., and Cool, T. A., "Experiments Concerning the Laser Enhanced Reaction Between Vibrationally Excited  $\text{O}_3$  and NO," J. Chem. Phys. 68, 1022 (1978)
- Huie, R. E., and Herron, J. T., "Temperature Dependence of the Rate Constants for Reactions of Ozone with Some Olefins," Int. J. Chem. Kinet., Symp. No. 1, 165-181 (1975)
- Huie, R. E., and Herron, J. T., "The Rate Constant for the Reaction  $\text{O}_3 + \text{NO}_2 \rightarrow \text{O}_2 + \text{NO}_3$  Over the Temperature Range 259-362° K," Chem. Phys. Lett. 27, 411-414 (1974a)



- Huie, R. E., Herron, J. T., and Davis, D. D., "Absolute Rate Constants for the Reaction  $O + O_2 + M \rightarrow O_3 + M$  Over the Temperature Range 200-346°K," J. Phys. Chem. 76, 2653-2658 (1972)
- Husain, D., Kirsch, L. J., and Donovan, R. J., "A Kinetic Study of  $O(2^3P_J)$  by Atomic Absorption Spectroscopy Following the Flash Photolysis of Ozone," J. Photochem. 1, 69-73 (1972)
- Jacob, A., and Winkler, C. A., "Kinetics of the Reactions of Oxygen Atoms and Nitrogen Atoms with Sulphur Trioxide," J. Chem. Soc., Faraday Trans. I 68, 2077-2082 (1972)
- James, F. C., Kerr, J. A., and Simons, J. P., "A Direct Measurement of the Rate of Reaction of the Methyl Radical with Sulphur Dioxide," J. Chem. Soc. Faraday Trans. I 69, 2124-2129 (1973)
- Japar, S. M., Wu, C. H., and Niki, H., "Rate Constants for the Reaction of Ozone with Olefins in the Gas Phase," J. Phys. Chem. 78, 2318 (1974)
- Jayanty, R. K. M., Simonaitis, R., and Heicklen, J., "The Reaction of  $NH_2$  with  $NO_2$ ," (Department of Chemistry and Ionosphere Research Laboratory, Penn. State Univ., Pa. 16802, 1975d)
- Jayanty, R. K. M., Simonaitis, R., and Heicklen, J., "The Reaction of  $NH_2$  with  $NO$  and  $O_2$ ," J. Phys. Chem. 80, 433-437 (1976b)
- Johnston, H. S., "Gas Phase Reaction Rate Theory," The Ronald Press Co., New York (1966), Chapter 1
- Johnston, H. S., "Gas Phase Reaction Kinetics of Neutral Oxygen Species," NBS-NSRDS-20, 1968. (Supt. of Documents, U.S. Govt. Printing Office, Washington, D.C. 20402)
- Johnston, H. S., "Catalytic Reduction of Stratospheric Ozone by Nitrogen Oxides," Adv. Environ. Sci. Technol. 4, 263-380 (1974), J. N. Pitts, Jr., and R. L. Metcalf, editors, Wiley-Interscience
- Johnston, H. S., private communication (1973)
- Johnston, H. S., Chang, S.-G., and Whitten, G., "Photolysis of Nitric Acid Vapor," J. Phys. Chem. 78, 1-7 (1974)
- Johnston, H. S., and Graham, R., "Gas Phase Ultraviolet Absorption Spectrum of Nitric Acid Vapor," J. Phys. Chem. 77, 62-63 (1973)
- Johnston, H. S., and Graham, R., "Photochemistry of  $NO_x$  and  $HN O_x$  Compounds," Can. J. Chem. 52, 1415-1423 (1974)
- Johnston, H. S., Morris, E. D., Jr., and Van den Bogaerde, J., "Molecular Modulation Kinetic Spectrometry.  $ClO$  and  $ClO_2$  Radicals in the Photolysis of Chlorine in Oxygen," J. Amer. Chem. Soc. 91, 7712-7727 (1969)
- Johnston, H. S., and Selwyn, G. S., "New Cross Sections for the Absorption of Near Ultraviolet Radiation by Nitrous Oxide ( $N_2O$ )," Geophys. Res. Lett. 2, 549-551 (1975)
- Jones, I. T. N., and Bayes, K. D., "Photolysis of Nitrogen Dioxide," J. Chem. Phys. 59, 4836-4844 (1973)
- Jones, E. J., and Wulf, G. R., "Absorption Coefficient of Nitrogen Pentoxide in the Ultraviolet and the Visible Absorption Spectrum of  $NO_3$ ," J. Chem. Phys. 5, 873-877 (1937)
- Kajimoto, G., and Cvetanović, R. J., "Formation of Nitrous Oxide in the Reaction of  $O(^1D_2)$  Atoms with Nitrogen," J. Chem. Phys. 64, 1005-1015 (1976)
- Kerr, J. A., and Parsonage, M. J., "Evaluated Kinetic Data on Gas Phase Addition Reactions: Reactions of Atoms and Radicals with Alkenes, Alkynes and Aromatic Compounds," (Butterworths, 1972)

- Klemm, R. B., and Stief, L. J., "Absolute Rate Parameters for the Reaction of Ground State Atomic Oxygen with Carbonyl Sulfide," *J. Chem. Phys.* **61**, 4900-4906 (1974)
- Klemm, R. B., Payne, W. A., and Stief, L. J., "Absolute Rate Parameters for the Reaction of Atomic Hydrogen with  $H_2O_2$ ," *Int. J. Chem. Kinet.*, Symp. No. 1, 61-72 (1975)
- Kondratiev, V. N., "Konstanty Skorosti Gazofaznykh Reaktsij Spravochnik," (Izdatelstvo "Nauka", Moskva, 1970); also issued as "Rate Constants of Gas Phase Reactions, Reference Book," R. M. Fristrom, Ed., Translated by L. J. Holtschlag CDM-72-10014, Office of Standard Reference Data, NBS (1972) (Distributed by National Technical Information Service, Springfield, Va. 22151)
- Kurylo, M. J., Hollinden, G. A., LeFevre, H. F., and Timmons, R. B., "ESR Study of the Kinetics of the Reactions of D Atoms and O Atoms with  $NH_3$ ," *J. Chem. Phys.* **51**, 4497-4501 (1969)
- Kurylo, M. J., "Absolute Rate Constants for the Reaction  $H + O_2 + M \rightarrow HO_2 + M$  over the Temperature Range 203-404 K," *J. Phys. Chem.* **76**, 3518-3526 (1972)
- Kurylo, M. J., "Kinetics of the Reactions  $OH(v = 0) + NH_3 \rightarrow H_2O + NH_2$  and  $OH(v = 0) + O_3 \rightarrow HO_2 + O_2$  at 298°K," *Chem. Phys. Letts.* **23**, 467-471 (1973)
- Kurylo, M. J., "Flash Photolysis Resonance Fluorescence Investigation of the Reaction of  $O(^3P)$  Atoms with  $ClONO_2$ ," *Chem. Phys. Lett.* **49**, 467-470 (1977)
- Kurylo, M. J., and Braun, W., "Flash Photolysis Resonance Fluorescence Study of the Reaction  $Cl + O_3 \rightarrow ClO + O_2$  over the Temperature Range 213-298 K," *Chem. Phys. Lett.* **37**, 232-235 (1976)
- Kurylo, M. J., and Manning, R. G., "Flash Photolysis Resonance Fluorescence Investigation of the Reaction of  $Cl(^2P)$  Atoms with  $ClONO_2$ ," *Chem. Phys. Lett.* **48**, 279-283 (1977)
- Kurylo, M. J., Braun, W., and Xuan, C. N., "Infrared Laser Enhanced Reactions: Temperature Resolution of the Chemical Dynamics of the  $O_3 + NO$  Reaction System," *J. Chem. Phys.* **62**, 2065-2071 (1975)
- Kurylo, M. J., Peterson, N. C., and Braun, W., "Absolute Rate of the Reaction  $H + H_2S$ ," *J. Chem. Phys.* **54**, 943-946 (1971)
- Laufer, A. H., and Bass, A. M., "Rate Constants of the Combination of Methyl Radicals with Nitric Oxide and Oxygen," *Int. J. Chem. Kinet.* **7**, 639-648 (1975)
- Lee, J. H., Michael, J. V., Payne, W. A., and Stief, L. J., "Absolute Rate of the Reaction of Hydrogen Atoms with Ozone from 219-360 K," paper presented at 13th Informal Conference on Photochemistry, Clearwater Beach, Florida, Jan. 1978
- Lee, J. H., Michael, J. V., Payne, W. A., Stief, L. J., and Whytock, D. A., "Absolute Rate of the Reaction of  $Cl(^2P)$  with Molecular Hydrogen from 200-500 K," *J. Chem. Soc., Faraday Trans. I* **73**, 1530-1536 (1977)
- Lesclaux, R., van Khe, P., Dezaudier, P., and Soulignac, J. C., "Flash Photolysis Studies of the Reaction of  $NH_2$  Radicals with  $NO$ ," *Chem. Phys. Lett.* **35**, 493-497 (1975)
- Leu, M.-T., and DeMore, W. B., "Rate Constants at 295 K for the Reactions of Atomic Chlorine with  $H_2O_2$ ,  $HO_2$ ,  $O_3$ ,  $CH_4$  and  $HNH_3$ ," *Chem Phys. Lett.* **41**, 121-124 (1976)
- Leu, M.-T., and DeMore, W. B., "Rate Constant for the Reaction of Atomic Bromine with Ozone," *Chem. Phys. Lett.* **48**, 317-320 (1977)

- Leu, M. T., Lin, C. L., and DeMore, W. B., "Rate Constant for Formation of Chlorine Nitrate by the Reaction  $\text{ClO} + \text{NO}_2 + \text{M}$ ," J. Phys. Chem. 81, 190-195 (1977)
- Lewis, R. S., Tang, K. Y., and Lee, E. K. C., "Photoexcited Chemiluminescence Spectroscopy: Detection of Hydrogen Atoms Produced from Single Vibronic Level Photolysis of Formaldehyde ( $\tilde{\text{A}}^1\text{A}_2$ )," J. Chem. Phys. 65, 2910-2911 (1976)
- Lewis, R. S., and Lee, E. K. C., "Photolysis of Formaldehyde-Butene Mixtures at Low Pressures: Quantum Yield of Radical Decomposition Process in Formaldehyde," typescript, Dept. of Chemistry, Univ. of California, Irvine, CA (1977)
- Lin, C. L., and DeMore, W. B., " $\text{O}(^1\text{D})$  Production in Ozone Photolysis Near  $3100\text{\AA}$ ," J. Photochem. 2, 161-164 (1973)
- Lloyd, A. C., "Evaluated and Estimated Kinetic Data for Gas Phase Reactions of the Hydroperoxyl Radical," Int. J. Chem. Kinet. 6, 169-228 (1974) Supersedes NBS Report 10447
- Lundell, O. R., Ketcheson, R. D., and Schiff, H. I., "The Production of  $\text{O}(^3\text{P})$  Atoms, Free From Excited Molecules, and Their Reaction with  $\text{O}_3$ ," Symp. Combust. 12th (Combustion Institute, Pittsburgh, 1969) 307-311
- Mack, G. P. R., and Thrush, B. A., "Reaction of Oxygen Atoms with Carbonyl Compounds. Part 1. Formaldehyde," J. Chem. Soc., Faraday Trans. I 69, 208-215 (1973)
- Mandelman, M., and Carrington, T., "The f-value of the  $\text{NO } \delta(0, 0)$  Band by the Line Absorption Method," J. Quant. Spectrosc. Radiat. Transfer 14, 509-521 (1974)
- Manning, R. G., Braun, W., and Kurylo, M. J., "The Effect of Infrared Laser Excitation on Reaction Dynamics:  $\text{O} + \text{C}_2\text{H}_4^\dagger$  and  $\text{O} + \text{OCS}^\dagger$ ," J. Chem. Phys. 65, 2609-2615 (1976)
- Manning, R. G., and Kurylo, M. J., "Flash Photolysis Resonance Fluorescence Investigation of the Temperature Dependencies of the Reaction of  $\text{Cl}(^2\text{P})$  Atoms with  $\text{CH}_4$ ,  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_3\text{F}$ ,  $\text{CH}_3\text{F}^\ddagger$ , and  $\text{C}_2\text{H}_6$ ," J. Phys. Chem. 81, 291-296 (1977)
- Margitan, J. J., Kaufman, F., and Anderson, J. G., "The Reaction of  $\text{OH}$  with  $\text{CH}_4$ ," Geophys. Res. Lett. 1, 80-81 (1974)
- Margitan, J. J., Kaufman, F., and Anderson, J. G., "Kinetics of the Reaction  $\text{OH} + \text{HNO}_3 \rightarrow \text{H}_2\text{O} + \text{NO}_3$ ," Int. J. Chem. Kinet., Symp. No. 1, 281-287 (1975)
- Marling, J., "Isotope Separation of Oxygen-17, Oxygen-18, Carbon-13, and Deuterium by Ion Laser Induced Formaldehyde Photopredissociation," J. Chem. Phys. 66, 4200-4225 (1977)
- McCrumb, J. L., and Kaufman, F., "Kinetics of the  $\text{O} + \text{O}_3$  Reaction," J. Chem. Phys. 57, 1270-1276 (1972)
- McKenzie, A., Mulcahy, M. F. R., and Steven, J. R., "Kinetics of Decay of Hydroxyl Radicals at Low Pressure," J. Chem. Phys. 59, 3244-3254 (1973)
- McQuigg, R. D., and Calvert, J. G., "The Photodecomposition of  $\text{CH}_2\text{O}$ ,  $\text{CD}_2\text{O}$ ,  $\text{CHDO}$  and  $\text{CH}_2\text{O}-\text{CD}_2\text{O}$  Mixtures at Xenon Flash Lamp Intensities," J. Amer. Chem. Soc. 91, 1590-1599 (1969)
- Meaburn, G. M., Perner, D., LeCalve, J., and Bourene, M., "A Pulsed-Radiolysis Study of Atomic Oxygen Reactions in the Gas Phase," J. Phys. Chem. 72, 3920-3925 (1968)
- Meagher, J. F., and Heicklen, J., "The Photolysis of Hydrogen Peroxide in the Presence of Carbon Monoxide," J. Photochem. 3, 455-466 (1975)



- Michael, J. V., Payne, W. A., and Whytock, D. A., "Absolute Rate Constants for  $\text{O} + \text{N}_2 + \text{M}$  ( $\text{M} = \text{He, Ne, Ar, Kr}$ )  $\rightarrow \text{N}_2\text{O} + \text{M}$  from 217 to 500 K," J. Chem. Phys. 65, 4830-4834 (1976)
- Milks, D., and Matula, R. A., "A Single-Pulse Shock-Tube Study of the Reaction between Nitrous Oxide and Carbon Monoxide," Symp. Combust. 14th (Combustion Institute, Pittsburgh, 1973) 83-96
- Miller, R. G., and Lee, E. K. C., "Single Vibronic Level Photochemistry of Formaldehydes in the  $\tilde{\text{X}}^1\text{A}_2$  State: Radiative and Non-Radiative Processes in  $\text{H}_2\text{CO}$ ,  $\text{HDCO}$  and  $\text{D}_2\text{CO}$ ," typescript, Dept. of Chemistry, Univ. of California, Irvine, CA (1977)
- Molina, L. T., Spencer, J. E., and Molina, M. J., "The Rate constant for the Reaction of  $\text{O}(^3\text{P})$  Atoms with  $\text{ClONO}_2$ ," Chem. Phys. Lett. 45, 168-162 (1977)
- Molina, M. J. (1977), "Cross Sections and Quantum Yields," in Chapter 1 of NASA Reference Publication 1010 "Chlorofluoromethanes and the Stratosphere", R. D. Hudson, Ed., August 1977
- Moortgat, G. K., and Allen, E. R., "The Room Temperature Reactions of Hydrogen-Atoms in Molecular Oxygen," National Center for Atmospheric Research, Boulder (1973). Abstract of unpublished work.
- Moortgat, G. K., Slemr, F., and Warneck, P., "Kinetics and Mechanism of the Reaction  $\text{H} + \text{CH}_3\text{ONO}$ ," Int. J. Chem. Kinet. 9, 249-265 (1977)
- Moortgat, G. K., Slemr, F., Seiler, W., and Warneck, P., "Wavelength Dependence of Relative and Absolute Quantum Yields in the Near UV Photolysis of Formaldehyde at 25°C," paper presented at 13th Informal Conference on Photochemistry, Clearwater Beach, Florida, Jan. (1978)
- Morley, C., and Smith, I. W. M., "Rate Measurements of Reactions of OH by Resonance Absorption. Part I. Reactions of OH with  $\text{N}_2\text{O}$  and  $\text{NO}$ ," J. Chem. Soc., Faraday Trans. II 68, 1016-1030 (1972)
- Morris, E. D., Jr., and Niki, H., "Mass Spectrometric Study of the Reaction of Hydroxyl Radical with Formaldehyde," J. Chem. Phys. 55, 1991-1992 (1971)
- Morris, E. D., Jr., and Niki, H., "Reaction of Methyl Radicals with Atomic Oxygen," Int. J. Chem. Kinet. 5, 47-53 (1972)
- Morris, E. D., Jr., and Niki, H., "Reaction of Dinitrogen Pentoxide with Water," J. Phys. Chem. 77, 1929-1932 (1973)
- Morris, E. D., Jr., Stedman, D. H., and Niki, H., "Mass Spectrometric Study of the Reactions of the Hydroxyl Radical with Ethylene, Propylene, and Acetaldehyde in a Discharge-Flow System," J. Amer. Chem. Soc. 93, 3570-3572 (1971)
- Moy, J., Bar-Ziv, E., and Gordon, R. J., "Temperature Dependence of the Laser Enhanced Reaction  $\text{NO} + \text{O}_3(001) \rightarrow \text{NO}_2(^2\text{B}_{1,2}) + \text{O}_2$ ," J. Chem. Phys. 66, 5439-5446 (1977)
- Mulcahy, M. F. R., and Williams, D. J., "Kinetics of Combination of Oxygen Atoms with Oxygen Molecules," Trans. Faraday Soc. 64, 59-70 (1968)
- Mulvihill, J. N., and Phillips, L. F., "Products of Reaction of NH with  $\text{NO}$ ," Chem. Phys. Lett. 35, 327-329 (1975)
- NASA (1977). "Chlorofluoromethanes and the Stratosphere," R. D. Hudson, Ed., NASA Reference Publication 1010, August 1977, Chapter 1. Recommendations of the Laboratory Measurements Committee for the NASA chlorofluoromethane assessment.
- Nicholas, J. E., and Norrish, R. G. W., "Some Reactions in the Chlorine and Oxygen System Studied by Flash Photolysis," Proc. Roy. Soc. (London) A 307, 391-397 (1968)

- Ogryzlo, E. A., and Thrush, B. A., "Vibrational Excitation of NO in its Collisional Quenching of  $\text{O}_2(^1\Delta_g)$ ," Chem. Phys. Lett. 23, 34-36 (1973)
- Overend, R., and Paraskevopoulos, G., "The Question of a Pressure Effect in the Reaction  $\text{OH} + \text{CO}$  at Room Temperature," Chem. Phys. Lett. 49, 109-111 (1977)
- Overend, R., and Paraskevopoulos, G., "Rates of OH Radical Reactions. III. The Reaction  $\text{OH} + \text{C}_2\text{H}_4 + \text{M}$  at 256°K," J. Chem. Phys. 67, 674-679 (1977)
- Overend, R., Paraskevopoulos, G., and Black, C., "Rates of OH Radicals Reactions. II. The Combination Reaction  $\text{OH} + \text{NO} + \text{M}$ ," J. Chem. Phys. 64, 4149-4154 (1976)
- Overend, R., Paraskevopoulos, G., and Cvetanović, R. J., "Rates of OH Radical Reactions. I. Reactions with  $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{C}_2\text{H}_6$ , and  $\text{C}_3\text{H}_8$  at 295 K," Can. J. Chem. 53, 3374-3382 (1975)
- Parkes, D. A., "The Roles of Alkylperoxy and Alkoxy Radicals in Alkyl Radical Oxidation at Room Temperature," Symp. Combust. 15th (Combustion Institute, Pittsburgh, 1975) 795-805
- Parkes, D. A., "Oxidation of Methyl Radicals at Room Temperature," Int. J. Chem. Kinet. 9, 451-469 (1977)
- Pastrana, A., and Carr, R. W., Jr., "Rate of the Reaction of Hydroxyl Radical with Acetylene," Int. J. Chem. Kinet. 6, 587-595 (1974)
- Pate, C. T., Finlayson, B. J., and Pitts, J. N., Jr., "A Long Path Infrared Spectroscopic Study of the Reaction of Methylperoxy Free Radicals with Nitric Oxide," J. Am. Chem. Soc. 96, 6554-6558 (1974)
- Payne, W. A., Stief, L. J., and Davis, D. D., "A Kinetics Study of the Reaction of  $\text{HO}_2$  with  $\text{SO}_2$  and  $\text{NO}$ ," J. Amer. Chem. Soc. 95, 7614-7619 (1973)
- Peeters, J., and Mahnen, G., "Reaction Mechanisms and Rate Constants of Elementary Steps in Methane-Oxygen Flames," Symp. Combust. 14th (Combustion Institute, Pittsburgh, 1973) 133-141
- Penzhorn, R-D., Gusten, H., Schurath, U., and Becker, K. H., "Quenching of Singlet Molecular Oxygen by Some Atmospheric Pollutants," Environ. Sci. Technol. 8, 907 (1974)
- Penzhorn, R-D., Gusten, H., Schurath, U., and Becker, K. H., "Halogenierte Kohlenwasserstoffe in der Atmosphäre: Geschwindigkeitskonstanten für die Löschung von Metastabilen Sauerstoffmolekülen," Staub - Reinhalt Luft 35, 95-98 (1975)
- Perry, R. A., Atkinson, R., and Pitts, J. N., Jr., "Rate Constants for the Reaction of OH Radicals with  $\text{CH}_2\text{Cl}_2$  and  $\text{CH}_3\text{Cl}$  over the Temperature Range 298-423°K, and with  $\text{CH}_2\text{Cl}_2$  at 298°K," J. Chem. Phys. 64, 1618-1620 (1976)
- Perry, R. A., Atkinson, R., and Pitts, J. N., Jr., "Rate Constants for the Reactions  $\text{OH} + \text{H}_2\text{S} \rightarrow \text{H}_2\text{O} + \text{SH}$  and  $\text{OH} + \text{NH}_3 \rightarrow \text{H}_2\text{O} + \text{NH}_2$  over the Temperature Range 297-427°K," J. Chem. Phys. 64, 3237-3239 (1976a)
- Perry, R. A., Atkinson, R., and Pitts, J. N., Jr., "Rate Constants for the Reaction of OH Radicals with n-Butane over the Temperature Range 297-420°K," J. Chem. Phys. 64, 5314-5316 (1976b)
- Perry, R. A., Atkinson, R., and Pitts, J. N., Jr., "Kinetics and Mechanism of the Gas Phase Reaction of OH Radicals with Aromatic Hydrocarbons over the Temperature Range 296-473 K," J. Phys. Chem. 81, 296-304 (1977)
- Perry, R. A., Atkinson, R., and Pitts, J. N., Jr., "Kinetics of the Reactions of OH Radicals with  $\text{C}_2\text{H}_2$  and  $\text{CO}$ ," J. Chem. Phys. 67, 5577-5584 (1977a)



- Phillips, L. F., and Schiff, H. I., "Mass Spectrometric Studies of Atomic Reactions. III. Reactions of Hydrogen Atoms with Nitrogen Dioxide and with Ozone," J. Chem. Phys. 37, 1233-1238 (1962)
- Pruss, F. J., Jr., Slagle, I. R., and Gutman, D., "Determination of Branching Ratios for the Reaction of Oxygen Atoms with Ethylene," J. Phys. Chem. 78, 663-665 (1974)
- Ravishankara, A. R., Davis, D. D., Smith, G., Tesi, G., and Spencer, J., "A Study of the Chemical Degradation of  $\text{ClONO}_2$  in the Stratosphere," Geophys. Res. Lett. 4, 7-9 (1977)
- Rawlins, W. T., and Gardiner, W. C., Jr., "Rate Constant of  $\text{OH} + \text{OH} = \text{H}_2\text{O} + \text{O}$  from 1500 to 2000 K," J. Chem. Phys. 60, 4676-4681 (1974)
- Ridley, B. A., Davenport, J. A., Stief, L. J., and Welge, K. H., "Absolute Rate Constant for the Reaction  $\text{H} + \text{H}_2\text{CO}$ ," J. Chem. Phys. 57, 520-523 (1972)
- Rowland, F. S., Spencer, J. E., and Molina, M. J., "Stratospheric Formation and Photolysis of Chlorine Nitrate," J. Phys. Chem. 80, 2711-2713 (1976)
- Sauer, M. C., Jr., "A Pulse-Radiolysis Study of the Dependence of the Reaction of Atomic Oxygen with Oxygen on the Nature of the Third Body," J. Phys. Chem. 71, 3311-3313 (1967)
- Schmatjko, K. J., and Wolfrum, J., "Reaction of Flash Photolytically Produced  $\text{CN}(\text{X}^2\Sigma^+, \text{v})$  Radicals with  $\text{O}(^3\text{P})$  Atoms," Photochem. Conference 12th (National Bureau of Standards, Washington, D. C. 20234 1976) R5-1
- Schmidt, C., and Schiff, H. I., "Reactions of  $\text{O}_2(^1\Delta_g)$  with Atomic Nitrogen and Hydrogen," Chem. Phys. Lett. 23, 339-342 (1973)
- Schofield, K., "The Rate of Destruction of  $\text{O}_2(^1\Delta_g)$  by Atomic Hydrogen," Intern. J. Chem. Kinet. 4, 255-263 (1972)
- Schofield, K., "Evaluated Chemical Kinetic Rate Constants for Various Gas Phase Reactions," J. Phys. Chem. Ref. Data 2, 25-84 (1973)
- Schott, G. L., Getzinger, R. W., and Seitz, W. A., "Transient Oxygen Atom Yields in  $\text{H}_2\text{O}_2$  Ignition and the Rate Coefficient for  $\text{O} + \text{H}_2 \rightarrow \text{OH} + \text{H}$ ," Int. J. Chem. Kinet. 6, 921-943 (1974)
- Selwyn, G., Podolske, J., and Johnston, H. S., "Nitrous Oxide Ultraviolet Absorption Spectrum at Stratospheric Temperatures," Geophys. Res. Lett. 4, 427-430 (1977)
- Sie, B. K. T., Simonaitis, R., and Heicklen, J., "The Reaction of  $\text{OH}$  with  $\text{CO}$ ," Int. J. Chem. Kinet. 8, 85-98 (1976a)
- Sie, B. K. T., Simonaitis, R., and Heicklen, J., "The Reaction of  $\text{OH}$  with  $\text{NO}$ ," Int. J. Chem. Kinet. 8, 99-106 (1976b)
- Simonaitis, R., and Heicklen, J., "The Reactions of  $\text{HO}_2$  with Carbon Monoxide and Nitric Oxide and the Reaction of  $\text{O}(^1\text{D})$  with Water," J. Phys. Chem. 77, 1096-1102 (1973a)
- Simonaitis, R., and Heicklen, J., "The Reaction of  $\text{HO}_2$  with  $\text{O}_3$ ," J. Phys. Chem. 77, 1932-1935 (1973b)
- Simonaitis, R., and Heicklen, J., "The Mechanism of the  $\text{H}_2\text{O}$ -Catalyzed Chain Photodecomposition of  $\text{O}_3$  and the Reaction of  $\text{HO}$  with  $\text{O}_3$ ," J. Photochem. 2, 309-314 (1973c)
- Simonaitis, R., and Heicklen, J., "The Reaction of  $\text{HO}_2$  with  $\text{NO}$  and  $\text{NO}_2$ ," J. Phys. Chem. 78, 653-657 (1974)
- Simonaitis, R., and Heicklen, J., "Reactions of  $\text{CH}_3\text{O}_2$  with  $\text{NO}$  and  $\text{NO}_2$ ," J. Phys. Chem. 78, 2417-2421 (1974)
- Simonaitis, R., and Heicklen, J., "Reactions of  $\text{CH}_3$ ,  $\text{CH}_3\text{O}$ , and  $\text{CH}_3\text{O}_2$  Radicals with  $\text{O}_3$ ," J. Phys. Chem. 79, 298-302 (1975)
- Simonaitis, R., and Heicklen, J., "Reactions of  $\text{HO}_2$  with  $\text{NO}$  and  $\text{NO}_2$  and of  $\text{OH}$  with  $\text{NO}$ ," J. Phys. Chem. 80, 1-7 (1976a)

- Simonaitis, R., and Heicklen, J., "The Temperature Dependence of the Reactions of  $\text{H}\dot{\text{O}}_2$  with  $\text{NO}$  and  $\text{NO}_2$ ," *Int. J. Chem. Kinet.* 10, 67-87 (1978)
- Singleton, D. L., and Cvetanović, R. J., "Temperature Dependence of the Reactions of Oxygen Atoms with Olefins," *J. Am. Chem. Soc.* 98, 6812-6819 (1976)
- Singleton, D. L., Furuyama, S., Cvetanović, R. J., and Irwin, R. S., "Temperature Dependence of the Rate Constants for the Reactions  $\dot{\text{O}}(^3\text{P}) + 2,3\text{-Dimethyl-2-Butene}$  and  $\dot{\text{O}}(^3\text{P}) + \text{NO} + \text{M}$  Determined by a Phase Shift Technique," *J. Chem. Phys.* 63, 1003-1007 (1975)
- Slack, M. W., "Rate Coefficient for  $\text{H} + \dot{\text{O}}_2 + \text{M} \rightarrow \text{HO}_2 + \text{M}$  Evaluated from Shock Tube Measurements of Induction Times," *Combust. Flame* 28, 241-249 (1977)
- Slagle, I. R., Pruss, F. J., Jr., and Gutman, D., "Kinetics into the Steady State. I. Study of the Reaction of Oxygen Atoms with Methyl Radicals," *Int. J. Chem. Kinet.* 6, 111-123 (1974)
- Slagle, I. R., Gilbert, J. R., and Gutman, D., "Kinetics of the Reaction between Oxygen Atoms and Carbon Disulfide," *J. Chem. Phys.* 62, 740-709 (1974)
- Slagle, I. R., Graham, R. E., Gilbert, J. R., and Gutman, D., "Direct Determination of the Rate Constant for the Reaction of Oxygen Atoms with Carbon Monosulphide," *Chem. Phys. Lett.* 32, 184-186 (1975)
- Slanger, T. G., and Black, G., "Reaction Rate Measurements of  $\dot{\text{O}}(^3\text{P})$  Atoms by Resonance Fluorescence. I.  $\dot{\text{O}}(^3\text{P}) + \dot{\text{O}}_2 + \text{M} \rightarrow \dot{\text{O}}_3 + \text{M}$  and  $\dot{\text{O}}(^3\text{P}) + \text{NO} + \text{M} \rightarrow \text{NO}_2 + \text{M}$ ," *J. Chem. Phys.* 53, 3717-3721 (1970)
- Slanger, T. G., and Black, G., " $\dot{\text{O}}(^1\text{S})$  Quenching by  $\dot{\text{O}}(^3\text{P})$ ," *J. Chem. Phys.* 64, 3763-3766 (1976a)
- Slanger, T. G., and Black, G., "Temperature Dependence for Quenching of  $\dot{\text{O}}(^1\text{S})$  by  $\text{N}_2\dot{\text{O}}$ ," *J. Chem. Phys.* 65, 2025-2026 (1976b)
- Slanger, T. G., Wood, B. J., and Black, G., "Investigation of the Rate Coefficient for  $\dot{\text{O}}(^3\text{P}) + \text{NO}_2 \rightarrow \dot{\text{O}}_2 + \text{NO}$ ," *Int. J. Chem. Kinet.* 5, 615-620 (1973)
- Slanger, T. G., and Welge, K. H., "Rate Constants for Reactions of  $\dot{\text{O}}(^1\text{S})$ ," in *Chemical Kinetics Data Survey V. Sixty-six Contributed Rate and Photochemical Data Evaluations on Ninety-four Reactions*, NBSIR 73-206, D. Garvin, Ed., National Bureau of Standards, Washington, D.C., (1973) pages 22-32
- Slemr, F., and Warneck, P., "Kinetics of the Reaction of Atomic Hydrogen with Methylhydroperoxide," *Int. J. Chem. Kinet.* 9, 267-282 (1977)
- Smith, I. W. M., and Zellner, R., "Rate Measurements of Reactions of  $\dot{\text{O}}\text{H}$  by Resonance Absorption Part 2. Reactions of  $\dot{\text{O}}\text{H}$  with  $\text{CO}$ ,  $\text{C}_2\text{H}_4$  and  $\text{C}_2\text{H}_2$ ," *J. Chem. Soc., Faraday Trans. II* 69, 1617-1627 (1973)
- Smith, I. W. M., and Zellner, R., "Rate Measurements of Reactions of  $\dot{\text{O}}\text{H}$  by Resonance Absorption. Part 3. Reactions of  $\dot{\text{O}}\text{H}$  with  $\text{H}_2$ ,  $\text{D}_2$ , Hydrogen and Deuterium Halides," *J. Chem. Soc., Faraday Trans. II* 70, 1045-1056 (1974a)
- Smith, I. W. M., and Zellner, R., "Rate Measurements of Reactions of  $\dot{\text{O}}\text{H}$  by Resonance Absorption. IV. - Reactions of  $\dot{\text{O}}\text{H}$  with  $\text{NH}_3$  and  $\text{HNO}_3$ ," *Int. J. Chem. Kinet., Symp. No. 1*, 341-351 (1975)
- Spencer, J. E., and Glass, G. P., "A Reexamination of the Reaction  $\text{H} + \text{HCl} \rightleftharpoons \text{H}_2 + \text{Cl}$ ," *J. Phys. Chem.* 79, 2329-2335 (1975)
- Sperling, H. P., and Toby, S., "The Photochemical Decomposition of Gaseous Formaldehyde," *Can. J. Chem.* 51, 471-475 (1973)
- Stedman, D. H., and Niki, H., "Kinetics and Mechanism for the Photolysis of Nitrogen Dioxide in Air," *J. Phys. Chem.* 77, 2604-2609 (1973)

- Stedman, D. H., Wu, C. H., and Niki, H., "Kinetics of Gas-Phase Reactions of Ozone with Some Olefins," J. Phys. Chem. 77, 2511-2514 (1973)
- Steinert, W., and Zellner, R., "Rates of Reaction of OH with CO and CH<sub>4</sub> over an Extended Temperature Range," Deuxieme Symp. European sur la Combustion (Orleans, 1975) 31-37
- Stief, L. J., and Payne, W. A., "Absolute Rate Parameters for the Reaction of Atomic Hydrogen with Hydrazine," J. Chem. Phys. 64, 4892-4896 (1976)
- Streit, G. E., and Johnston, H. S., "Reactions and Quenching of Vibrationally Excited Hydroxyl Radicals," J. Chem. Phys. 64 95-103 (1976)
- Streit, G. E., Howard, C. J., Schmeltekopf, A. L., Davidson, J. A., and Schiff, H. I., "Temperature Dependence of O(<sup>1</sup>D) Rate Constants for Reactions with O<sub>2</sub>, N<sub>2</sub>, CO<sub>2</sub>, O<sub>3</sub>, and H<sub>2</sub>O," J. Chem. Phys. 65, 4761-4764 (1976)
- Stuhl, F., "Absolute Rate Constant for the Reaction OH + NH<sub>3</sub> → NH<sub>2</sub> + H<sub>2</sub>O," J. Chem. Phys. 59, 535-537 (1973a)
- Stuhl, F., "Rate Constant for the Reaction of OH with n-C<sub>4</sub>H<sub>10</sub>," Z. Naturforsch. Pt A 28, 1383-1384 (1973b)
- Stuhl, F., "Determination of Rate Constants for the Reactions of OH with Propylene and Ethylene by a Pulsed Photolysis-Resonance Fluorescence Method," Ber. Bunsenges. Physik. Chem. 77, 674-677 (1973c).
- Stuhl, F., "Determination of the Rate Constant for the Reaction OH + H<sub>2</sub>S by a Pulsed Photolysis - Resonance Fluorescence Method," Ber. Bunsenges. Physik. Chem. 78, 230-232 (1974)
- Stuhl, F., and Niki, H., "Detection of Oxygen (<sup>3</sup>P) Atoms in Pulsed Vacuum UV Photolysis of NO and Its Application to O Atom Reactions," Chem. Phys. Lett. 7, 197-200 (1970)
- Stuhl, F., and Niki, H., "Measurements of Rate Constants for Termolecular Reactions of O(<sup>3</sup>P) with NO, O<sub>2</sub>, CO, N<sub>2</sub>, and CO<sub>2</sub> Using a Pulsed Vacuum-uv Photolysis-Chemiluminescent Method," J. Chem. Phys. 55, 3943-3953 (1971)
- Stuhl, F., and Niki, H., "Pulsed Vacuum-uv Photochemical Study of Reactions of OH with H<sub>2</sub>, D<sub>2</sub>, and CO Using a Resonance-Fluorescent Detection Method," J. Chem. Phys. 57, 3671-3677 (1972)
- Takacs, G. A., and Glass, G. P., "Reactions of Hydrogen Atoms and Hydroxyl Radicals with Hydrogen Bromide," J. Phys. Chem. 77, 1060-1064 (1973)
- Takacs, G. A., and Glass, G. P., "Reactions of Hydroxyl Radicals with Some Hydrogen Halides," J. Phys. Chem. 77, 1948-1951 (1973)
- Takacs, G. A., and Glass, G. P., "Reaction of Atomic Oxygen with Hydrogen Bromide," J. Phys. Chem. 77, 1182-1186 (1973)
- Taylor, R. L., "High Temperature Reactions of Air" pages 5-154 to 5-157 in "The Natural Stratosphere of 1974" Climatic Impact Assessment Program Monograph 1, DOT-TST-75-51, Department of Transportation, Washington D.C., September 1975
- Thrush, B. A., Letter to D. Garvin (Sept. 1973) reevaluating work in his laboratory
- Toby, F. S., and Toby, S., "Reaction between Ozone and Allene in the Gas Phase," Int. J. Chem. Kinet. 6, 417-428 (1974)
- Toby, F. S., and Toby, S., "The Reaction of Ozone with 1,3-Butadiene and with Allene," Int. J. Chem. Kinet., Symp. No 1, 197-204 (1975)
- Toby, F. S., Toby, S., and O'Neal, H. E., "The Kinetics of the Gas-Phase Reaction between Ozone and Alkenes," Int. J. Chem. Kinet. 8, 25-35 (1976)
- Trainor, D. W., and von Rosenberg, C. W., Jr., "Flash Photolysis Study of the Gas Phase Recombination of Hydroxyl Radicals," J. Chem. Phys. 61, 1010-1015 (1974)



- Trainor, D. W., and von Rosenberg, C. W., Jr., "Energy Partitioning in the Products of Elementary Reactions Involving OH-Radicals," Symp. Combust. 15th (Combustion Institute, Pittsburgh, 1975) 755-764
- Tsang, W., "Comparison between Experimental and Calculated Rate Constants for Dissociation and Combination Reactions Involving Small Polyatomic Molecules," Int. J. Chem. Kinet. 5, 947-963 (1973)
- Vardanyan, I. A., Dangyan, T. M., and Sachyan, G. A., "Rate Constants for the Reaction  $\text{H}\ddot{\text{O}}_2 + \text{CO} \rightarrow \text{CO}_2 + \text{OH}$ ," Dokl. Phys. Chem. 205, 632-634 (1972); tr. of Dokl. Akad. Nauk SSSR 205, 619-621 (1972)
- van den Bergh, H. E., and Callear, A. B., "Spectroscopic Measurement of the Rate of the Gas-Phase Combination of Methyl Radicals with Nitric Oxide and Oxygen at 295 K," Trans. Faraday Soc. 67, 2017-2024 (1971)
- Vandooren, J., Peeters, J., and Van Tiegelen, P. J., "Rate Constant of the Elementary Reaction of Carbon Monoxide with Hydroxyl Radical," Symp. Combust. 15th (Combustion Institute, Pittsburgh, 1975) 745-752
- Volman, D. H., and Gorse, R. A., "Rate Constant for the Reaction of  $\text{H}\ddot{\text{O}}_2$  with Carbon Monoxide," J. Phys. Chem. 76, 3301-3302 (1972)
- von Rosenberg, C. W., Jr., and Trainor, D. W., "Vibrational Excitation of Ozone Formed by Recombination," J. Chem. Phys. 61, 2442-2456 (1974)
- von Rosenberg, C. W., Jr., and Trainor, D. W., "Excitation of Ozone formed by Recombination. II," J. Chem. Phys. 63, 5348-5353 (1975)
- Wagner, H. Gg., Welzbacher, U., and Zellner, R., "Rate Measurements for the Reaction  $\text{H} + \text{Cl}_2 \rightarrow \text{HCl} + \text{Cl}$  by Lyman- $\alpha$  Fluorescence," Ber. Bunsenges. Phys. Chem. 80, 902-908 (1976)
- Wagner, H. Gg., Welzbacher, U., and Zellner, R., "Rate Measurements for the Reactions  $\text{H} + \text{NO}_2 \rightarrow \text{OH} + \text{NO}$  and  $\text{H} + \text{NOCl} \rightarrow \text{HCl} + \text{NO}$  by Lyman- $\alpha$  Fluorescence," Ber. Bunsenges. Phys. Chem. 80, 1023-1027 (1976)
- Walker, R. W., Ph.D., thesis, Univ. of London, Queen Mary College, 1973
- Walker, R. W., Comments on "Gas-Phase Oxidation of Butene-2: The Role of Acetaldehyde in the Reaction," by Ray, D. J. M., Diaz, R. R., and Wassington, D. J., Symp. Combust. 14th (Combustion Institute, Pittsburgh, 1973) 265-266
- Walker, R. W., "Rate Constants for Reactions in Gas-Phase Hydrocarbon Oxidation," Chapter 7 in "Gas Kinetics and Energy Transfer - Volume 2" A Specialist Periodical Report, The Chemical Society, London (1977)
- Washida, N., and Bayes, K. D., "The Reactions of Methyl Radicals with Atomic and Molecular Oxygen," Int. J. Chem. Kinet. 8, 777-794 (1976)
- Washida, N., Martinez, R. I., and Bayes, K. D., "The Oxidation of Formyl Radicals," Z. Naturforsch. A 29, 251-255 (1974)
- Watson, R. T., "Rate Constants for Reactions of  $\text{Cl}\ddot{\text{O}}_x$  of Atmospheric Interest," J. Phys. Chem. Ref. Data 6, 871-918 (1977)
- Watson, R., Machado, G., Fischer, S., and Davis, D. D., "A Temperature Dependence Kinetics Study of the Reaction of  $\text{Cl}(\text{}^2\text{P}_{3/2})$  with  $\text{O}_3$ ,  $\text{CH}_4$  and  $\text{H}_2\ddot{\text{O}}_2$ ," J. Chem. Phys. 65, 2126-2138 (1976)
- Watson, R. T., Machado, G., Conaway, B., Wagner, S., and Davis, D. D., "A Temperature Dependent Kinetics Study of the Reaction of OH with  $\text{CH}_2\text{ClF}$ ,  $\text{CHCl}_2\text{F}$ ,  $\text{CHClF}_2$ ,  $\text{CH}_3\text{CCl}_3$ ,  $\text{CH}_3\text{CF}_2\text{Cl}$ , and  $\text{CF}_2\text{ClCFCl}_2$ ," J. Phys. Chem. 81, 256-262 (1977)
- Weaver, J., Meagher, J., Shortridge, R., and Heicklen, J., "The Oxidation of Acetyl Radicals," J. Photochem. 4, 341-360 (1975)



- Wei, C. N., and Timmons, R. B., "ESR Study of the Kinetics of the Reactions of  $\Theta(^3P)$  Atoms with  $CS_2$  and  $\Theta CS$ ," J. Chem. Phys. **62**, 3240-3245 (1975)
- Westenberg, A. A., and deHaas, N., "Steady-State Intermediate Concentrations and Rate Constants. Some  $H\Theta_2$  Results," J. Phys. Chem. **76**, 1586-1593 (1972a)
- Westenberg, A. A., and deHaas, N., "Measurement of the Rate Constant for  $H + H_2CO \rightarrow H_2 + HCO$  at 297-652°K," J. Phys. Chem. **76**, 2213-2214 (1972b)
- Westenberg, A. A., and deHaas, N., "Rates of  $C\Theta + \Theta H$  and  $H_2 + \Theta H$  Over an Extended Temperature Range," J. Chem. Phys. **58**, 4061-4065 (1973a)
- Westenberg, A. A., and deHaas, N., "Rate of the Reaction  $\Theta H + \Theta H \rightarrow H_2\Theta + \Theta$ ," J. Chem. Phys. **58**, 4066-4071 (1973b)
- Westenberg, A. A., and deHaas, N., "Rates of  $H + CH_3X$  Reactions," J. Chem. Phys. **62**, 3321-3325 (1975a)
- Westenberg, A. A., and deHaas, N., "Reaction Rates of  $\Theta + CH_3Br$  and  $\Theta + CH_3Cl$ ," J. Chem. Phys. **62**, 4477-4479 (1975b)
- Westenberg, A. A., and deHaas, N., "Rate of the  $\Theta + S\Theta_3$  Reaction," J. Chem. Phys. **62**, 725-730 (1975c)
- Westenberg, A. A., and deHaas, N., "Rate of the Reaction of  $\Theta + S\Theta_2 + M \rightarrow S\Theta_3 + M$ ," J. Chem. Phys. **63**, 5411-5415 (1975d)
- Westenberg, A. A., Roscoe, J. M., and DeHaas, N., "Rate Measurements on  $N + \Theta_2(^1\Delta_g) \rightarrow N\Theta + \Theta$  and  $H + \Theta_2(^1\Delta_g) \rightarrow \Theta H + \Theta$ ," Chem. Phys. Lett. **7**, 597-599 (1970)
- Whytock, D. A., Michael, J. V., and Payne, W. A., "Absolute Rate Constants for  $\Theta + N\Theta + N_2 \rightarrow N\Theta_2 + N_2$  from 217 - 500 K," Chem. Phys. Lett. **42**, 466-471 (1976)
- Whytock, D. A., Lee, J. H., Michael, J. V., Payne, W. A., and Stief, L. J., "Absolute Rate of the Reaction of  $Cl(^2P)$  with Methane from 200-500 K," J. Chem. Phys. **66**, 2690-2695 (1977)
- Whytock, D. A., Timmons, R. B., Lee, J. H., Michael, J. V., Payne, W. A., and Stief, L. J., "Absolute Rate of the Reaction of  $\Theta(^3P)$  with Hydrogen Sulfide Over the Temperature Range 263 to 495 K," J. Chem. Phys. **65**, 2052-2055 (1976)
- Wilson, Wm. E., Jr., "A Critical Review of the Gas-Phase Reaction Kinetics of the Hydroxyl Radical," J. Phys. Chem. Ref. Data **1**, 535-573 (1972)
- Wong, E. L., and Belles, F. E., "Rate Measurements for the Reaction of Hydrogen Chloride and Deuterium Chloride with Atomic Oxygen," Lewis Research Center, NASA, Cleveland, Ohio, NASA Tech. Note, NASA TN-D-6495 (1971); Chem. Abstr. **76**:18326q (1972)
- Wong, W., and Davis, D. D., "A Flash Photolysis-Resonance Fluorescence Study of the Reaction of Atomic Hydrogen with Molecular Oxygen:  $H + \Theta_2 + M \rightarrow H\Theta_2 + M$ ," Int. J. Chem. Kinet. **6**, 401-416 (1974)
- Wu, C. H., Morris, E. D., Jr., and Niki, H., "The Reaction of Nitrogen Dioxide with Ozone," J. Phys. Chem. **77**, 2507-2511 (1973)
- Wyrsh, D., Wendt, H. R., and Hunziker, H. E., "Modulation Kinetic Spectroscopy of  $Hg/H_2/\Theta_2/CO$  Mixtures: Reactions of  $HgH$  and  $H\Theta_2$ ," XI. Int. Symp. on Free Radicals (Berchtesgaden-Konigssee, Sept. 4-7, 1973) Paper No. 36; see also, Ber. Bunsenges. Physik. Chem. **78**, 204 (1974)
- Yaron, M., von Engel, A., and Vidaud, P. H., "The Collisional Quenching of  $\Theta_2(^1\Delta_g)$  by  $N\Theta$  and  $CO_2$ ," Chem Phys. Lett. **37**, 159-161 (1976)
- Zahniser, M. S., Berquist, B. M., and Kaufman, F., "Kinetics of the Reaction  $Cl + CH_4 \rightarrow CH_3 + HCl$  from 200 to 500 K," Int. J. Chem. Kinet. **10**, 15-29 (1978)

- Zahniser, M. S., Chang, J. S., and Kaufman, F., "Chlorine Nitrate: Kinetics of Formation by  $\text{ClO} + \text{NO}_2 + \text{M}$  and of Reaction with  $\text{OH}$ ," J. Chem. Phys. 67, 997-1003 (1977)
- Zahniser, M. S., and Kaufman, F., "Kinetics of the Reactions of  $\text{ClO}$  with  $\text{O}$  and with  $\text{NO}$ ," J. Chem. Phys. 66, 3673-3681 (1977)
- Zahniser, M. S., Kaufman, F., and Anderson, J. G., "Kinetics of the Reaction of  $\text{OH}$  with  $\text{HCl}$ ," Chem. Phys. Lett. 27, 507-510 (1974)
- Zahniser, M. S., Kaufman, F., and Anderson, J. G., "Kinetics of the Reaction  $\text{Cl} + \text{O}_3 \rightarrow \text{ClO} + \text{O}_2$ ," Chem. Phys. Lett. 37, 226-231 (1976)
- Zellner, R., Erler, K., and Field, D., "Kinetics of the Recombination Reaction  $\text{OH} + \text{H} + \text{M} \rightarrow \text{H}_2\text{O} + \text{M}$  at Low Temperatures," Symp. Combust. 16th (Combustion Institute, Pittsburgh, 1977) 939-948
- Zellner, R., and Steinert, W., "Flash Photolysis Study of the Rate of the Reaction  $\text{OH} + \text{CH}_4 \rightarrow \text{CH}_3 + \text{H}_2\text{O}$  over an Extended Temperature Range," Int. J. Chem. Kinet. 8, 397-409 (1976)

## APPENDIX 1.

### Chemical Thermodynamic Properties of Selected Molecular Species

E. S. Domalski, D. Garvin, and D. D. Wagman

Two tables are provided showing chemical thermodynamic properties of selected molecular species of interest in stratospheric chemistry. In these tables there are sufficient data for the usual calculations of heats of reaction and equilibria.

The first table lists enthalpy and Gibbs energy of formation, entropy and heat capacity data. The information was extracted from "Selected Values of Chemical Thermodynamic Properties" (D. D. Wagman, et al, NBS Technical Note 270-3, January 1968) except as indicated in the Notes. The thermodynamic properties tabulated are:

1. enthalpy of formation at 0K and 298.15K,
2. Gibbs energy of formation at 298.15K,
3. enthalpy increment between 0K and 298.15K,
4. entropy at 298.15K,
5. heat capacity at constant pressure at 298.15K.

The values given for the heats of formation are based on experimental thermodynamic and spectroscopic measurements except where otherwise indicated; the values of  $S^\circ$ ,  $C_p^\circ$  and  $H_{298}^\circ - H_0^\circ$  for gaseous species were calculated by standard statistical-mechanical equations, using corrections for vibrational anharmonicities, rotational stretching, and rotation-vibration interactions where these data were available.

The second table presents bond dissociation energies or bond strengths. These usually are the enthalpies of simple bond scission reactions. However, there are some cases in which there is a substantial contribution from rearrangement of electronic structure in the products (reducing the bond energy).

The values given here for some molecules may differ slightly from values adopted by the CODATA Task Group on Key Values for Thermodynamics (see CODATA Bulletin 17, published by the International Council of Scientific Unions Committee on Data for Science and Technology, January 1976). These differences will have no significant effect on the enthalpies or free energies of reaction calculated from the tables.

National Bureau of Standards

CHEMICAL THERMODYNAMIC PROPERTIES OF SELECTED MOLECULAR SPECIES

Washington, D. C.

Enthalpy and Gibbs Energy of Formation; Entropy and Heat Capacity *							
Substance		$\Delta H_f^\circ$	$\Delta G_f^\circ$	$H_{298}^\circ - H_0^\circ$	$S^\circ$	$C_p^\circ$	
Formula and Description		State	Formula Weight	298.15°K (25°C)			
				0°K	kcal/mol	cal/deg mol	
O oxygen, atomic	g	15.9994	58.983	59.553	1.607	38.467	
O( <sup>1</sup> D)	g	15.9994	104.34	104.78	1.481		
O( <sup>1</sup> S)	g	15.9994	155.60	156.04	1.481		
O <sub>2</sub> oxygen, molecular, ( <sup>3</sup> Σ <sub>g</sub> <sup>-</sup> )	g	31.9988	0	0	2.0746	7.016	
O <sub>2</sub> ( <sup>1</sup> Δ)	g	31.9988	22.54	22.54			
O <sub>2</sub> ( <sup>1</sup> Σ)	g	31.9988	37.51	37.51			
O <sub>3</sub> ozone	g	47.9982	34.74	34.1	39.0	57.08	
H hydrogen, atomic	g	1.0080	51.633	52.103	48.588	27.391	
H <sub>2</sub> hydrogen, molecular	g	2.0159	0	0	2.0238	31.208	
OH hydroxyl	g	17.0074	9.25	9.31	8.18	43.890	
HO <sub>2</sub> hydroperoxy 1	g	33.0068	1.2+2	0.5+2	3.4	54.73	
H <sub>2</sub> O water	g	18.0153	-57.102	-57.796	-54.634	45.104	
H <sub>2</sub> O <sub>2</sub> hydrogen peroxide	g	34.0147	-31.08	-32.58	-25.24	55.6	
N nitrogen, atomic	g	14.0067	112.534	112.979	108.883	36.622	
N <sub>2</sub> nitrogen, molecular	g	28.0134	0	0	0	45.77	
NO nitric oxide	g	30.0061	21.45	21.57	20.69	50.347	
NO <sub>2</sub> nitrogen dioxide	g	46.0055	8.60	7.93	12.26	57.35	
NO <sub>3</sub> nitrogen trioxide	g	62.008	18.5	17.0	27.7	60.4	
N <sub>2</sub> O nitrous oxide	g	44.0128	20.435	19.61	24.90	52.52	
N <sub>2</sub> O <sub>4</sub> dinitrogen tetroxide	g	92.0110	4.49	2.19	23.38	72.70	
NISTCOMM-NBS-DC							

USCOMM-NBS-DC



## Enthalpy and Gibbs Energy of Formation; Entropy and Heat Capacity

Substance		Formula Weight	State	Formula and Description	$\Delta H_f^\circ$	$\Delta H_f^\circ$	$\Delta G_f^\circ$	$H_{298}^\circ - H_0^\circ$	$S^\circ$	$C_p^\circ$		
298.15°K (25°C)												
0°K												
					kcal/mol		cal/deg mol					
N <sub>2</sub> O <sub>5</sub> dinitrogen pentoxide					108.0104	g	5.7	2.7	27.5	4.237	85.0	20.2
NH imino (imidogen)					15.0147	g	82.	82.	80.6	2.060	43.29	6.97
NH <sub>2</sub> amino					16.0226	g	46.2	45.5	47.8	2.37	46.51	8.02
NH <sub>3</sub> ammonia					17.0306	g	-9.31	-10.98	-3.93	2.40	46.05	8.38
HNO nitroxyl hydride					31.0142	g	24.5	23.8	26.859	2.387	52.729	8.279
HNO <sub>2</sub> cis, nitrous acid					47.0135	g	-17.12	-18.64	-10.27	2.608	59.43	10.70
trans, nitrous acid					47.0135	g	-17.68	-19.15	-10.82	2.652	59.54	11.01
cis-trans mixture, equil.					47.0135	g	-29.94	-19.0	-11.0		60.7	10.9
HNO <sub>3</sub> nitric acid					63.0129	g	-29.94	-32.28	-17.87	2.815	63.64	12.75
S sulfur, rhombic					32.064	c	0	0	0	1.054	7.60	5.41
					32.064	g	65.7	66.20	56.52	1.591	40.085	5.658
SO monosulfur monoxide					48.0634	g	1.2	1.2	-5.0	2.087	53.02	7.21
SO <sub>2</sub> sulfur dioxide					64.0628	g	-70.33	-70.94	-71.74	2.521	59.30	9.53
SO <sub>3</sub> sulfur trioxide					80.0622	c		-108.63	-88.19		12.5	
					80.0622	liq		-105.41	-88.04		22.85	
					80.0622	g	-93.21	-94.58	-88.69	2.796	61.34	12.11
H <sub>2</sub> SO <sub>4</sub> sulfuric acid					98.0775	c	-194.069					
					98.0775	liq		-194.548	-164.938	6.748	37.501	33.20
S <sub>2</sub> O disulfur monoxide					80.1274	g				2.66	63.8	10.5
HS sulfur monohydride					33.0720	g	33.1	33.3	26.3	2.22	46.73	7.76

USCOMM-NBS-DC

USCOMM-NBS-DC

## Enthalpy and Gibbs Energy of Formation; Entropy and Heat Capacity

Substance		$\Delta H_f^\circ$	$\Delta H_f^\circ$	$\Delta G_f^\circ$	$H_{298}^\circ - H_0^\circ$	$S^\circ$	$C_p^\circ$
Formula and Description		State	Formula Weight	298.15°K (25°C)			
				$\Delta H_f^\circ$	$\Delta G_f^\circ$	$H_{298}^\circ - H_0^\circ$	cal/deg mol
				0°K			
H <sub>2</sub> S	hydrogen sulfide	g	34.0799	-4.232	-8.02	2.379	49.16
C	graphite, Acheson spectroscopic	c	12.0112	0	0	0.251	1.372
CO	carbon monoxide	g	28.0106	-27.199	-32.780	2.0716	47.219
CO <sub>2</sub>	carbon dioxide	g	44.0100	-93.963	-94.254	2.2378	6.959
CH <sub>3</sub>	methyl	g	15.0351	35.6	35.3	2.49	51.06
CH <sub>4</sub>	methane	g	16.0430	-15.970	-12.13	2.388	46.38
HCO	formyl	g	29.0185	8.9	5.36	2.386	44.492
HCHO	formaldehyde	g	30.0265	-25.03	-24.51	2.394	53.68
CH <sub>3</sub> O	methoxy	g	31.0345	5.8	6.4	2.41	52.26
CH <sub>3</sub> O <sub>2</sub>	methylperoxy	g	47.0339	6.7			64.2
CH <sub>3</sub> OOH	methylhydroperoxide	g	48.0347	-30.8	-17.4		15.
CH <sub>3</sub> NO <sub>2</sub>	nitromethane	g	61.0406	-14.546	-1.65	3.083	67.9
CH <sub>3</sub> ONO	methyl nitrite	g	61.0406	-13.5	-1.5	3.36	65.69
CH <sub>3</sub> NO <sub>3</sub>	methyl nitrate	g	77.0400	-25.9	-9.4	3.55	69.7
COCl <sub>2</sub>	phosgene	g	98.9166	-52.195	-49.20	3.067	76.1
							67.74
							13.78

JUSCOMM-NBS-DC

# SELECTED VALUES OF CHEMICAL THERMODYNAMIC PROPERTIES - SERIES 1

National Bureau of Standards

Washington, D. C.

Enthalpy and Gibbs Energy of Formation; Entropy and Heat Capacity									
Substance		Formula Weight	State	$\Delta H_f^\circ$	$\Delta H_f^\circ$	$\Delta G_f^\circ$	$H_{298}^\circ - H_0^\circ$	$S^\circ$	$C_p^\circ$
Formula and Description									
				0°K	298.15°K (25°C)			cal/deg mol	
					kcal/mol				
$C_2H_2$	acetylene	26.0382	g	54.324	54.19	50.00	2.392	48.00	10.50
$C_2H_4$	ethylene	28.0542	g	14.515	12.49	16.28	2.525	52.45	10.41
$C_2H_5$	ethyl radical	29.0627	g		25.7	32.0		59.6	10.9
$C_2H_6$	ethane	30.0701	g	-16.323	-20.04	-7.66	2.856	54.85	12.58
$C_3H_6$	propylene	42.0816	g	8.467	4.88	15.00	3.237	63.80	15.27
$C_3H_8$	propane	44.0976	g	-19.684	-25.02	-5.81	3.512	64.51	17.57
$C_4H_{10}$	n-butane	58.1248	g	-23.552	-30.03	-3.97	4.645	74.12	23.29
$C_4H_{10}$	isobutane	58.1248	g	-25.223	-32.07	-4.91	4.276	70.42	23.14
$CH_2CO$	ketene	42.0376	g	-10.66	-11.4	-11.6	2.819	59.16	12.37
$CH_3CO$	acetyl radical	43.0456	g		-5.8	-3.0		64.5	12.7
$CH_3CHO$	acetaldehyde	44.0536	g	-37.14	-39.72	-30.31	3.01	59.8	13.7
$CH_3COCl$	acetyl chloride	78.4986	g	-57.93	-60.07	-51.08	3.529	70.56	16.25
Cl	chlorine, atomic	35.453	g	28.59	28.992	25.173	1.499	39.457	5.220
ClO	chlorine monoxide	51.4524	g	24.33	24.47	23.68	2.278	53.78	8.23
ClO <sub>2</sub>	chlorine dioxide (sym.)	67.4518	g	25.09	24.5	28.8	2.580	61.36	10.03

USCOMM-NBS-DC

USCOMM-NBS-DC

# SELECTED VALUES OF CHEMICAL THERMODYNAMIC PROPERTIES - SERIES I

National Bureau of Standards

Washington, D. C.

Enthalpy and Gibbs Energy of Formation; Entropy and Heat Capacity						
Substance		$\Delta H_f^\circ$	$\Delta H_f^\circ$	$\Delta G_f^\circ$	$H_{298}^\circ - H_0^\circ$	$C_p^\circ$
Formula and Description	State	Formula Weight	0°K	298.15°K (25°C)		
			kcal/mol			
			cal/deg mol			
ClO <sub>2</sub> chlorine dioxide (unsym.)	g	67.4518	21.7	21.3	25.1	2.78
ClO <sub>3</sub> chlorine trioxide	g	83.4512		37		
HCl hydrogen chloride	g	36.4610	-22.020	-22.062	-22.777	2.066
HOCl hypochlorous acid	g	52.4604	-18.0	-18.7	-15.7	2.440
NOCl nitrosyl chloride	g	65.4591	12.81	12.36	15.77	2.716
NO <sub>2</sub> Cl nitryl chloride	g	81.4585	4.29	3.0	13.0	2.915
NO <sub>3</sub> Cl chlorine nitrate	g	97.458		6.28		
Cl <sub>2</sub> chlorine, molecular	g	70.906	0	0	0	2.193
Cl <sub>2</sub> O dichlorine monoxide	g	86.9054	19.62	19.20	23.3	2.806
Cl <sub>2</sub> O <sub>4</sub> chlorine perchlorate	g	134.9036				4.16
CF <sub>4</sub> carbon tetrafluoride	g	88.0048	-221.61	-223.04	-212.37	3.043
CClF <sub>3</sub> chlorotrifluoromethane	g	104.4594	-168.0	-169.2	-159.5	3.294
CF <sub>2</sub> Cl <sub>2</sub> dichlorodifluoromethane	g	120.9140	-116.5	-117.5	-108.2	3.553
CCl <sub>3</sub> F trichlorofluoromethane	g	137.3686	-68.24	-69.0	-59.6	3.838
CCl <sub>4</sub> carbon tetrachloride	g	153.8232	-22.42	-22.94	-12.83	4.120

ITS/COMM-NBS-DC

USCOMM-NBS-DC



# SELECTED VALUES OF CHEMICAL THERMODYNAMIC PROPERTIES - SERIES I

National Bureau of Standards

Washington, D. C.

Enthalpy and Gibbs Energy of Formation; Entropy and Heat Capacity									
Table	Substance		$\Delta H_f^\circ$	$\Delta H_f^\circ$	$\Delta G_f^\circ$	$H_{298}^\circ - H_0^\circ$	$S^\circ$	$C_p^\circ$	
	Formula and Description	State	Formula Weight	298.15°K (25°C)					
				kcal/mol					
				cal/deg mol					
	CH <sub>3</sub> Cl methyl chloride	g	50.4881	-17.69	-19.59	-13.98	2.489	55.99	9.73
	CH <sub>2</sub> Cl <sub>2</sub> methylene dichloride	g	84.9331	-21.19	-22.83	-16.49	2.830	64.57	12.16
	CHCl <sub>3</sub> chloroform	g	119.3781	-23.49	-24.66	-16.83	3.383	70.63	15.63
	CF <sub>3</sub> trifluoromethyl radical	g	69.0064	-111.7	-112.4	-109.21	2.754	63.33	11.90
	CCl <sub>3</sub> trichloromethyl radical	g	118.3702	19.15	19.	22.09	3.392	70.92	15.21
	C <sub>2</sub> Cl <sub>4</sub> tetrachloroethylene	g	165.8343	-2.83	-2.97	5.15	4.759	82.05	22.84
	C <sub>2</sub> HCl <sub>3</sub> trichloroethylene	g	131.3893	-1.032	-1.86	4.31	3.975	77.6	19.18
	CH <sub>3</sub> CCl <sub>3</sub> 1,1,1-trichloroethane (methyl chloroform)	g	133.4052	-34.65	-34.01	-18.21	4.30	76.49	22.07

USCOMM-NBS-DC

USCOMM-NBS-DC

## Notes

1. Hampson, R. F. et al, J. Phys. Chem. Ref. Data 2, 267-312 (1973)  
[Data sheet on  $O_3 + h\nu$  (u.v.)] p. 303.
2. Contributions of the higher electronic states of molecular oxygen to the enthalpy, entropy, and heat capacity at 298 K are insignificant.
3. Stull, D. R. and Prophet, H. "JANAF Thermochemical Tables" 2nd. ed. NSRDS-NBS 37 (1971).
4. Tsang, W. Estimated from kinetic data on the decomposition of  $N_2H_4$ . Chapter 12, NBS Report 10904, 1 July 1972.
5. NBS Report 10904, 1 July 1972, pp. 239-307.
6. R. A. Fletcher and G. Pilcher, Trans Faraday Soc. 66, 794-799 (1970) provide more recent data for  $\Delta H_f^\circ$ , 298.15 K.
7. Data obtained from Benson and O'Neal ("Kinetic Data on Gas Phase Unimolecular Reactions", NSRDS-NBS-21, Feb 1970, page 55).
8. Heat formation value from S. W. Benson "Thermochemical Kinetics" Second Edition, John Wiley and Sons (1976).
9. Values for  $\Delta H_f^\circ$ ,  $S^\circ$ , and  $C_p^\circ$  at 298.15 K were calculated using the group additivity method described by Benson and O'Neal ("Kinetic Data on Gas Phase Unimolecular Reactions", NSRDS-NBS-21, Feb. 1970, pp. 40-42).

10. The value for  $\Delta H_f^\circ [\text{CH}_3\text{ONO}(\text{g})] = -16.5 \text{ kcal mol}^{-1}$  was derived from the combustion data of Geiseler and Thierfelder (Z. physik. Chem., N. F., 29, 248-257 (1961)), and the equilibrium data of Leermakers and Ramsperger (J. Am. Chem. Soc. 54, 1837-1845 (1932)) for:  $\text{CH}_3\text{OH}(\text{g}) + \text{NOCl}(\text{g}) = \text{CH}_3\text{ONO}(\text{g}) + \text{HCl}(\text{g})$ .
11. The value for  $\Delta H_f^\circ (\text{CH}_3\text{ONO}_2(\text{g})) = -29.8 \text{ kcal mol}^{-1}$  was derived from the work of Ray and Ogg (J. Phys. Chem. 63, 1522-1523 (1959)) on the enthalpy of reaction of  $\text{N}_2\text{O}_5$  with  $\text{CH}_3\text{ONO}$ .
12. We have chosen  $S^\circ [\text{CH}_3\text{ONO}_2(\text{g})] = 76.1 \text{ cal mol}^{-1}\text{K}^{-1}$  based upon low temperature thermal data of Gray and Smith (J. Chem. Soc. 1953, 2380-2385). Calculations from microwave studies by Dixon and Wilson (J. Chem. Phys. 35, 191-198 (1961)) give  $S^\circ = 72.15 \text{ cal mol}^{-1}\text{K}^{-1}$ .
13. Davies, J. V., and Pritchard, H. O., J. Chem. Thermodynamics 4, 23-29 (1972) provide more recent  $\Delta H_f^\circ$  data.
14. Data on  $\Delta H_f^\circ$ ,  $S^\circ$ ,  $C_p^\circ$  obtained from S. W. Benson and H. E. O'Neal ("Kinetic Data on Gas Phase Unimolecular Reactions", NSRDS-NBS-21, Feb. 1970, page 567).
15. Rossini, F. D., Pitzer, K. S., Arnett, R. L., Braun, R. M. and Pimentel, G. C., "Selected Values of Physical and Thermodynamic Properties of Hydrocarbons and Related Compounds", API Research Project 44, Carnegie Press, Pittsburgh, 1953.

16.  $\Delta H_f^\circ$  data from Pittam and Pilcher (J. Chem. Soc., Faraday Trans. I, 68, 2224-2229 (1972)).
17.  $\Delta H_f^\circ$  data from R. L. Nuttall, A. H. Laufer and M. V. Kilday., J. Chem. Thermodynamics 3, 167-174 (1971).
18. Devore, J. A., and O'Neal, H. E., J. Phys. Chem. 73, 2644-2648 (1969).
19. Data on  $S^\circ$  and  $C_p^\circ$  were obtained from Note (14), page 589; data on  $\Delta H_f^\circ$  came from Note (18).
20. Spectroscopic studies of N. Basco and R. D. Morse, J. Molec. Spectros. 45, 35 (1973); R. A. Durie and D. A. Ramsay, Can. J. Phys. 36, 35 (1958); P. A. G. O'Hare and A. C. Wahl, J. Chem. Phys. 54, 3770 (1971); L. Andrews and J. I. Raymond, J. Chem. Phys. 55, 3087 (1971); D. R. Johnson and F. X. Powell, Bull. Am. Phys. Soc. 13, 594 (1968); A. Carrington, P. N. Dyer, and D. H. Levy, J. Chem. Phys. 47, 1756 (1967) were examined and used to calculate the thermodynamic properties.
21. Arkell, A., and Schwager, I., J. Am. Chem. Soc. 89, 5999 (1967); Johnston, H., Morris, E. D., and Van den Bogaerde, J., J. Am. Chem. Soc. 91, 7712 (1969), Clyne, M. A. A. and White, I. F., Trans. Faraday Soc. 67, 2068 (1971).
22. Studies by D. J. Gardiner, J. Mol. Spectroscopy 38, 476 (1971) and G. E. Herberich, R. H. Jackson, and D. J. Miller, J. Chem. Soc. (A) 1966, 336 were examined and the data used to calculate the thermodynamic properties.



23. The spectroscopic data of K. O. Christe, C. J. Schack and E. C. Curtis, *Inorg. Chem.* 10, 1589 (1971) were used.
24. Thermal Constants of Substances (Termichesky Konstanty Veshchestv), Vol. IV, V. P. Glushko, Editor, Moscow (1970).
25. The combustion data of A. T. Hu and G. C. Sinke, *J. Chem. Thermodynamics* 1, 507-513 (1969) were used to derive  $\Delta H_f^\circ$  298.15 K values for  $\text{CCl}_4$ ,  $\text{CHCl}_3$ , and  $\text{CH}_2\text{Cl}_2$  in the liquid state.
26. The vaporization data of D. L. Hildenbrand and R. A. McDonald, *J. Phys. Chem.* 63, 1521-1522 (1959) were used.
27. The combustion data of R. A. Fletcher and G. Pilcher, *Trans. Faraday Soc.* 67, 3191-3201 (1971) were used to calculate  $\Delta H_f^\circ [\text{CH}_3\text{Cl(g)}]$ .
28. J. Chao, A. S. Rodgers, R. C. Wilhoit, and B. J. Zwolinski, *J. Phys. & Chem. Ref. Data* 3, 141-162 (1974) provide a recent evaluation of the thermodynamic properties of  $\text{CH}_3\text{CCl}_3$ .
29. Based on data in H. Okabe and M. Lenzi, *J. Chem. Phys.* 47, 5241 (1967) and in J. Masanet, A. Gilles and C. Vermeil, *J. Photochemistry* 3, 417 (1974/75).
30. Parker, V. B., Wagman, D. D., and Garvin, D., NBSIR 75-968. Selected thermochemical data compatible with the CODATA recommendations, January 1976.
31. Value for heat of formation derived from equilibrium studies reported by H. Imagawa, *Denki Kagaku* 19, 271 (1951).

32. JANAF Thermochemical Tables Supplement No. 48, June 1977.
33. Heat of formation value based on data in H. D. Knauth, H. Martin and W. Stockmann, Z. Naturforsch. 29A, 200 (1974).
34. Wagman, D.D. Private communication.

Bond Dissociation Energies,  $D_0^\circ$  at 0 K  
Calculated from heats of formation in the accompanying table

Reaction	kJ/mol	kcal/mol	Note
$O_2 \rightarrow 2O$	493.59±0.4	117.97±0.1	
$O_3 \rightarrow O_2 + O$	101.4±2.1	24.24±0.5	
$H_2 \rightarrow 2H$	432.00±0.04	103.25±0.01	
$HO \rightarrow H + O$	423.8±2.1	101.3±0.5	
$HO_2 \rightarrow HO + O$	230±8	67±2	
$\rightarrow H + O_2$	210±8	50±2	
$H_2O \rightarrow HO + H$	493.7±0.8	118.0±0.2	
$H_2O_2 \rightarrow 2HO$	207.1±2.1	49.5±0.5	
$H_2O_2 \rightarrow H + HO_2$	350±8	84±2	
$N_2 \rightarrow N + N$	941.69±0.04	225.07±0.01	
$NO \rightarrow N + O$	627.9	150.1	
$NO_2 \rightarrow NO + O$	300.5	71.83	
$NO_3 \rightarrow NO_2 + O$	205±21	49±5	
$N_2O \rightarrow N + NO$	474.9±4	113.5±1	
$\rightarrow N_2 + O$	161.3	38.58	
$N_2O_3 \rightarrow NO + NO_2$	35.1±0.8	8.4±0.2	
$N_2O_4 \rightarrow NO_2 + NO_2$	53.2	12.7	
$N_2O_5 \rightarrow NO_2 + NO_3$	89.5	21.4	
$HNO \rightarrow H + NO$	203	48.6	
$\rightarrow NH + O$	487	116.5	
$trans-HNO_2 \rightarrow HO + NO$	202	48.4	
$HNO_3 \rightarrow HO + NO_2$	200	47.8	
$NH \rightarrow H + N$	343	82	

Bond Dissociation Energies,  $D_0^0$  at 0 K  
Calculated from heats of formation in the accompanying table

Reaction	kJ/mol	kcal/mol	Note
$\text{NH}_2 \rightarrow \text{H} + \text{HN}$	366	87.4	
$\text{NH}_3 \rightarrow \text{NH}_2 + \text{H}$	448	107	
$\text{CO} \rightarrow \text{C} + \text{O}$	1071.94±0.4	256.2±0.1	
$\text{CO}_2 \rightarrow \text{CO} + \text{O}$	525.9±0.4	125.7±0.1	
$\text{CHO} \rightarrow \text{CO} + \text{H}$	70	16.7	(1)
$\text{CH}_2\text{O} \rightarrow \text{CHO} + \text{H}$	364	87	(1)
$\text{CH}_3\text{O} \rightarrow \text{CH}_3 + \text{O}$	378	90.4	(1)
$\quad \rightarrow \text{H} + \text{H}_2\text{CO}$	93	22	(1)
$\text{CH}_3\text{O}_2 \rightarrow \text{CH}_3\text{O} + \text{O}$	237	56.8	(1)
$\quad \rightarrow \text{CH}_3 + \text{O}_2$	118	28.1	(1)
$\text{CH}_4 \rightarrow \text{CH}_3 + \text{H}$	432	103.2	
$\text{CH}_3\text{OOH} \rightarrow \text{CH}_3\text{O} + \text{HO}$	184.	44.0	(1)
$\quad \rightarrow \text{CH}_3 + \text{HO}_2$	276±8	66±2	(1)
$\text{CH}_3\text{ONO} \rightarrow \text{CH}_3 + \text{NO}_2$	248	59.2	(1)
$\text{CH}_3\text{NO}_2 \rightarrow \text{CH}_3 + \text{NO}_2$	254	60.6	(1)
$\text{CH}_3\text{ONO}_2 \rightarrow \text{CH}_3\text{O} + \text{NO}_2$	174	41.6	(1)
$\quad \rightarrow \text{CH}_3 + \text{NO}_3$	341	81.6	(1)
$\text{SO} \rightarrow \text{S} + \text{O}$	515.4±8	123.2±2	
$\text{SO}_2 \rightarrow \text{SO} + \text{O}$	547.3±8	130.8±2.0	
$\text{SO}_3 \rightarrow \text{SO}_2 + \text{O}$	342.7±4	81.9±1	

Note 1. Calculated for T = 298K



APPENDIX 2. CONVERSION TABLES  
EQUIVALENT SECOND ORDER RATE CONSTANTS

A \ B	$\text{cm}^3 \text{mol}^{-1} \text{s}^{-1}$	$\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$	$\text{m}^3 \text{mol}^{-1} \text{s}^{-1}$	$\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$	$(\text{mm Hg})^{-1} \text{s}^{-1}$	$\text{atm}^{-1} \text{s}^{-1}$	$\text{ppm}^{-1} \text{min}^{-1}$	$\text{m}^2 \text{KN}^{-1} \text{s}^{-1}$
$1 \text{ cm}^3 \text{mol}^{-1} \text{s}^{-1} =$	1	$10^{-3}$	$10^{-6}$	$1.66 \times 10^{-24}$	$1.604 \times 10^{-5} \text{T}^{-1}$	$1.219 \times 10^{-2} \text{T}^{-1}$	$2.453 \times 10^{-9}$	$1.203 \times 10^{-4} \text{T}^{-1}$
$1 \text{ dm}^3 \text{mol}^{-1} \text{s}^{-1} =$	$10^3$	1	$10^{-3}$	$1.66 \times 10^{-21}$	$1.604 \times 10^{-2} \text{T}^{-1}$	$12.19 \text{T}^{-1}$	$2.453 \times 10^{-6}$	$1.203 \times 10^{-1} \text{T}^{-1}$
$1 \text{ m}^3 \text{mol}^{-1} \text{s}^{-1} =$	$10^6$	$10^3$	1	$1.66 \times 10^{-18}$	$16.04 \text{T}^{-1}$	$1.219 \times 10^4 \text{T}^{-1}$	$2.453 \times 10^{-3}$	$120.3 \text{T}^{-1}$
$1 \text{ cm}^3 \text{molecule}^{-1} \text{s}^{-1} =$	$6.023 \times 10^{23}$	$6.023 \times 10^{20}$	$6.023 \times 10^{17}$	1	$9.658 \times 10^{18} \text{T}^{-1}$	$7.34 \times 10^{21} \text{T}^{-1}$	$1.478 \times 10^{15}$	$7.244 \times 10^{19} \text{T}^{-1}$
$1 (\text{mm Hg})^{-1} \text{s}^{-1} =$	$6.236 \times 10^4 \text{T}$	$62.36 \text{T}$	$6.236 \times 10^{-2} \text{T}$	$1.035 \times 10^{-19} \text{T}$	1	760	$4.56 \times 10^{-2}$	7.500
$1 \text{ atm}^{-1} \text{s}^{-1} =$	82.06 T	$8.206 \times 10^{-2} \text{T}$	$8.206 \times 10^{-5} \text{T}$	$1.362 \times 10^{-22} \text{T}$	$1.316 \times 10^{-3}$	1	$6 \times 10^{-5}$	$9.869 \times 10^{-3}$
$1 \text{ ppm}^{-1} \text{min}^{-1} =$ at 298K, 1 atm. total pressure	$4.077 \times 10^8$	$4.077 \times 10^5$	407.7	$6.76 \times 10^{-16}$	21.93	$1.667 \times 10^4$	1	164.5
$1 \text{ m}^2 \text{KN}^{-1} \text{s}^{-1} =$	8314 T	$8.314 \text{T}$	$8.314 \times 10^{-3} \text{T}$	$1.38 \times 10^{-20} \text{T}$	0.1333	101.325	$6.079 \times 10^{-3}$	1

To convert a rate constant from one set of units A to a new set B find the conversion factor for the row A under Column B and multiply the old value by it, e.g. to convert  $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$  to  $\text{m}^3 \text{mol}^{-1} \text{s}^{-1}$  multiply by  $6.023 \times 10^{17}$ .

Table adapted from Evaluated Kinetic Data for High Temperature Reactions, Volume 1: Homogeneous Gas Phase Reactions of the  $\text{H}_2\text{-O}_2$  System, Butterworths, London, 1972.

# EQUIVALENT THIRD ORDER RATE CONSTANTS

A \ B	$6 \text{ cm mol}^{-2} \text{ s}^{-1}$	$6 \text{ dm mol}^{-2} \text{ s}^{-1}$	$6 \text{ mol}^{-2} \text{ s}^{-1}$	$6 \text{ cm molecule}^{-2} \text{ s}^{-1}$	$(\text{mm Hg})^{-2} \text{ s}^{-1}$	$\text{atm}^{-2} \text{ s}^{-1}$	$\text{ppm}^{-2} \text{ min}^{-1}$	$4 \text{ m KN}^{-2} \text{ s}^{-1}$
$1 \text{ cm mol}^{-2} \text{ s}^{-1} =$	1	$10^{-6}$	$10^{-12}$	$2.76 \times 10^{-48}$	$2.57 \times 10^{-10} \text{ T}^{-2}$	$1.48 \times 10^{-4} \text{ T}^{-2}$	$1.003 \times 10^{-19}$	$1.447 \times 10^{-8} \text{ T}^{-2}$
$1 \text{ dm mol}^{-2} \text{ s}^{-1} =$	$10^6$	1	$10^{-6}$	$2.76 \times 10^{-42}$	$2.57 \times 10^{-4} \text{ T}^{-2}$	$148 \text{ T}^{-2}$	$1.003 \times 10^{-13}$	$1.447 \times 10^{-2} \text{ T}^{-2}$
$1 \text{ m mol}^{-2} \text{ s}^{-1} =$	$10^{12}$	$10^6$	1	$2.76 \times 10^{-36}$	$257 \text{ T}^{-2}$	$1.48 \times 10^8 \text{ T}^{-2}$	$1.003 \times 10^{-7}$	$1.447 \times 10^4 \text{ T}^{-2}$
$1 \text{ cm molecule}^{-2} \text{ s}^{-1} =$	$3.628 \times 10^{47}$	$3.628 \times 10^{41}$	$3.628 \times 10^{35}$	1	$9.328 \times 10^{37} \text{ T}^{-2}$	$5.388 \times 10^{43} \text{ T}^{-2}$	$3.64 \times 10^{28}$	$5.248 \times 10^{39} \text{ T}^{-2}$
$1 (\text{mm Hg})^{-2} \text{ s}^{-1} =$	$3.89 \times 10^{9.2} \text{ T}^{-2}$	$3.89 \times 10^{3.2} \text{ T}^{-2}$	$3.89 \times 10^{-3.2} \text{ T}^{-2}$	$1.07 \times 10^{-38.2} \text{ T}^{-2}$	1	$5.776 \times 10^5$	$3.46 \times 10^{-5}$	56.25
$1 \text{ atm}^{-2} \text{ s}^{-1} =$	$6.733 \times 10^{3.2} \text{ T}^{-2}$	$6.733 \times 10^{-3.2} \text{ T}^{-2}$	$6.733 \times 10^{-9.2} \text{ T}^{-2}$	$1.86 \times 10^{-44.2} \text{ T}^{-2}$	$1.73 \times 10^{-6}$	1	$6 \times 10^{-11}$	$9.74 \times 10^{-5}$
$1 \text{ ppm}^{-2} \text{ min}^{-1} =$ at 298K, 1 atm. total pressure	$9.97 \times 10^{18}$	$9.97 \times 10^{12}$	$9.97 \times 10^6$	$2.75 \times 10^{-29}$	$2.89 \times 10^4$	$1.667 \times 10^{10}$	1	$1.623 \times 10^6$
$1 \text{ m KN}^{-2} \text{ s}^{-1} =$	$6.91 \times 10^{7.2} \text{ T}^{-2}$	$69.1 \text{ T}^{-2}$	$6.91 \times 10^{-5.2} \text{ T}^{-2}$	$1.904 \times 10^{-40.2} \text{ T}^{-2}$	0.0178	$1.027 \times 10^4$	$6.16 \times 10^{-7}$	1

See note to Table for Second Order Rate Constants

CONVERSION FACTORS FOR UNITS OF OPTICAL ABSORPTION COEFFICIENTS

B A	(cross section $\sigma$ ) $\frac{2}{\text{cm molecule}}^{-1}$ base e	(atm at 273) $^{-1}$ cm base e	$\frac{3}{\text{dm mol}}^{-1}$ cm $^{-1}$ base 10	$\frac{2}{\text{cm mol}}^{-1}$ base 10
1 (atm at 298) $^{-1}$ cm $^{-1}$ base e =	$4.06 \times 10^{-20}$	1.09	10.6	$1.06 \times 10^4$
1 (atm at 298) $^{-1}$ cm $^{-1}$ base 10 =	$9.35 \times 10^{-20}$	2.51	24.4	$2.44 \times 10^4$
1 (mm Hg at 298) $^{-1}$ cm $^{-1}$ base 10 =	$7.11 \times 10^{-17}$	$1.91 \times 10^3$	$1.86 \times 10^4$	$1.86 \times 10^7$
1 (atm at 273) $^{-1}$ cm $^{-1}$ base e =	$3.72 \times 10^{-20}$	1	9.73	$9.73 \times 10^3$
1 (atm at 273) $^{-1}$ cm $^{-1}$ base 10 =	$8.57 \times 10^{-20}$	2.303	22.4	$2.24 \times 10^4$
1 dm $^3$ mol $^{-1}$ cm $^{-1}$ base 10 =	$3.82 \times 10^{-21}$	0.103	1	$10^3$
1 cm $^2$ mol $^{-1}$ base 10 =	$3.82 \times 10^{-24}$	$1.03 \times 10^{-4}$	$10^{-3}$	1
1 cm $^2$ molecule $^{-1}$ base e =	1	$2.69 \times 10^{19}$	$2.62 \times 10^{20}$	$2.62 \times 10^{23}$

To convert an absorption coefficient from one set of units A to a new set B, multiply by the value tabulated for row A under column B, e.g. to convert the value of the absorption coefficient expressed in dm $^3$  mol $^{-1}$  cm $^{-1}$  base 10 to (atm at 273) $^{-1}$  cm $^{-1}$  base e, multiply by 0.103.

U.S. DEPT. OF COMM. BIBLIOGRAPHIC DATA SHEET	1. PUBLICATION OR REPORT NO.  NBS SP-513	2. Gov't Accession No.	3. Recipient's Accession No.
4. TITLE AND SUBTITLE  Reaction Rate and Photochemical Data for Atmospheric Chemistry - 1977		5. Publication Date  May 1978	
		6. Performing Organization Code	
7. AUTHOR(S)  Robert F. Hampson, Jr., and David Garvin, Editors		8. Performing Organ. Report No.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS  NATIONAL BUREAU OF STANDARDS DEPARTMENT OF COMMERCE WASHINGTON, D.C. 20234		10. Project/Task/Work Unit No.	
		11. Contract/Grant No.	
12. Sponsoring Organization Name and Complete Address (Street, City, State, ZIP) Department of Transportation, FAA, Washington, DC 20591 N.A.S.A., Washington, DC 20546 Office of Standard Reference Data, NBS, Washington, DC 20234 Office of Environmental Measurements, NBS, Washington DC 20234		13. Type of Report & Period Covered	
		14. Sponsoring Agency Code	
15. SUPPLEMENTARY NOTES  Library of Congress Catalog Card Number: 78-606033			
16. ABSTRACT (A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here.)  A table of data for gas phase chemical reactions and photochemistry of neutral species is presented. Specifically, it gives preferred values for reaction rate constants, photoabsorption cross sections, and quantum yields of primary photochemical processes and also cites recent experimental work (to December 1977). It is intended to provide the basic physical chemical data needed as input data for calculations modelling atmospheric chemistry. An auxiliary table of thermochemical data for the pertinent chemical species is given in the appendix.			
17. KEY WORDS (six to twelve entries; alphabetical order; capitalize only the first letter of the first key word unless a proper name; separated by semicolons) Air pollution; atmospheric chemistry; chemical kinetics; data evaluation; gas phase; photoabsorption cross section; photochemistry; quantum yield; rate constant			
18. AVAILABILITY  <input checked="" type="checkbox"/> Unlimited  <input type="checkbox"/> For Official Distribution. Do Not Release to NTIS  <input checked="" type="checkbox"/> Order From Sup. of Doc., U.S. Government Printing Office Washington, D.C. 20402, SD Stock No. SN003-003-  <input type="checkbox"/> Order From National Technical Information Service (NTIS) Springfield, Virginia 22151		19. SECURITY CLASS (THIS REPORT)  UNCLASSIFIED	21. NO. OF PAGES  111
		20. SECURITY CLASS (THIS PAGE)  UNCLASSIFIED	22. Price





# NBS TECHNICAL PUBLICATIONS

## PERIODICALS

**JOURNAL OF RESEARCH**—The Journal of Research of the National Bureau of Standards reports NBS research and development in those disciplines of the physical and engineering sciences in which the Bureau is active. These include physics, chemistry, engineering, mathematics, and computer sciences. Papers cover a broad range of subjects, with major emphasis on measurement methodology, and the basic technology underlying standardization. Also included from time to time are survey articles on topics closely related to the Bureau's technical and scientific programs. As a special service to subscribers each issue contains complete citations to all recent NBS publications in NBS and non-NBS media. Issued six times a year. Annual subscription: domestic \$17.00; foreign \$21.25. Single copy, \$3.00 domestic; \$3.75 foreign.

Note: The Journal was formerly published in two sections: Section A "Physics and Chemistry" and Section B "Mathematical Sciences."

### DIMENSIONS/NBS

This monthly magazine is published to inform scientists, engineers, businessmen, industry, teachers, students, and consumers of the latest advances in science and technology, with primary emphasis on the work at NBS. The magazine highlights and reviews such issues as energy research, fire protection, building technology, metric conversion, pollution abatement, health and safety, and consumer product performance. In addition, it reports the results of Bureau programs in measurement standards and techniques, properties of matter and materials, engineering standards and services, instrumentation, and automatic data processing.

Annual subscription: Domestic, \$12.50; Foreign \$15.65.

## NONPERIODICALS

**Monographs**—Major contributions to the technical literature on various subjects related to the Bureau's scientific and technical activities.

**Handbooks**—Recommended codes of engineering and industrial practice (including safety codes) developed in cooperation with interested industries, professional organizations, and regulatory bodies.

**Special Publications**—Include proceedings of conferences sponsored by NBS, NBS annual reports, and other special publications appropriate to this grouping such as wall charts, pocket cards, and bibliographies.

**Applied Mathematics Series**—Mathematical tables, manuals, and studies of special interest to physicists, engineers, chemists, biologists, mathematicians, computer programmers, and others engaged in scientific and technical work.

**National Standard Reference Data Series**—Provides quantitative data on the physical and chemical properties of materials, compiled from the world's literature and critically evaluated. Developed under a world-wide program coordinated by NBS. Program under authority of National Standard Data Act (Public Law 90-396).

NOTE: At present the principal publication outlet for these data is the Journal of Physical and Chemical Reference Data (JPCRD) published quarterly for NBS by the American Chemical Society (ACS) and the American Institute of Physics (AIP). Subscriptions, reprints, and supplements available from ACS, 1155 Sixteenth St. N.W., Wash., D.C. 20056.

**Building Science Series**—Disseminates technical information developed at the Bureau on building materials, components, systems, and whole structures. The series presents research results, test methods, and performance criteria related to the structural and environmental functions and the durability and safety characteristics of building elements and systems.

**Technical Notes**—Studies or reports which are complete in themselves but restrictive in their treatment of a subject. Analogous to monographs but not so comprehensive in scope or definitive in treatment of the subject area. Often serve as a vehicle for final reports of work performed at NBS under the sponsorship of other government agencies.

**Voluntary Product Standards**—Developed under procedures published by the Department of Commerce in Part 10, Title 15, of the Code of Federal Regulations. The purpose of the standards is to establish nationally recognized requirements for products, and to provide all concerned interests with a basis for common understanding of the characteristics of the products. NBS administers this program as a supplement to the activities of the private sector standardizing organizations.

**Consumer Information Series**—Practical information, based on NBS research and experience, covering areas of interest to the consumer. Easily understandable language and illustrations provide useful background knowledge for shopping in today's technological marketplace.

*Order above NBS publications from: Superintendent of Documents, Government Printing Office, Washington, D.C. 20402.*

*Order following NBS publications—NBSIR's and FIPS from the National Technical Information Services, Springfield, Va. 22161.*

**Federal Information Processing Standards Publications (FIPS PUB)**—Publications in this series collectively constitute the Federal Information Processing Standards Register. Register serves as the official source of information in the Federal Government regarding standards issued by NBS pursuant to the Federal Property and Administrative Services Act of 1949 as amended, Public Law 89-306 (79 Stat. 1127), and as implemented by Executive Order 11717 (38 FR 12315, dated May 11, 1973) and Part 6 of Title 15 CFR (Code of Federal Regulations).

**NBS Interagency Reports (NBSIR)**—A special series of interim or final reports on work performed by NBS for outside sponsors (both government and non-government). In general, initial distribution is handled by the sponsor; public distribution is by the National Technical Information Services (Springfield, Va. 22161) in paper copy or microfiche form.

## BIBLIOGRAPHIC SUBSCRIPTION SERVICES

The following current-awareness and literature-survey bibliographies are issued periodically by the Bureau:

**Cryogenic Data Center Current Awareness Service.** A literature survey issued biweekly. Annual subscription: Domestic, \$25.00; Foreign, \$30.00.

**Liquified Natural Gas.** A literature survey issued quarterly. Annual subscription: \$20.00.

**Superconducting Devices and Materials.** A literature survey issued quarterly. Annual subscription: \$30.00. Send subscription orders and remittances for the preceding bibliographic services to National Bureau of Standards, Cryogenic Data Center (275.02) Boulder, Colorado 80302.

**U.S. DEPARTMENT OF COMMERCE**  
**National Bureau of Standards**  
Washington, D.C. 20234

OFFICIAL BUSINESS

Penalty for Private Use, \$300

POSTAGE AND FEES PAID  
U.S. DEPARTMENT OF COMMERCE  
COM-215



SPECIAL FOURTH-CLASS RATE  
BOOK

---